

Table 1.1C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 12 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	942	795	508	455	366	332
Marijuana and Hashish	872	775	428	398	295	281
Cocaine	596	519	188	150	130	112
Crack	293	260	120	68	78	49
Heroin	229	202	54	46	34	34
Hallucinogens	557	487	101	131	59	64
LSD	479	433	81	88	43	38
PCP	298	279	28	29	11	10
Inhalants	443	381	88	90	52	67
Nonmedical Use of Any Psychotherapeutic ²	645	582	296	259	209	178
Pain Relievers	490	458	261	218	178	142
Tranquilizers	460	403	155	162	114	99
Stimulants	466	392	130	118	98	77
Methamphetamine	383	296	101	92	69	62
Sedatives	354	309	85	71	61	38
Any Illicit Drug Other Than Marijuana ¹	787	704	341	306	242	209

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.1D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 12 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.43	0.36	0.23	0.20	0.17	0.15
Marijuana and Hashish	0.39	0.35	0.19	0.18	0.13	0.13
Cocaine	0.27	0.23	0.08	0.07	0.06	0.05
Crack	0.13	0.12	0.05	0.03	0.04	0.02
Heroin	0.10	0.09	0.02	0.02	0.02	0.02
Hallucinogens	0.25	0.22	0.05	0.06	0.03	0.03
LSD	0.22	0.19	0.04	0.04	0.02	0.02
PCP	0.13	0.12	0.01	0.01	0.00	0.00
Inhalants	0.20	0.17	0.04	0.04	0.02	0.03
Nonmedical Use of Any Psychotherapeutic ²	0.29	0.26	0.13	0.12	0.09	0.08
Pain Relievers	0.22	0.21	0.12	0.10	0.08	0.06
Tranquilizers	0.21	0.18	0.07	0.07	0.05	0.04
Stimulants	0.21	0.18	0.06	0.05	0.04	0.03
Methamphetamine	0.17	0.13	0.05	0.04	0.03	0.03
Sedatives	0.16	0.14	0.04	0.03	0.03	0.02
Any Illicit Drug Other Than Marijuana ¹	0.36	0.32	0.15	0.14	0.11	0.09

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.1P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 12 or Older: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.1639		0.0845		0.8949	
Marijuana and Hashish	0.4517		0.1936		0.6920	
Cocaine	0.3230		0.0660		0.0520	
Crack	0.0941		0.0180		0.0915	
Heroin	0.3171		0.4925		0.5928	
Hallucinogens	0.2561		0.1092		0.6325	
LSD	0.7023		0.1406		0.1503	
PCP	0.8955		0.6937		0.9211	
Inhalants	0.2871		0.7619		0.5823	
Nonmedical Use of Any Psychotherapeutic ²	0.0230		0.1617		0.6054	
Pain Relievers	0.1832		0.6044		0.5448	
Tranquilizers	0.0952		0.9148		0.4782	
Stimulants	0.0173		0.2428		0.1804	
Methamphetamine	0.1567		0.3812		0.5852	
Sedatives	0.1258		0.8104		0.4365	
Any Illicit Drug Other Than Marijuana ¹	0.2462		0.2589		0.3178	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.2C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	83	86	75	73	53	56
Marijuana and Hashish	74	72	66	62	47	49
Cocaine	29	31	24	27	14	17
Crack	14	13	12	10	4	5
Heroin	13	10	11	8	10	4
Hallucinogens	45	43	36	36	20	22
LSD	35	36	29	28	16	14
PCP	17	18	11	11	6	4
Inhalants	54	49	37	32	18	17
Nonmedical Use of Any Psychotherapeutic ²	58	61	47	47	30	31
Pain Relievers	49	54	41	41	26	27
Tranquilizers	28	29	23	21	13	11
Stimulants	37	35	27	29	13	16
Methamphetamine	20	19	15	15	8	10
Sedatives	16	15	12	11	8	7
Any Illicit Drug Other Than Marijuana ¹	73	74	59	60	38	39

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.2D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.36	0.37	0.32	0.31	0.23	0.24
Marijuana and Hashish	0.32	0.31	0.29	0.27	0.20	0.21
Cocaine	0.13	0.13	0.10	0.12	0.06	0.07
Crack	0.06	0.06	0.05	0.04	0.02	0.02
Heroin	0.06	0.04	0.05	0.03	0.04	0.02
Hallucinogens	0.19	0.18	0.16	0.16	0.08	0.09
LSD	0.15	0.15	0.12	0.12	0.07	0.06
PCP	0.07	0.08	0.05	0.05	0.02	0.02
Inhalants	0.23	0.21	0.16	0.14	0.08	0.07
Nonmedical Use of Any Psychotherapeutic ²	0.25	0.26	0.20	0.20	0.13	0.13
Pain Relievers	0.21	0.23	0.18	0.17	0.11	0.12
Tranquilizers	0.12	0.12	0.10	0.09	0.05	0.05
Stimulants	0.16	0.15	0.12	0.12	0.06	0.07
Methamphetamine	0.09	0.08	0.06	0.06	0.03	0.04
Sedatives	0.07	0.06	0.05	0.05	0.03	0.03
Any Illicit Drug Other Than Marijuana ¹	0.31	0.32	0.26	0.26	0.17	0.17

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.2P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 12 to 17: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.1413		0.0119		0.8229	
Marijuana and Hashish	0.3569		0.0503		0.8917	
Cocaine	0.9117		0.7337		0.7400	
Crack	0.9667		0.8031		0.8349	
Heroin	0.2640		0.2241		0.0257	
Hallucinogens	0.7335		0.8101		0.6370	
LSD	0.3984		0.2542		0.1484	
PCP	0.2710		0.4237		0.2931	
Inhalants	0.4586		0.0783		0.3584	
Nonmedical Use of Any Psychotherapeutic ²	0.9950		0.8173		0.6305	
Pain Relievers	0.4221		0.7913		0.3217	
Tranquilizers	0.6702		0.7823		0.5380	
Stimulants	0.9260		0.1553		0.1748	
Methamphetamine	0.5994		0.3093		0.2335	
Sedatives	0.8534		0.8828		0.4352	
Any Illicit Drug Other Than Marijuana ¹	0.6289		0.3114		0.6965	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.3C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	154	150	137	132	115	105
Marijuana and Hashish	149	150	131	126	109	98
Cocaine	93	83	60	52	35	32
Crack	47	42	26	19	12	9
Heroin	38	33	18	15	9	7
Hallucinogens	118	118	67	71	36	34
LSD	105	99	54	47	26	24
PCP	39	37	16	13	9	6
Inhalants	98	94	47	42	22	19
Nonmedical Use of Any Psychotherapeutic ²	117	102	86	73	52	50
Pain Relievers	106	89	76	65	43	44
Tranquilizers	75	72	47	45	26	26
Stimulants	78	68	51	42	27	22
Methamphetamine	58	54	35	30	19	14
Sedatives	37	33	20	21	11	9
Any Illicit Drug Other Than Marijuana ¹	139	133	104	98	64	62

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.3D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.54	0.52	0.48	0.46	0.40	0.36
Marijuana and Hashish	0.52	0.52	0.46	0.43	0.38	0.34
Cocaine	0.33	0.29	0.21	0.18	0.12	0.11
Crack	0.17	0.15	0.09	0.06	0.04	0.03
Heroin	0.13	0.11	0.06	0.05	0.03	0.03
Hallucinogens	0.42	0.41	0.24	0.25	0.13	0.12
LSD	0.37	0.34	0.19	0.16	0.09	0.08
PCP	0.14	0.13	0.06	0.05	0.03	0.02
Inhalants	0.34	0.32	0.16	0.14	0.08	0.07
Nonmedical Use of Any Psychotherapeutic ²	0.41	0.35	0.30	0.25	0.18	0.17
Pain Relievers	0.37	0.31	0.27	0.22	0.15	0.15
Tranquilizers	0.26	0.25	0.16	0.16	0.09	0.09
Stimulants	0.28	0.23	0.18	0.15	0.10	0.07
Methamphetamine	0.20	0.19	0.12	0.10	0.07	0.05
Sedatives	0.13	0.12	0.07	0.07	0.04	0.03
Any Illicit Drug Other Than Marijuana ¹	0.49	0.46	0.37	0.34	0.22	0.21

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.3P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 18 to 25: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.0542		0.0571		0.3377	
Marijuana and Hashish	0.1309		0.1937		0.2250	
Cocaine	0.0204		0.0022		0.0552	
Crack	0.0476		0.0095		0.0377	
Heroin	0.0424		0.3228		0.3734	
Hallucinogens	0.9899		0.9661		0.8172	
LSD	0.1915		0.0310		0.1911	
PCP	0.5382		0.1924		0.5504	
Inhalants	0.0047		0.2930		0.4667	
Nonmedical Use of Any Psychotherapeutic ²	0.0077		0.0930		0.6614	
Pain Relievers	0.2107		0.3552		0.6925	
Tranquilizers	0.2503		0.6754		0.6615	
Stimulants	0.0001		0.0172		0.0163	
Methamphetamine	0.0000		0.0744		0.0588	
Sedatives	0.0532		0.7729		0.2381	
Any Illicit Drug Other Than Marijuana ¹	0.0383		0.0825		0.5779	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.4C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 26 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	898	751	451	398	332	300
Marijuana and Hashish	841	734	379	339	261	248
Cocaine	580	505	170	135	123	102
Crack	290	252	115	66	77	48
Heroin	224	198	50	42	31	33
Hallucinogens	525	453	59	81	42	39
LSD	456	408	49	56	30	20
PCP	295	275	19	23	*	7
Inhalants	416	355	61	68	40	59
Nonmedical Use of Any Psychotherapeutic ²	621	566	272	243	195	170
Pain Relievers	468	438	241	204	173	134
Tranquilizers	451	383	142	154	108	94
Stimulants	452	386	116	107	91	71
Methamphetamine	369	289	93	85	66	58
Sedatives	350	306	81	67	59	36
Any Illicit Drug Other Than Marijuana ¹	746	662	308	274	225	195

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.4D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 26 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.53	0.44	0.27	0.23	0.20	0.18
Marijuana and Hashish	0.50	0.43	0.22	0.20	0.15	0.15
Cocaine	0.34	0.30	0.10	0.08	0.07	0.06
Crack	0.17	0.15	0.07	0.04	0.05	0.03
Heroin	0.13	0.12	0.03	0.02	0.02	0.02
Hallucinogens	0.31	0.26	0.03	0.05	0.02	0.02
LSD	0.27	0.24	0.03	0.03	0.02	0.01
PCP	0.17	0.16	0.01	0.01	*	0.00
Inhalants	0.25	0.21	0.04	0.04	0.02	0.03
Nonmedical Use of Any Psychotherapeutic ²	0.37	0.33	0.16	0.14	0.11	0.10
Pain Relievers	0.28	0.26	0.14	0.12	0.10	0.08
Tranquilizers	0.27	0.22	0.08	0.09	0.06	0.05
Stimulants	0.27	0.23	0.07	0.06	0.05	0.04
Methamphetamine	0.22	0.17	0.06	0.05	0.04	0.03
Sedatives	0.21	0.18	0.05	0.04	0.03	0.02
Any Illicit Drug Other Than Marijuana ¹	0.44	0.39	0.18	0.16	0.13	0.11

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.4P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 26 or Older: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.3184		0.3454		0.6538	
Marijuana and Hashish	0.6525		0.4785		0.3787	
Cocaine	0.5398		0.3156		0.1003	
Crack	0.1715		0.0713		0.1652	
Heroin	0.5378		0.8842		0.8644	
Hallucinogens	0.2595		0.0157		0.5384	
LSD	0.4486		0.8100		0.8293	
PCP	0.8832		0.3119		*	
Inhalants	0.7315		0.2160		0.1969	
Nonmedical Use of Any Psychotherapeutic ²	0.0614		0.3253		0.6019	
Pain Relievers	0.2187		0.7922		0.7497	
Tranquilizers	0.1230		0.9624		0.5655	
Stimulants	0.0796		0.4842		0.3160	
Methamphetamine	0.4534		0.6163		0.7888	
Sedatives	0.1920		0.8765		0.6612	
Any Illicit Drug Other Than Marijuana ¹	0.4316		0.6127		0.3108	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.5C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 12 or 13: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	35	36	30	28	20	18
Marijuana and Hashish	21	22	20	17	13	11
Cocaine	7	6	7	6	4	4
Crack	2	4	2	4	1	1
Heroin	2	3	2	3	1	1
Hallucinogens	13	13	10	9	6	5
LSD	8	9	6	7	3	2
PCP	6	6	5	4	1	1
Inhalants	27	27	20	18	12	8
Nonmedical Use of Any Psychotherapeutic ²	25	23	21	18	13	12
Pain Relievers	21	21	17	15	12	11
Tranquilizers	10	9	8	6	5	3
Stimulants	12	14	9	11	5	6
Methamphetamine	5	5	4	5	3	3
Sedatives	8	7	7	6	4	3
Any Illicit Drug Other Than Marijuana ¹	34	32	27	25	17	15

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.5D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 12 or 13: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.46	0.46	0.39	0.36	0.26	0.23
Marijuana and Hashish	0.28	0.29	0.26	0.21	0.17	0.15
Cocaine	0.10	0.08	0.09	0.07	0.05	0.05
Crack	0.03	0.05	0.02	0.05	0.02	0.02
Heroin	0.03	0.04	0.03	0.04	0.02	0.01
Hallucinogens	0.17	0.17	0.13	0.12	0.08	0.06
LSD	0.10	0.11	0.08	0.09	0.04	0.03
PCP	0.07	0.07	0.06	0.05	0.02	0.01
Inhalants	0.36	0.34	0.26	0.23	0.15	0.11
Nonmedical Use of Any Psychotherapeutic ²	0.32	0.30	0.27	0.23	0.18	0.16
Pain Relievers	0.28	0.26	0.22	0.19	0.15	0.14
Tranquilizers	0.13	0.12	0.10	0.08	0.07	0.04
Stimulants	0.16	0.19	0.12	0.14	0.07	0.07
Methamphetamine	0.06	0.07	0.05	0.06	0.04	0.04
Sedatives	0.11	0.09	0.09	0.07	0.05	0.04
Any Illicit Drug Other Than Marijuana ¹	0.45	0.41	0.35	0.32	0.23	0.19

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.5P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 12 or 13: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.2639		0.0060		0.0078	
Marijuana and Hashish	0.8527		0.1169		0.1535	
Cocaine	0.4814		0.3559		0.7521	
Crack	0.4890		0.2867		0.7319	
Heroin	0.9111		0.8258		0.5457	
Hallucinogens	0.5275		0.8304		0.5173	
LSD	0.6884		0.7102		0.4954	
PCP	0.6213		0.8178		0.3625	
Inhalants	0.9788		0.0410		0.0034	
Nonmedical Use of Any Psychotherapeutic ²	0.8072		0.1926		0.4676	
Pain Relievers	0.7185		0.2392		0.6131	
Tranquilizers	0.9250		0.6593		0.5998	
Stimulants	0.3626		0.2497		0.8290	
Methamphetamine	0.7486		0.4031		0.9497	
Sedatives	0.7004		0.7081		0.6050	
Any Illicit Drug Other Than Marijuana ¹	0.4485		0.0318		0.0509	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.6C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 14 or 15: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	49	47	44	41	32	31
Marijuana and Hashish	42	39	37	34	27	26
Cocaine	14	15	12	14	7	8
Crack	9	7	8	7	3	4
Heroin	5	6	5	4	5	2
Hallucinogens	22	22	19	19	9	10
LSD	18	18	14	15	8	6
PCP	11	10	7	7	4	2
Inhalants	33	31	22	20	11	11
Nonmedical Use of Any Psychotherapeutic ²	33	32	30	26	19	18
Pain Relievers	29	29	26	23	17	16
Tranquilizers	16	15	13	11	8	6
Stimulants	20	19	17	16	9	9
Methamphetamine	12	11	10	8	5	4
Sedatives	9	7	8	5	5	3
Any Illicit Drug Other Than Marijuana ¹	43	40	36	33	23	23

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.6D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 14 or 15: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.63	0.58	0.55	0.51	0.41	0.38
Marijuana and Hashish	0.53	0.48	0.48	0.43	0.35	0.32
Cocaine	0.18	0.18	0.15	0.17	0.09	0.10
Crack	0.11	0.09	0.10	0.09	0.04	0.04
Heroin	0.07	0.07	0.07	0.05	0.07	0.02
Hallucinogens	0.28	0.27	0.24	0.24	0.12	0.12
LSD	0.22	0.22	0.18	0.18	0.10	0.07
PCP	0.14	0.12	0.09	0.09	0.05	0.03
Inhalants	0.42	0.39	0.27	0.25	0.14	0.13
Nonmedical Use of Any Psychotherapeutic ²	0.42	0.40	0.38	0.32	0.24	0.22
Pain Relievers	0.37	0.36	0.33	0.29	0.21	0.20
Tranquilizers	0.20	0.19	0.17	0.14	0.10	0.07
Stimulants	0.26	0.24	0.21	0.19	0.12	0.11
Methamphetamine	0.15	0.13	0.13	0.10	0.06	0.05
Sedatives	0.11	0.09	0.10	0.06	0.07	0.04
Any Illicit Drug Other Than Marijuana ¹	0.54	0.49	0.46	0.41	0.30	0.28

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.6P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 14 or 15: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.2463		0.3277		0.9928	
Marijuana and Hashish	0.8831		0.7892		0.9436	
Cocaine	0.9084		0.2332		0.9127	
Crack	0.3584		0.7418		0.9043	
Heroin	0.1790		0.7090		0.0567	
Hallucinogens	0.3375		0.3200		0.7476	
LSD	0.9205		0.4184		0.1583	
PCP	0.4942		0.3318		0.3772	
Inhalants	0.8993		0.7565		0.4787	
Nonmedical Use of Any Psychotherapeutic ²	0.3586		0.1391		0.3004	
Pain Relievers	0.6985		0.1291		0.7675	
Tranquilizers	0.4228		0.2719		0.1023	
Stimulants	0.6346		0.6315		0.9884	
Methamphetamine	0.7280		0.3040		0.4157	
Sedatives	0.5186		0.1806		0.1807	
Any Illicit Drug Other Than Marijuana ¹	0.4706		0.5694		0.5428	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.7C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 16 or 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	53	51	48	48	35	40
Marijuana and Hashish	50	48	45	45	34	37
Cocaine	23	24	19	20	11	12
Crack	11	10	9	7	3	3
Heroin	12	7	10	5	8	4
Hallucinogens	33	34	27	29	15	19
LSD	27	29	23	22	13	13
PCP	13	13	7	8	3	3
Inhalants	33	28	21	19	9	10
Nonmedical Use of Any Psychotherapeutic ²	38	41	29	33	18	22
Pain Relievers	32	36	27	30	16	20
Tranquilizers	20	21	16	17	8	9
Stimulants	26	25	18	20	8	11
Methamphetamine	16	16	10	12	5	9
Sedatives	10	12	6	8	4	5
Any Illicit Drug Other Than Marijuana ¹	46	47	36	40	24	29

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.7D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 16 or 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.69	0.67	0.62	0.63	0.45	0.53
Marijuana and Hashish	0.64	0.64	0.59	0.59	0.44	0.49
Cocaine	0.30	0.32	0.24	0.27	0.14	0.16
Crack	0.15	0.13	0.11	0.09	0.03	0.04
Heroin	0.15	0.09	0.13	0.07	0.10	0.05
Hallucinogens	0.43	0.45	0.35	0.38	0.20	0.25
LSD	0.35	0.39	0.29	0.29	0.16	0.18
PCP	0.16	0.17	0.10	0.10	0.04	0.05
Inhalants	0.42	0.38	0.27	0.25	0.12	0.13
Nonmedical Use of Any Psychotherapeutic ²	0.49	0.54	0.37	0.44	0.23	0.29
Pain Relievers	0.42	0.48	0.34	0.40	0.21	0.26
Tranquilizers	0.26	0.28	0.20	0.23	0.10	0.12
Stimulants	0.34	0.33	0.23	0.26	0.10	0.15
Methamphetamine	0.21	0.21	0.13	0.17	0.06	0.11
Sedatives	0.12	0.15	0.08	0.10	0.05	0.06
Any Illicit Drug Other Than Marijuana ¹	0.59	0.62	0.47	0.53	0.31	0.39

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.7P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 16 or 17: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.8851		0.4536		0.1647	
Marijuana and Hashish	0.7597		0.2334		0.4259	
Cocaine	0.7967		0.8423		0.4560	
Crack	0.5977		0.7232		0.6778	
Heroin	0.0443		0.0841		0.1840	
Hallucinogens	0.6306		0.9046		0.4606	
LSD	0.5290		0.0785		0.5094	
PCP	0.3870		0.8303		0.7663	
Inhalants	0.3153		0.3901		0.4825	
Nonmedical Use of Any Psychotherapeutic ²	0.2235		0.0482		0.0254	
Pain Relievers	0.1250		0.0733		0.0317	
Tranquilizers	0.1727		0.1220		0.3876	
Stimulants	0.6213		0.2121		0.0402	
Methamphetamine	0.7438		0.0639		0.0357	
Sedatives	0.5739		0.1843		0.6624	
Any Illicit Drug Other Than Marijuana ¹	0.4524		0.5227		0.0168	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.8C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 18 to 20: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	92	88	88	84	74	72
Marijuana and Hashish	91	88	84	82	70	69
Cocaine	52	53	39	36	22	24
Crack	30	29	19	14	9	8
Heroin	21	18	13	9	6	4
Hallucinogens	65	70	47	54	28	28
LSD	60	58	41	37	22	20
PCP	26	26	13	10	7	5
Inhalants	58	57	35	33	17	16
Nonmedical Use of Any Psychotherapeutic ²	69	68	55	53	37	36
Pain Relievers	63	61	50	46	31	30
Tranquilizers	44	42	31	32	17	18
Stimulants	51	46	36	32	22	18
Methamphetamine	40	32	25	22	13	11
Sedatives	24	22	14	17	8	6
Any Illicit Drug Other Than Marijuana ¹	82	80	69	67	44	46

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.8D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 18 to 20: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.77	0.73	0.73	0.70	0.61	0.60
Marijuana and Hashish	0.75	0.73	0.70	0.68	0.58	0.57
Cocaine	0.43	0.44	0.32	0.30	0.19	0.20
Crack	0.25	0.24	0.16	0.11	0.08	0.06
Heroin	0.17	0.15	0.11	0.08	0.05	0.03
Hallucinogens	0.54	0.58	0.39	0.45	0.23	0.23
LSD	0.50	0.48	0.34	0.31	0.18	0.17
PCP	0.21	0.22	0.11	0.09	0.06	0.04
Inhalants	0.48	0.47	0.29	0.28	0.14	0.13
Nonmedical Use of Any Psychotherapeutic ²	0.57	0.57	0.45	0.44	0.31	0.30
Pain Relievers	0.52	0.50	0.41	0.38	0.26	0.25
Tranquilizers	0.37	0.35	0.25	0.26	0.14	0.15
Stimulants	0.43	0.38	0.30	0.26	0.18	0.15
Methamphetamine	0.33	0.26	0.21	0.18	0.11	0.09
Sedatives	0.20	0.18	0.12	0.14	0.07	0.05
Any Illicit Drug Other Than Marijuana ¹	0.68	0.66	0.58	0.56	0.37	0.38

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.8P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 18 to 20: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.6958		0.7434		0.8837	
Marijuana and Hashish	0.8882		0.8373		0.9418	
Cocaine	0.1928		0.0286		0.8178	
Crack	0.3599		0.2974		0.3804	
Heroin	0.2794		0.1298		0.3603	
Hallucinogens	0.1859		0.4604		0.7589	
LSD	0.7968		0.2308		0.4740	
PCP	0.7259		0.4282		0.6888	
Inhalants	0.6014		0.8200		0.6933	
Nonmedical Use of Any Psychotherapeutic ²	0.5443		0.9078		0.9329	
Pain Relievers	0.4160		0.5427		0.7835	
Tranquilizers	0.4661		0.7463		0.8541	
Stimulants	0.0239		0.3359		0.3799	
Methamphetamine	0.0010		0.3897		0.1961	
Sedatives	0.3106		0.5305		0.3655	
Any Illicit Drug Other Than Marijuana ¹	0.9266		0.7328		0.7254	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.9C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 21 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	106	110	100	91	80	72
Marijuana and Hashish	103	110	95	84	78	64
Cocaine	72	65	42	37	24	20
Crack	35	31	19	13	8	4
Heroin	30	26	12	11	6	6
Hallucinogens	89	83	49	45	23	20
LSD	79	70	32	29	13	13
PCP	29	26	10	8	5	4
Inhalants	72	66	28	23	14	11
Nonmedical Use of Any Psychotherapeutic ²	86	78	61	51	32	34
Pain Relievers	77	65	54	45	27	30
Tranquilizers	56	56	35	31	20	17
Stimulants	58	55	33	27	17	12
Methamphetamine	41	46	23	19	13	9
Sedatives	28	25	14	11	8	7
Any Illicit Drug Other Than Marijuana ¹	100	97	73	68	43	41

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.9D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 21 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.64	0.65	0.61	0.54	0.49	0.43
Marijuana and Hashish	0.63	0.65	0.58	0.49	0.47	0.38
Cocaine	0.44	0.38	0.26	0.22	0.15	0.12
Crack	0.21	0.19	0.11	0.07	0.05	0.02
Heroin	0.18	0.15	0.07	0.07	0.04	0.04
Hallucinogens	0.54	0.49	0.30	0.26	0.14	0.12
LSD	0.48	0.41	0.20	0.17	0.08	0.08
PCP	0.17	0.16	0.06	0.05	0.03	0.02
Inhalants	0.44	0.39	0.17	0.14	0.08	0.06
Nonmedical Use of Any Psychotherapeutic ²	0.52	0.46	0.37	0.30	0.20	0.20
Pain Relievers	0.47	0.39	0.33	0.27	0.16	0.18
Tranquilizers	0.34	0.33	0.21	0.18	0.12	0.10
Stimulants	0.35	0.32	0.20	0.16	0.11	0.07
Methamphetamine	0.25	0.27	0.14	0.11	0.08	0.05
Sedatives	0.17	0.15	0.08	0.07	0.05	0.04
Any Illicit Drug Other Than Marijuana ¹	0.61	0.57	0.44	0.40	0.26	0.24

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.9P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 21 to 25: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.0212		0.0259		0.1598	
Marijuana and Hashish	0.0534		0.1301		0.1215	
Cocaine	0.0418		0.0206		0.0102	
Crack	0.0509		0.0097		0.0199	
Heroin	0.0731		0.9532		0.7160	
Hallucinogens	0.3142		0.4773		0.5009	
LSD	0.0517		0.0706		0.2177	
PCP	0.2367		0.3070		0.6577	
Inhalants	0.0008		0.0533		0.4983	
Nonmedical Use of Any Psychotherapeutic ²	0.0021		0.0121		0.4505	
Pain Relievers	0.0167		0.0572		0.7279	
Tranquilizers	0.3178		0.7996		0.4198	
Stimulants	0.0016		0.0107		0.0055	
Methamphetamine	0.0103		0.0799		0.1517	
Sedatives	0.0969		0.2148		0.4276	
Any Illicit Drug Other Than Marijuana ¹	0.0031		0.0048		0.2527	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.10C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 26 to 34: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	252	221	171	145	121	119
Marijuana and Hashish	241	220	151	136	106	109
Cocaine	186	147	74	58	54	41
Crack	115	76	40	23	28	14
Heroin	60	44	18	7	17	3
Hallucinogens	187	160	33	52	14	44
LSD	160	141	21	27	7	20
PCP	68	57	*	11	*	3
Inhalants	155	136	27	31	24	23
Nonmedical Use of Any Psychotherapeutic ²	181	155	90	78	49	63
Pain Relievers	148	127	78	68	38	51
Tranquilizers	126	110	51	43	32	25
Stimulants	132	104	40	43	23	34
Methamphetamine	110	89	29	37	18	25
Sedatives	71	59	16	21	12	10
Any Illicit Drug Other Than Marijuana ¹	233	204	112	101	76	84

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.10D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 26 to 34: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.75	0.67	0.51	0.44	0.36	0.36
Marijuana and Hashish	0.72	0.67	0.45	0.41	0.32	0.33
Cocaine	0.55	0.45	0.22	0.18	0.16	0.13
Crack	0.34	0.23	0.12	0.07	0.08	0.04
Heroin	0.18	0.13	0.05	0.02	0.05	0.01
Hallucinogens	0.56	0.49	0.10	0.16	0.04	0.13
LSD	0.48	0.43	0.06	0.08	0.02	0.06
PCP	0.20	0.17	*	0.03	*	0.01
Inhalants	0.46	0.41	0.08	0.09	0.07	0.07
Nonmedical Use of Any Psychotherapeutic ²	0.54	0.47	0.27	0.24	0.15	0.19
Pain Relievers	0.44	0.38	0.23	0.21	0.11	0.16
Tranquilizers	0.37	0.33	0.15	0.13	0.10	0.08
Stimulants	0.39	0.31	0.12	0.13	0.07	0.10
Methamphetamine	0.33	0.27	0.09	0.11	0.05	0.08
Sedatives	0.21	0.18	0.05	0.06	0.03	0.03
Any Illicit Drug Other Than Marijuana ¹	0.69	0.62	0.33	0.31	0.22	0.26

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.10P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 26 to 34: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.0149		0.9017		0.0648	
Marijuana and Hashish	0.0571		0.9816		0.2615	
Cocaine	0.0001		0.1555		0.0516	
Crack	0.0011		0.0227		0.0068	
Heroin	0.3942		0.2665		0.1627	
Hallucinogens	0.3501		0.0027		0.0495	
LSD	0.3680		0.4949		0.1863	
PCP	0.1271		*		*	
Inhalants	0.5750		0.0989		0.5795	
Nonmedical Use of Any Psychotherapeutic ²	0.0204		0.6790		0.0074	
Pain Relievers	0.3106		0.8171		0.0008	
Tranquilizers	0.0075		0.5878		0.5977	
Stimulants	0.0037		0.9906		0.2396	
Methamphetamine	0.1826		0.6720		0.4484	
Sedatives	0.1719		0.3215		0.6543	
Any Illicit Drug Other Than Marijuana ¹	0.0090		0.9695		0.2236	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.11C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 35 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	839	704	409	375	309	270
Marijuana and Hashish	789	691	341	309	246	217
Cocaine	528	485	147	126	108	91
Crack	266	235	108	62	71	45
Heroin	215	191	47	42	26	33
Hallucinogens	483	427	49	55	40	23
LSD	424	387	44	43	29	*
PCP	285	263	19	20	*	7
Inhalants	384	333	54	60	32	54
Nonmedical Use of Any Psychotherapeutic ²	581	536	254	231	187	158
Pain Relievers	431	402	227	189	169	126
Tranquilizers	430	354	135	149	103	88
Stimulants	435	370	106	96	85	64
Methamphetamine	353	275	88	72	57	52
Sedatives	340	301	78	64	58	34
Any Illicit Drug Other Than Marijuana ¹	686	625	278	258	204	184

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.11D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 35 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.62	0.51	0.30	0.27	0.23	0.20
Marijuana and Hashish	0.58	0.50	0.25	0.22	0.18	0.16
Cocaine	0.39	0.35	0.11	0.09	0.08	0.07
Crack	0.20	0.17	0.08	0.05	0.05	0.03
Heroin	0.16	0.14	0.03	0.03	0.02	0.02
Hallucinogens	0.36	0.31	0.04	0.04	0.03	0.02
LSD	0.31	0.28	0.03	0.03	0.02	*
PCP	0.21	0.19	0.01	0.01	*	0.00
Inhalants	0.28	0.24	0.04	0.04	0.02	0.04
Nonmedical Use of Any Psychotherapeutic ²	0.43	0.39	0.19	0.17	0.14	0.11
Pain Relievers	0.32	0.29	0.17	0.14	0.12	0.09
Tranquilizers	0.32	0.26	0.10	0.11	0.08	0.06
Stimulants	0.32	0.27	0.08	0.07	0.06	0.05
Methamphetamine	0.26	0.20	0.06	0.05	0.04	0.04
Sedatives	0.25	0.22	0.06	0.05	0.04	0.02
Any Illicit Drug Other Than Marijuana ¹	0.51	0.45	0.20	0.19	0.15	0.13

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.11P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 35 or Older: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.8473		0.3996		0.8151	
Marijuana and Hashish	0.8188		0.5305		0.6043	
Cocaine	0.4778		0.6898		0.3517	
Crack	0.8404		0.2842		0.6325	
Heroin	0.6652		0.8576		0.4453	
Hallucinogens	0.1011		0.3977		0.2936	
LSD	0.2203		0.9608		*	
PCP	0.6360		0.7834		*	
Inhalants	0.9806		0.5482		0.2341	
Nonmedical Use of Any Psychotherapeutic ²	0.2168		0.3948		0.1547	
Pain Relievers	0.3494		0.7504		0.5118	
Tranquilizers	0.4382		0.8710		0.6664	
Stimulants	0.3364		0.4442		0.1145	
Methamphetamine	0.7221		0.4627		0.5388	
Sedatives	0.2831		0.6679		0.7275	
Any Illicit Drug Other Than Marijuana ¹	0.8831		0.6208		0.1130	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.12C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Males: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	612	562	372	336	283	260
Marijuana and Hashish	582	549	331	297	250	230
Cocaine	448	409	148	113	105	78
Crack	246	203	101	53	70	38
Heroin	201	155	31	34	28	19
Hallucinogens	390	380	87	115	51	62
LSD	345	345	69	77	36	33
PCP	235	210	25	26	9	7
Inhalants	366	319	78	78	48	61
Nonmedical Use of Any Psychotherapeutic ²	461	422	197	168	140	128
Pain Relievers	341	318	182	144	133	105
Tranquilizers	324	268	116	89	83	59
Stimulants	343	295	89	83	51	58
Methamphetamine	309	234	77	65	41	42
Sedatives	264	221	67	46	58	32
Any Illicit Drug Other Than Marijuana ¹	525	544	242	224	174	163

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.12D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Males: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.58	0.52	0.35	0.31	0.27	0.24
Marijuana and Hashish	0.55	0.51	0.31	0.28	0.23	0.21
Cocaine	0.42	0.38	0.14	0.11	0.10	0.07
Crack	0.23	0.19	0.09	0.05	0.07	0.04
Heroin	0.19	0.14	0.03	0.03	0.03	0.02
Hallucinogens	0.37	0.35	0.08	0.11	0.05	0.06
LSD	0.33	0.32	0.06	0.07	0.03	0.03
PCP	0.22	0.20	0.02	0.02	0.01	0.01
Inhalants	0.34	0.30	0.07	0.07	0.05	0.06
Nonmedical Use of Any Psychotherapeutic ²	0.43	0.39	0.19	0.16	0.13	0.12
Pain Relievers	0.32	0.30	0.17	0.13	0.13	0.10
Tranquilizers	0.31	0.25	0.11	0.08	0.08	0.06
Stimulants	0.32	0.27	0.08	0.08	0.05	0.05
Methamphetamine	0.29	0.22	0.07	0.06	0.04	0.04
Sedatives	0.25	0.21	0.06	0.04	0.05	0.03
Any Illicit Drug Other Than Marijuana ¹	0.49	0.51	0.23	0.21	0.16	0.15

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.12P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Males: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.8644		0.0572		0.2780	
Marijuana and Hashish	0.9721		0.0748		0.2695	
Cocaine	0.5455		0.1980		0.0984	
Crack	0.0243		0.0593		0.2392	
Heroin	0.1040		0.9372		0.3592	
Hallucinogens	0.2781		0.2204		0.8697	
LSD	0.7553		0.5582		0.8142	
PCP	0.7520		0.9179		0.8744	
Inhalants	0.0701		0.7759		0.7971	
Nonmedical Use of Any Psychotherapeutic ²	0.1396		0.1696		0.5007	
Pain Relievers	0.1938		0.8238		0.7293	
Tranquilizers	0.0512		0.2675		0.3604	
Stimulants	0.0431		0.4855		0.7288	
Methamphetamine	0.0677		0.3646		0.4530	
Sedatives	0.0768		0.8803		0.6420	
Any Illicit Drug Other Than Marijuana ¹	0.9889		0.2486		0.3213	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.13C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Females: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	666	512	310	286	218	211
Marijuana and Hashish	609	509	236	219	147	162
Cocaine	356	323	106	99	71	74
Crack	145	152	63	43	33	30
Heroin	100	138	44	32	18	29
Hallucinogens	354	304	56	67	30	37
LSD	301	273	40	36	23	18
PCP	168	176	12	13	6	8
Inhalants	217	218	40	39	20	26
Nonmedical Use of Any Psychotherapeutic ²	447	381	211	201	149	124
Pain Relievers	337	307	180	164	124	100
Tranquilizers	296	266	102	137	79	72
Stimulants	293	270	90	79	77	49
Methamphetamine	213	197	66	58	48	44
Sedatives	222	205	51	55	19	20
Any Illicit Drug Other Than Marijuana ¹	539	441	229	219	165	143

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.13D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Females: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.58	0.44	0.27	0.25	0.19	0.18
Marijuana and Hashish	0.53	0.44	0.21	0.19	0.13	0.14
Cocaine	0.31	0.28	0.09	0.09	0.06	0.06
Crack	0.13	0.13	0.06	0.04	0.03	0.03
Heroin	0.09	0.12	0.04	0.03	0.02	0.02
Hallucinogens	0.31	0.26	0.05	0.06	0.03	0.03
LSD	0.26	0.24	0.04	0.03	0.02	0.02
PCP	0.15	0.15	0.01	0.01	0.01	0.01
Inhalants	0.19	0.19	0.03	0.03	0.02	0.02
Nonmedical Use of Any Psychotherapeutic ²	0.39	0.33	0.18	0.17	0.13	0.11
Pain Relievers	0.29	0.26	0.16	0.14	0.11	0.09
Tranquilizers	0.26	0.23	0.09	0.12	0.07	0.06
Stimulants	0.25	0.23	0.08	0.07	0.07	0.04
Methamphetamine	0.19	0.17	0.06	0.05	0.04	0.04
Sedatives	0.19	0.18	0.04	0.05	0.02	0.02
Any Illicit Drug Other Than Marijuana ¹	0.47	0.38	0.20	0.19	0.14	0.12

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.13P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Females: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.0666		0.6957		0.1220	
Marijuana and Hashish	0.2493		0.7753		0.0177	
Cocaine	0.4124		0.1333		0.1994	
Crack	0.7029		0.1376		0.1501	
Heroin	0.5063		0.3388		0.8754	
Hallucinogens	0.6251		0.3010		0.5647	
LSD	0.8363		0.0330		0.0183	
PCP	0.4966		0.5037		0.9788	
Inhalants	0.3748		0.8987		0.4146	
Nonmedical Use of Any Psychotherapeutic ²	0.0801		0.5210		0.9470	
Pain Relievers	0.5561		0.5996		0.6246	
Tranquilizers	0.6404		0.4075		0.8979	
Stimulants	0.2304		0.3078		0.1256	
Methamphetamine	0.8648		0.7581		0.9182	
Sedatives	0.7467		0.8522		0.3713	
Any Illicit Drug Other Than Marijuana ¹	0.0934		0.6585		0.7048	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.14C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Males Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	55	61	54	51	41	38
Marijuana and Hashish	52	51	49	44	37	35
Cocaine	21	20	16	18	10	12
Crack	10	9	8	7	2	3
Heroin	10	7	9	5	7	3
Hallucinogens	31	31	25	26	13	15
LSD	24	28	20	23	10	12
PCP	13	13	9	10	3	4
Inhalants	39	38	27	24	13	12
Nonmedical Use of Any Psychotherapeutic ²	39	43	32	32	21	21
Pain Relievers	34	38	28	28	19	19
Tranquilizers	19	19	15	14	9	7
Stimulants	24	23	16	18	7	10
Methamphetamine	14	14	9	10	4	7
Sedatives	11	10	8	7	5	5
Any Illicit Drug Other Than Marijuana ¹	49	53	42	41	27	26

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.14D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Males Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.47	0.51	0.45	0.42	0.34	0.31
Marijuana and Hashish	0.44	0.42	0.41	0.36	0.31	0.29
Cocaine	0.17	0.17	0.14	0.15	0.08	0.10
Crack	0.09	0.07	0.07	0.06	0.02	0.03
Heroin	0.09	0.06	0.07	0.04	0.06	0.02
Hallucinogens	0.26	0.26	0.21	0.21	0.11	0.13
LSD	0.20	0.23	0.17	0.19	0.08	0.10
PCP	0.11	0.11	0.07	0.08	0.03	0.03
Inhalants	0.33	0.32	0.23	0.20	0.11	0.10
Nonmedical Use of Any Psychotherapeutic ²	0.33	0.36	0.27	0.26	0.18	0.18
Pain Relievers	0.29	0.32	0.24	0.24	0.16	0.16
Tranquilizers	0.16	0.16	0.13	0.11	0.07	0.06
Stimulants	0.20	0.19	0.14	0.15	0.06	0.09
Methamphetamine	0.12	0.11	0.08	0.09	0.03	0.05
Sedatives	0.10	0.08	0.07	0.06	0.04	0.04
Any Illicit Drug Other Than Marijuana ¹	0.42	0.45	0.35	0.34	0.23	0.22

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.14P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Males Aged 12 to 17: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.0956		0.0175		0.6434	
Marijuana and Hashish	0.0980		0.0150		0.9758	
Cocaine	0.6278		0.3047		0.4438	
Crack	0.9113		0.8695		0.2110	
Heroin	0.1998		0.1817		0.1042	
Hallucinogens	0.5470		0.6539		0.1384	
LSD	0.6588		0.9805		0.4153	
PCP	0.3888		0.3529		0.9040	
Inhalants	0.7382		0.4633		0.5937	
Nonmedical Use of Any Psychotherapeutic ²	0.5485		0.3620		0.8130	
Pain Relievers	0.0813		0.5753		0.7171	
Tranquilizers	0.8045		0.9651		0.4878	
Stimulants	0.7843		0.1104		0.0459	
Methamphetamine	0.8612		0.0769		0.0721	
Sedatives	0.2218		0.7780		0.7525	
Any Illicit Drug Other Than Marijuana ¹	0.9837		0.6076		0.5893	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.15C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Females Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	58	55	52	50	37	38
Marijuana and Hashish	50	48	44	43	32	32
Cocaine	20	22	17	18	10	10
Crack	10	10	8	8	4	4
Heroin	8	7	7	6	5	3
Hallucinogens	30	28	24	24	14	14
LSD	24	22	18	16	12	8
PCP	11	12	7	7	4	3
Inhalants	37	32	27	22	13	12
Nonmedical Use of Any Psychotherapeutic ²	42	40	35	33	22	23
Pain Relievers	37	36	30	29	18	21
Tranquilizers	21	22	17	17	9	8
Stimulants	26	27	21	21	11	12
Methamphetamine	15	14	12	11	7	8
Sedatives	11	12	9	9	6	5
Any Illicit Drug Other Than Marijuana ¹	51	48	42	41	27	28

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.15D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Females Aged 12 to 17: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.51	0.48	0.46	0.44	0.33	0.34
Marijuana and Hashish	0.44	0.42	0.39	0.38	0.28	0.28
Cocaine	0.17	0.19	0.15	0.16	0.09	0.09
Crack	0.09	0.09	0.07	0.07	0.03	0.03
Heroin	0.07	0.06	0.06	0.05	0.05	0.03
Hallucinogens	0.27	0.24	0.21	0.21	0.12	0.12
LSD	0.21	0.19	0.16	0.14	0.10	0.07
PCP	0.10	0.10	0.06	0.06	0.04	0.02
Inhalants	0.33	0.28	0.24	0.20	0.12	0.10
Nonmedical Use of Any Psychotherapeutic ²	0.37	0.35	0.31	0.29	0.19	0.20
Pain Relievers	0.32	0.31	0.27	0.25	0.16	0.18
Tranquilizers	0.19	0.19	0.15	0.15	0.08	0.07
Stimulants	0.23	0.23	0.18	0.19	0.10	0.10
Methamphetamine	0.13	0.12	0.10	0.10	0.06	0.07
Sedatives	0.09	0.10	0.08	0.08	0.05	0.04
Any Illicit Drug Other Than Marijuana ¹	0.45	0.42	0.37	0.36	0.24	0.25

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.15P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Females Aged 12 to 17: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.6600		0.2487		0.8762	
Marijuana and Hashish	0.7135		0.8123		0.8685	
Cocaine	0.5273		0.6083		0.7627	
Crack	0.9607		0.8538		0.4712	
Heroin	0.8210		0.7291		0.0951	
Hallucinogens	0.8917		0.9081		0.4161	
LSD	0.0710		0.0670		0.0028	
PCP	0.5039		0.9500		0.1206	
Inhalants	0.4711		0.0769		0.4232	
Nonmedical Use of Any Psychotherapeutic ²	0.5652		0.2441		0.6670	
Pain Relievers	0.5362		0.3627		0.3224	
Tranquilizers	0.7365		0.6960		0.8377	
Stimulants	0.7142		0.5598		0.8377	
Methamphetamine	0.3927		0.8058		0.8877	
Sedatives	0.3709		0.9665		0.4241	
Any Illicit Drug Other Than Marijuana ¹	0.4660		0.3424		0.9787	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.16C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Males Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	97	103	92	89	82	75
Marijuana and Hashish	95	98	90	84	79	69
Cocaine	71	63	49	39	27	24
Crack	37	32	21	16	10	8
Heroin	29	24	14	11	7	7
Hallucinogens	86	78	54	53	27	28
LSD	77	66	42	36	21	20
PCP	30	29	14	11	8	5
Inhalants	70	69	38	32	18	16
Nonmedical Use of Any Psychotherapeutic ²	81	74	61	54	40	35
Pain Relievers	73	67	52	50	33	32
Tranquilizers	55	50	36	33	21	18
Stimulants	59	48	37	28	23	16
Methamphetamine	43	38	27	20	15	9
Sedatives	27	26	15	16	9	8
Any Illicit Drug Other Than Marijuana ¹	94	93	73	71	48	45

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.16D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Males Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.69	0.71	0.65	0.62	0.58	0.52
Marijuana and Hashish	0.67	0.68	0.64	0.58	0.56	0.48
Cocaine	0.50	0.44	0.34	0.27	0.19	0.16
Crack	0.26	0.22	0.15	0.11	0.07	0.05
Heroin	0.21	0.17	0.10	0.08	0.05	0.05
Hallucinogens	0.61	0.54	0.38	0.37	0.19	0.19
LSD	0.55	0.46	0.30	0.25	0.15	0.14
PCP	0.22	0.20	0.10	0.07	0.06	0.03
Inhalants	0.49	0.48	0.27	0.22	0.12	0.11
Nonmedical Use of Any Psychotherapeutic ²	0.57	0.52	0.43	0.37	0.28	0.24
Pain Relievers	0.52	0.47	0.37	0.35	0.23	0.22
Tranquilizers	0.39	0.34	0.25	0.23	0.15	0.13
Stimulants	0.41	0.34	0.26	0.20	0.16	0.11
Methamphetamine	0.31	0.27	0.19	0.14	0.11	0.06
Sedatives	0.19	0.18	0.11	0.11	0.06	0.06
Any Illicit Drug Other Than Marijuana ¹	0.67	0.65	0.51	0.49	0.34	0.31

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.16P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Males Aged 18 to 25: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.1371		0.0857		0.1197	
Marijuana and Hashish	0.1125		0.0998		0.0991	
Cocaine	0.0344		0.0045		0.0732	
Crack	0.0474		0.0474		0.0165	
Heroin	0.0703		0.3688		0.7382	
Hallucinogens	0.8178		0.6255		0.8018	
LSD	0.3612		0.0330		0.3865	
PCP	0.8112		0.1242		0.5075	
Inhalants	0.0096		0.1207		0.2226	
Nonmedical Use of Any Psychotherapeutic ²	0.1733		0.5353		0.3909	
Pain Relievers	0.8193		0.6697		0.9268	
Tranquilizers	0.5182		0.9397		0.6918	
Stimulants	0.0009		0.0250		0.0377	
Methamphetamine	0.0005		0.0123		0.0132	
Sedatives	0.4118		0.6167		0.6990	
Any Illicit Drug Other Than Marijuana ¹	0.2168		0.1627		0.2191	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.17C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Females Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	100	95	83	83	63	62
Marijuana and Hashish	98	97	77	78	59	58
Cocaine	50	54	33	34	20	19
Crack	26	27	14	10	6	5
Heroin	22	19	10	9	5	3
Hallucinogens	67	75	41	44	22	20
LSD	58	61	29	28	15	14
PCP	23	24	9	8	5	3
Inhalants	57	55	24	25	14	12
Nonmedical Use of Any Psychotherapeutic ²	73	69	55	52	29	32
Pain Relievers	65	58	50	42	25	27
Tranquilizers	45	47	27	29	16	16
Stimulants	47	46	29	29	15	14
Methamphetamine	37	36	19	22	10	10
Sedatives	24	20	12	14	6	4
Any Illicit Drug Other Than Marijuana ¹	84	87	64	64	37	41

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.17D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Females Aged 18 to 25: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.70	0.65	0.58	0.57	0.44	0.42
Marijuana and Hashish	0.69	0.66	0.54	0.53	0.41	0.40
Cocaine	0.35	0.37	0.23	0.23	0.14	0.13
Crack	0.18	0.19	0.10	0.07	0.04	0.04
Heroin	0.15	0.13	0.07	0.06	0.03	0.02
Hallucinogens	0.47	0.51	0.29	0.30	0.16	0.14
LSD	0.41	0.42	0.20	0.19	0.11	0.09
PCP	0.16	0.16	0.06	0.05	0.03	0.02
Inhalants	0.40	0.38	0.17	0.17	0.09	0.08
Nonmedical Use of Any Psychotherapeutic ²	0.51	0.47	0.38	0.35	0.20	0.22
Pain Relievers	0.45	0.40	0.35	0.29	0.17	0.19
Tranquilizers	0.32	0.32	0.19	0.20	0.11	0.11
Stimulants	0.33	0.32	0.21	0.20	0.10	0.10
Methamphetamine	0.26	0.25	0.14	0.15	0.07	0.07
Sedatives	0.17	0.14	0.09	0.09	0.04	0.03
Any Illicit Drug Other Than Marijuana ¹	0.59	0.60	0.45	0.44	0.26	0.28

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.17P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Females Aged 18 to 25: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.1587		0.2780		0.6868	
Marijuana and Hashish	0.5005		0.8282		0.9474	
Cocaine	0.2470		0.1979		0.3729	
Crack	0.4291		0.0695		0.8764	
Heroin	0.2494		0.6113		0.2114	
Hallucinogens	0.7737		0.5674		0.9596	
LSD	0.2689		0.3844		0.3147	
PCP	0.4903		0.9763		0.9586	
Inhalants	0.1145		0.6808		0.7097	
Nonmedical Use of Any Psychotherapeutic ²	0.0055		0.0558		0.7020	
Pain Relievers	0.0670		0.0488		0.5773	
Tranquilizers	0.2424		0.5491		0.8061	
Stimulants	0.0177		0.2308		0.1948	
Methamphetamine	0.0181		0.9235		0.9442	
Sedatives	0.0440		0.2972		0.0980	
Any Illicit Drug Other Than Marijuana ¹	0.0460		0.2501		0.4975	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.18C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Illicit Drugs Among Persons Aged 18 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	927	785	492	442	360	322
Marijuana and Hashish	868	771	419	387	290	273
Cocaine	594	515	187	146	130	110
Crack	292	258	119	67	78	49
Heroin	229	202	53	45	32	34
Hallucinogens	555	482	94	122	55	57
LSD	478	431	75	82	40	35
PCP	296	279	25	26	9	9
Inhalants	435	378	77	83	47	65
Nonmedical Use of Any Psychotherapeutic ²	639	581	290	258	206	178
Pain Relievers	487	458	256	215	176	141
Tranquilizers	459	401	153	161	113	98
Stimulants	463	394	127	116	97	76
Methamphetamine	381	296	100	91	68	61
Sedatives	353	310	84	71	60	38
Any Illicit Drug Other Than Marijuana ¹	776	693	330	300	239	208

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.18D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 18 or Older: 1999 and 2000

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.47	0.39	0.25	0.22	0.18	0.16
Marijuana and Hashish	0.44	0.39	0.21	0.19	0.15	0.14
Cocaine	0.30	0.26	0.09	0.07	0.07	0.06
Crack	0.15	0.13	0.06	0.03	0.04	0.02
Heroin	0.12	0.10	0.03	0.02	0.02	0.02
Hallucinogens	0.28	0.24	0.05	0.06	0.03	0.03
LSD	0.24	0.22	0.04	0.04	0.02	0.02
PCP	0.15	0.14	0.01	0.01	0.00	0.00
Inhalants	0.22	0.19	0.04	0.04	0.02	0.03
Nonmedical Use of Any Psychotherapeutic ²	0.32	0.29	0.15	0.13	0.10	0.09
Pain Relievers	0.25	0.23	0.13	0.11	0.09	0.07
Tranquilizers	0.23	0.20	0.08	0.08	0.06	0.05
Stimulants	0.23	0.20	0.06	0.06	0.05	0.04
Methamphetamine	0.19	0.15	0.05	0.05	0.03	0.03
Sedatives	0.18	0.15	0.04	0.04	0.03	0.02
Any Illicit Drug Other Than Marijuana ¹	0.39	0.35	0.17	0.15	0.12	0.10

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.18P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Illicit Drugs Among Persons Aged 18 or Older: 2000 Versus 1999

Drug	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Any Illicit Drug ¹	0.2051		0.1755		0.8582	
Marijuana and Hashish	0.4973		0.3217		0.6625	
Cocaine	0.3246		0.0559		0.0455	
Crack	0.0936		0.0189		0.0885	
Heroin	0.3456		0.6471		0.9631	
Hallucinogens	0.2707		0.0999		0.7252	
LSD	0.6578		0.2446		0.3304	
PCP	0.9484		0.9279		0.6453	
Inhalants	0.3249		0.6377		0.3773	
Nonmedical Use of Any Psychotherapeutic ²	0.0222		0.1713		0.5515	
Pain Relievers	0.1552		0.6333		0.6564	
Tranquilizers	0.0886		0.8851		0.5166	
Stimulants	0.0170		0.1253		0.1127	
Methamphetamine	0.1641		0.2961		0.4740	
Sedatives	0.1275		0.8277		0.5042	
Any Illicit Drug Other Than Marijuana ¹	0.2597		0.3386		0.2835	

*Low precision; no estimate reported.

¹ Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically. Any Illicit Drug Other Than Marijuana indicates use at least once of any of these listed drugs, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

² Nonmedical use of any prescription-type pain reliever, tranquilizer, stimulant, or sedative; does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.19C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	942	795	508	455	366	332
12	22	23	19	17	13	12
13	27	27	22	21	15	13
14	34	29	28	25	21	20
15	34	35	32	33	24	25
16	38	36	34	34	25	27
17	39	37	34	33	27	28
18	52	50	48	49	40	41
19	50	48	49	43	43	39
20	46	47	45	45	40	40
21	52	43	45	41	37	34
22	43	43	48	37	37	31
23	38	44	36	41	30	30
24	41	45	38	37	31	28
25	42	41	33	32	26	25
26-29	168	146	113	101	86	78
30-34	181	160	122	101	88	86
35-39	302	286	200	168	141	118
40-44	314	281	214	178	184	138
45-49	290	274	153	136	119	114
50-54	321	316	120	132	106	107
55-59	255	237	87	113	75	81
60-64	184	167	47	80	18	78
65 or Older	234	221	109	56	90	41

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.19D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.43	0.36	0.23	0.20	0.17	0.15
12	0.61	0.60	0.52	0.46	0.35	0.32
13	0.66	0.67	0.55	0.52	0.39	0.31
14	0.84	0.72	0.70	0.61	0.51	0.49
15	0.89	0.86	0.84	0.81	0.63	0.62
16	0.94	0.92	0.85	0.86	0.63	0.69
17	1.05	1.02	0.92	0.91	0.73	0.77
18	1.21	1.17	1.13	1.14	0.94	0.96
19	1.26	1.22	1.23	1.10	1.09	0.99
20	1.22	1.21	1.20	1.15	1.06	1.02
21	1.46	1.23	1.27	1.17	1.04	0.96
22	1.28	1.26	1.41	1.09	1.09	0.91
23	1.19	1.26	1.15	1.17	0.95	0.86
24	1.28	1.41	1.18	1.15	0.96	0.87
25	1.35	1.24	1.06	0.97	0.83	0.74
26-29	1.13	1.03	0.77	0.71	0.58	0.55
30-34	0.96	0.85	0.65	0.54	0.47	0.46
35-39	1.33	1.33	0.88	0.78	0.62	0.55
40-44	1.46	1.27	0.99	0.80	0.85	0.62
45-49	1.52	1.36	0.80	0.68	0.63	0.57
50-54	1.89	1.66	0.70	0.70	0.62	0.56
55-59	1.97	1.76	0.67	0.84	0.58	0.60
60-64	1.81	1.64	0.46	0.79	0.18	0.77
65 or Older	0.72	0.71	0.34	0.18	0.28	0.13

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.19P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug, by Detailed Age Categories: 2000 Versus 1999

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.1639		0.0845		0.8949	
12	0.1199		0.0192		0.1330	
13	0.9440		0.1185		0.0321	
14	0.0033		0.0304		0.5834	
15	0.3578		0.7462		0.7404	
16	0.9259		0.6250		0.8104	
17	0.8899		0.5822		0.0835	
18	0.9951		0.9831		0.3872	
19	0.6160		0.9128		0.9887	
20	0.8004		0.4665		0.5222	
21	0.8118		0.8441		0.5806	
22	0.6052		0.2282		0.1114	
23	0.0077		0.0317		0.0667	
24	0.0778		0.7610		0.7662	
25	0.6123		0.1240		0.5012	
26-29	0.6428		0.6855		0.2735	
30-34	0.0035		0.6225		0.1082	
35-39	0.0330		0.1000		0.2441	
40-44	0.9705		0.1271		0.2006	
45-49	0.5538		0.9064		0.1587	
50-54	0.8462		0.4213		0.9303	
55-59	0.0533		0.0476		0.2627	
60-64	0.8261		0.4334		0.1335	
65 or Older	0.4399		0.5405		0.4721	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.20C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	872	775	428	398	295	281
12	10	10	8	9	4	6
13	18	19	17	14	12	10
14	27	23	22	19	16	15
15	30	30	28	29	21	22
16	36	35	32	31	25	25
17	36	34	32	32	25	26
18	49	51	45	49	38	39
19	51	48	49	42	42	37
20	46	47	42	42	38	37
21	50	44	43	39	35	31
22	45	42	43	34	36	28
23	39	42	34	36	28	27
24	42	46	34	34	28	25
25	42	41	31	30	24	23
26-29	165	146	105	92	82	72
30-34	175	165	106	94	73	76
35-39	308	282	185	148	123	99
40-44	311	287	187	157	153	113
45-49	285	276	142	120	105	98
50-54	310	307	80	105	67	84
55-59	238	216	54	70	32	66
60-64	162	146	11	60	11	*
65 or Older	172	153	4	10	2	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.20D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.39	0.35	0.19	0.18	0.13	0.13
12	0.28	0.28	0.24	0.24	0.12	0.15
13	0.45	0.48	0.41	0.34	0.30	0.24
14	0.66	0.56	0.55	0.47	0.40	0.36
15	0.78	0.75	0.73	0.72	0.55	0.54
16	0.90	0.88	0.80	0.79	0.61	0.63
17	0.96	0.95	0.87	0.88	0.68	0.71
18	1.16	1.19	1.06	1.13	0.89	0.91
19	1.27	1.23	1.23	1.07	1.05	0.95
20	1.21	1.21	1.10	1.09	1.00	0.96
21	1.41	1.26	1.21	1.11	0.99	0.89
22	1.35	1.24	1.29	1.00	1.07	0.84
23	1.25	1.22	1.09	1.03	0.89	0.77
24	1.30	1.43	1.06	1.05	0.86	0.79
25	1.33	1.23	0.99	0.91	0.76	0.68
26-29	1.12	1.03	0.71	0.65	0.56	0.50
30-34	0.93	0.88	0.56	0.50	0.39	0.40
35-39	1.36	1.31	0.81	0.69	0.54	0.46
40-44	1.45	1.29	0.87	0.71	0.71	0.51
45-49	1.49	1.37	0.74	0.60	0.55	0.49
50-54	1.83	1.62	0.47	0.55	0.40	0.44
55-59	1.83	1.61	0.42	0.52	0.25	0.49
60-64	1.59	1.44	0.11	0.59	0.11	*
65 or Older	0.53	0.49	0.01	0.03	0.01	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.20P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana, by Detailed Age Categories: 2000 Versus 1999

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.4517		0.1936		0.6920	
12	0.8438		0.7943		0.4487	
13	0.8566		0.0664		0.0643	
14	0.0534		0.0731		0.1322	
15	0.2332		0.4552		0.3881	
16	0.4992		0.2204		0.7876	
17	0.7678		0.6607		0.1581	
18	0.4490		0.9870		0.6076	
19	0.1687		0.8876		0.9792	
20	0.7928		0.8051		0.4896	
21	0.8779		0.4622		0.9011	
22	0.8963		0.0938		0.1051	
23	0.0018		0.0404		0.0512	
24	0.1095		0.8331		0.5344	
25	0.7074		0.3204		0.3873	
26-29	0.6109		0.9069		0.8363	
30-34	0.0441		0.9489		0.1312	
35-39	0.0196		0.2830		0.9406	
40-44	0.5690		0.2200		0.2751	
45-49	0.8541		0.9485		0.1836	
50-54	0.1597		0.2740		0.8551	
55-59	0.1754		0.3804		0.1421	
60-64	0.7722		0.2472		*	
65 or Older	0.6402		0.5817		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.21C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	596	519	188	150	130	112
12	4	4	4	3	1	2
13	6	5	5	5	4	3
14	8	7	6	6	4	3
15	11	13	10	12	6	8
16	15	16	12	14	8	8
17	17	17	14	14	7	9
18	27	27	22	18	12	11
19	30	30	24	22	14	12
20	31	32	22	22	11	16
21	35	30	21	19	12	8
22	31	27	22	18	14	11
23	27	27	16	17	9	10
24	28	27	17	12	9	8
25	27	27	17	16	9	7
26-29	118	93	51	38	38	21
30-34	146	114	52	44	40	35
35-39	262	237	85	59	58	39
40-44	274	238	99	71	78	52
45-49	233	205	42	48	27	24
50-54	143	188	40	64	39	58
55-59	115	120	*	11	*	5
60-64	32	87	*	*	*	*
65 or Older	64	50	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.21D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.27	0.23	0.08	0.07	0.06	0.05
12	0.13	0.10	0.12	0.08	0.02	0.06
13	0.14	0.13	0.13	0.13	0.09	0.08
14	0.20	0.17	0.16	0.15	0.09	0.07
15	0.30	0.31	0.25	0.30	0.17	0.19
16	0.36	0.41	0.30	0.35	0.19	0.20
17	0.47	0.49	0.37	0.40	0.19	0.24
18	0.64	0.62	0.51	0.42	0.29	0.25
19	0.76	0.76	0.61	0.55	0.36	0.32
20	0.82	0.81	0.58	0.57	0.28	0.40
21	1.00	0.85	0.61	0.55	0.34	0.24
22	0.91	0.80	0.64	0.54	0.41	0.32
23	0.84	0.79	0.49	0.49	0.27	0.30
24	0.88	0.83	0.52	0.39	0.29	0.24
25	0.87	0.80	0.55	0.48	0.30	0.21
26-29	0.80	0.65	0.34	0.27	0.26	0.15
30-34	0.78	0.61	0.28	0.23	0.21	0.19
35-39	1.15	1.10	0.37	0.27	0.25	0.18
40-44	1.27	1.08	0.46	0.32	0.36	0.24
45-49	1.22	1.02	0.22	0.24	0.14	0.12
50-54	0.84	0.99	0.24	0.34	0.23	0.31
55-59	0.89	0.89	*	0.08	*	0.03
60-64	0.32	0.86	*	*	*	*
65 or Older	0.20	0.16	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.21P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine, by Detailed Age Categories: 2000 Versus 1999

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.3230		0.0660		0.0520	
12	0.9200		0.4956		0.3205	
13	0.4400		0.5357		0.4213	
14	0.9649		0.3967		0.6406	
15	0.9385		0.4135		0.9720	
16	0.8050		0.3463		0.7105	
17	0.8811		0.5967		0.4814	
18	0.3533		0.2776		0.7681	
19	0.2236		0.0466		0.3857	
20	0.8417		0.5683		0.4597	
21	0.5312		0.6541		0.0803	
22	0.4201		0.1531		0.0645	
23	0.0213		0.1241		0.5066	
24	0.4452		0.0960		0.4313	
25	0.5574		0.7083		0.6471	
26-29	0.0059		0.4821		0.1644	
30-34	0.0062		0.2009		0.1780	
35-39	0.7158		0.3995		0.4247	
40-44	0.7569		0.3775		0.3286	
45-49	0.5038		0.5373		0.7631	
50-54	0.0143		0.4046		0.8177	
55-59	0.1523		*		*	
60-64	0.0839		*		*	
65 or Older	0.9848		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.22C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	557	487	101	131	59	64
12	7	5	5	4	3	2
13	10	11	9	8	6	4
14	14	12	11	10	6	5
15	16	18	14	16	6	9
16	22	22	19	18	9	11
17	23	25	19	22	12	14
18	38	37	28	30	17	16
19	40	39	31	31	16	19
20	40	39	27	28	15	11
21	43	36	25	26	9	13
22	40	34	22	21	11	7
23	35	36	22	18	11	10
24	32	35	14	14	9	6
25	33	31	16	11	7	7
26-29	132	105	29	34	13	20
30-34	131	117	17	33	6	29
35-39	222	212	27	48	23	22
40-44	250	224	36	14	29	*
45-49	223	210	19	14	15	6
50-54	155	183	3	*	*	*
55-59	93	85	*	17	*	*
60-64	27	87	*	*	*	*
65 or Older	24	53	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.22D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.25	0.22	0.05	0.06	0.03	0.03
12	0.20	0.14	0.13	0.11	0.07	0.04
13	0.25	0.29	0.22	0.21	0.14	0.11
14	0.34	0.29	0.28	0.25	0.16	0.12
15	0.42	0.44	0.38	0.40	0.16	0.23
16	0.55	0.56	0.47	0.47	0.23	0.28
17	0.62	0.71	0.51	0.60	0.32	0.39
18	0.88	0.86	0.65	0.69	0.40	0.37
19	1.00	0.99	0.78	0.80	0.40	0.49
20	1.05	1.01	0.70	0.72	0.39	0.28
21	1.22	1.04	0.71	0.74	0.27	0.36
22	1.17	1.01	0.65	0.63	0.32	0.20
23	1.11	1.03	0.70	0.53	0.36	0.29
24	0.99	1.09	0.43	0.42	0.27	0.18
25	1.06	0.93	0.50	0.32	0.23	0.22
26-29	0.89	0.74	0.19	0.24	0.09	0.14
30-34	0.70	0.62	0.09	0.18	0.03	0.16
35-39	0.98	0.98	0.12	0.22	0.10	0.10
40-44	1.16	1.01	0.17	0.06	0.13	*
45-49	1.17	1.05	0.10	0.07	0.08	0.03
50-54	0.91	0.96	0.02	*	*	*
55-59	0.72	0.63	*	0.13	*	*
60-64	0.27	0.85	*	*	*	*
65 or Older	0.07	0.17	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.22P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens, by Detailed Age Categories: 2000 Versus 1999

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.2561		0.1092		0.6325	
12	0.2351		0.6113		0.5909	
13	0.9936		0.5693		0.6570	
14	0.2037		0.4886		0.4261	
15	0.0386		0.1111		0.3579	
16	0.6459		0.7025		0.3190	
17	0.7785		0.8477		0.8309	
18	0.6924		0.1375		0.6804	
19	0.9038		0.5459		0.1734	
20	0.0796		0.5896		0.0820	
21	0.5120		0.3014		0.1015	
22	0.4999		0.8388		0.0549	
23	0.0068		0.1374		0.2115	
24	0.6993		0.9974		0.6085	
25	0.7955		0.0817		0.7599	
26-29	0.1283		0.0383		0.2445	
30-34	0.8453		0.0137		0.0479	
35-39	0.0703		0.0461		0.9166	
40-44	0.6436		0.2224		*	
45-49	0.5091		0.6254		0.5545	
50-54	0.0353		*		*	
55-59	0.8932		*		*	
60-64	0.1083		*		*	
65 or Older	0.0959		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.23C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	443	381	88	90	52	67
12	16	16	13	11	8	7
13	21	20	15	14	8	5
14	24	22	16	15	8	9
15	21	21	15	15	7	6
16	21	21	14	14	6	8
17	24	20	15	13	7	6
18	34	29	20	18	9	8
19	34	33	23	19	14	10
20	32	33	17	19	4	8
21	35	30	17	13	7	5
22	34	31	14	10	5	4
23	29	29	9	10	1	6
24	31	28	8	9	5	5
25	31	26	13	9	10	4
26-29	112	93	13	19	8	8
30-34	114	99	24	24	23	22
35-39	187	170	30	18	21	17
40-44	205	155	31	26	17	22
45-49	177	141	32	35	17	34
50-54	133	135	7	19	*	15
55-59	78	100	*	22	*	22
60-64	66	41	3	*	3	*
65 or Older	22	56	3	22	3	16

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.23D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.20	0.17	0.04	0.04	0.02	0.03
12	0.45	0.42	0.37	0.28	0.23	0.17
13	0.52	0.50	0.36	0.34	0.20	0.12
14	0.60	0.55	0.40	0.36	0.20	0.22
15	0.55	0.53	0.39	0.36	0.17	0.16
16	0.52	0.53	0.34	0.34	0.16	0.20
17	0.65	0.57	0.40	0.36	0.18	0.16
18	0.80	0.67	0.46	0.42	0.21	0.19
19	0.86	0.85	0.58	0.50	0.35	0.26
20	0.85	0.86	0.45	0.49	0.11	0.20
21	0.99	0.85	0.47	0.38	0.18	0.14
22	1.01	0.90	0.42	0.29	0.14	0.12
23	0.91	0.84	0.27	0.29	0.04	0.16
24	0.95	0.87	0.25	0.27	0.16	0.14
25	0.99	0.77	0.41	0.28	0.33	0.13
26-29	0.75	0.65	0.09	0.13	0.05	0.06
30-34	0.61	0.53	0.13	0.13	0.12	0.12
35-39	0.82	0.79	0.13	0.09	0.09	0.08
40-44	0.95	0.70	0.14	0.12	0.08	0.10
45-49	0.93	0.70	0.17	0.17	0.09	0.17
50-54	0.78	0.71	0.04	0.10	*	0.08
55-59	0.60	0.74	*	0.16	*	0.16
60-64	0.65	0.40	0.03	*	0.03	*
65 or Older	0.07	0.18	0.01	0.07	0.01	0.05

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.23P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants, by Detailed Age Categories: 2000 Versus 1999

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.2871		0.7619		0.5823	
12	0.2747		0.0237		0.0954	
13	0.3438		0.4361		0.0059	
14	0.4417		0.4790		0.5955	
15	0.5130		0.7436		0.6170	
16	0.1644		0.7036		0.1508	
17	0.0094		0.1066		0.5288	
18	0.1092		0.7942		0.7661	
19	0.3672		0.8159		0.0950	
20	0.7829		0.6480		0.0743	
21	0.0756		0.1305		0.1624	
22	0.3253		0.0327		0.7374	
23	0.0976		0.4616		0.0042	
24	0.0337		0.8221		0.7713	
25	0.1542		0.4481		0.2657	
26-29	0.7915		0.1570		0.6766	
30-34	0.6297		0.3419		0.4907	
35-39	0.5940		0.5127		0.8482	
40-44	0.3869		0.8900		0.7547	
45-49	0.3665		0.9556		0.4874	
50-54	0.9633		0.2976		*	
55-59	0.1745		*		*	
60-64	0.8047		*		*	
65 or Older	0.0539		0.1260		0.4333	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.24C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	645	582	296	259	209	178
12	16	15	13	10	9	7
13	18	18	15	14	10	10
14	23	20	20	16	13	13
15	25	24	22	20	14	13
16	27	29	22	23	13	15
17	26	29	20	22	12	16
18	38	38	30	29	20	20
19	40	39	33	30	21	20
20	41	39	32	31	21	21
21	40	35	28	26	16	18
22	42	34	31	24	17	14
23	35	36	25	24	14	15
24	35	32	23	21	12	14
25	36	30	24	18	12	10
26-29	122	101	62	55	35	46
30-34	137	116	66	56	36	39
35-39	251	208	119	109	82	67
40-44	257	237	138	104	107	73
45-49	230	210	79	77	63	67
50-54	237	208	89	85	78	53
55-59	174	156	69	87	68	41
60-64	147	110	45	54	14	54
65 or Older	170	138	109	51	90	38

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.24D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.29	0.26	0.13	0.12	0.09	0.08
12	0.45	0.39	0.37	0.27	0.26	0.20
13	0.44	0.45	0.37	0.35	0.24	0.24
14	0.56	0.50	0.50	0.41	0.32	0.31
15	0.65	0.61	0.58	0.51	0.37	0.31
16	0.66	0.72	0.55	0.59	0.32	0.37
17	0.70	0.80	0.53	0.62	0.33	0.45
18	0.90	0.87	0.70	0.66	0.47	0.47
19	1.00	1.01	0.83	0.77	0.52	0.51
20	1.08	1.01	0.84	0.81	0.55	0.53
21	1.12	1.01	0.80	0.73	0.46	0.52
22	1.24	1.00	0.92	0.70	0.49	0.43
23	1.10	1.04	0.78	0.70	0.45	0.43
24	1.09	1.00	0.71	0.64	0.36	0.43
25	1.15	0.90	0.78	0.55	0.39	0.30
26-29	0.82	0.71	0.42	0.39	0.24	0.33
30-34	0.73	0.62	0.35	0.30	0.19	0.21
35-39	1.10	0.96	0.52	0.50	0.36	0.31
40-44	1.20	1.07	0.64	0.47	0.50	0.33
45-49	1.20	1.04	0.41	0.38	0.33	0.33
50-54	1.40	1.09	0.53	0.45	0.46	0.28
55-59	1.34	1.16	0.53	0.65	0.53	0.31
60-64	1.45	1.09	0.45	0.53	0.14	0.53
65 or Older	0.52	0.44	0.34	0.16	0.28	0.12

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.24P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic, by Detailed Age Categories: 2000 Versus 1999

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.0230		0.1617		0.6054	
12	0.1413		0.0291		0.2170	
13	0.3460		1.0000		0.9046	
14	0.0965		0.0546		0.7704	
15	0.8502		0.7332		0.2093	
16	0.8514		0.4041		0.3226	
17	0.1176		0.0467		0.0292	
18	0.3192		0.7330		0.6233	
19	0.2171		0.7315		0.6003	
20	0.2010		0.6635		0.4169	
21	0.5608		0.6233		0.9896	
22	0.0256		0.4020		0.6479	
23	0.1455		0.0714		0.5389	
24	0.6554		0.9958		0.5169	
25	0.0235		0.0179		0.1656	
26-29	0.4677		0.8738		0.0717	
30-34	0.0109		0.6922		0.0303	
35-39	0.6528		0.5073		0.1026	
40-44	0.4596		0.1036		0.1005	
45-49	0.4165		0.9167		0.5117	
50-54	0.6276		0.7640		0.9570	
55-59	0.4353		0.1007		0.8742	
60-64	0.5367		0.9700		0.1805	
65 or Older	0.1374		0.3130		0.3965	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.25C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	787	704	341	306	242	209
12	21	22	18	15	13	11
13	26	23	20	19	12	11
14	31	26	24	21	16	15
15	29	30	26	25	17	17
16	31	33	26	29	16	20
17	32	33	25	28	17	21
18	46	44	37	35	26	26
19	47	44	41	36	28	26
20	47	44	39	37	24	26
21	50	40	37	34	20	22
22	46	41	36	31	22	19
23	39	44	28	29	17	19
24	41	40	28	26	18	16
25	39	36	27	23	16	14
26-29	155	127	77	68	51	51
30-34	172	145	81	73	56	60
35-39	294	259	140	123	97	80
40-44	288	271	153	124	123	89
45-49	284	253	92	97	73	78
50-54	249	252	95	91	84	75
55-59	207	185	69	92	68	48
60-64	150	130	45	54	14	54
65 or Older	181	175	109	55	90	41

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.25D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana, by Detailed Age Categories: 1999 and 2000

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.36	0.32	0.15	0.14	0.11	0.09
12	0.60	0.58	0.50	0.40	0.35	0.28
13	0.64	0.57	0.49	0.47	0.31	0.26
14	0.75	0.65	0.58	0.52	0.39	0.38
15	0.77	0.74	0.68	0.63	0.45	0.41
16	0.77	0.83	0.66	0.73	0.39	0.49
17	0.85	0.92	0.68	0.77	0.45	0.57
18	1.08	1.02	0.87	0.82	0.61	0.61
19	1.19	1.12	1.02	0.94	0.69	0.66
20	1.23	1.15	1.04	0.95	0.64	0.68
21	1.41	1.13	1.05	0.97	0.57	0.63
22	1.37	1.21	1.06	0.92	0.66	0.56
23	1.22	1.26	0.89	0.83	0.55	0.56
24	1.26	1.25	0.86	0.80	0.55	0.50
25	1.26	1.09	0.85	0.70	0.52	0.41
26-29	1.05	0.90	0.52	0.48	0.35	0.36
30-34	0.91	0.77	0.43	0.39	0.30	0.32
35-39	1.29	1.20	0.61	0.57	0.43	0.37
40-44	1.34	1.22	0.71	0.56	0.57	0.40
45-49	1.49	1.26	0.48	0.48	0.38	0.39
50-54	1.47	1.32	0.56	0.48	0.49	0.39
55-59	1.59	1.37	0.53	0.68	0.53	0.36
60-64	1.48	1.28	0.45	0.53	0.14	0.53
65 or Older	0.56	0.56	0.34	0.18	0.28	0.13

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.25P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana, by Detailed Age Categories: 2000 Versus 1999

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
Total	0.2462		0.2589		0.3178	
12	0.0976		0.0067		0.0534	
13	0.6507		0.6033		0.3814	
14	0.0664		0.1125		0.8541	
15	0.4880		0.5815		0.4683	
16	0.3195		0.8638		0.1429	
17	0.8723		0.4473		0.0410	
18	0.4995		0.3603		0.3729	
19	0.3933		0.9119		0.6050	
20	0.9862		0.7136		0.7672	
21	0.6265		0.4976		0.4967	
22	0.1638		0.4154		0.2458	
23	0.0531		0.0275		0.4454	
24	0.0979		0.3446		0.4923	
25	0.1082		0.0573		0.4014	
26-29	0.4077		0.6463		0.4672	
30-34	0.0044		0.7388		0.3072	
35-39	0.7322		0.4582		0.0663	
40-44	0.6577		0.1531		0.0758	
45-49	0.2983		0.6861		0.6467	
50-54	0.3618		0.7975		0.9882	
55-59	0.1647		0.0408		0.7850	
60-64	1.0000		0.9935		0.2008	
65 or Older	0.3422		0.4835		0.4843	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.26C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	942	795	508	455	366	332
AGE						
12-17	83	86	75	73	53	56
18-25	154	150	137	132	115	105
26 or Older	898	751	451	398	332	300
GENDER						
Male	612	562	372	336	283	260
Female	666	512	310	286	218	211
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	812	668	424	399	310	292
Black Only	282	240	177	134	130	92
American Indian or Alaska Native Only	62	53	43	37	23	31
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	13
Asian Only	134	151	55	48	41	38
More Than One Race	77	71	36	68	31	68
Hispanic	290	247	146	121	110	83

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.26D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.43	0.36	0.23	0.20	0.17	0.15
AGE						
12-17	0.36	0.37	0.32	0.31	0.23	0.24
18-25	0.54	0.52	0.48	0.46	0.40	0.36
26 or Older	0.53	0.44	0.27	0.23	0.20	0.18
GENDER						
Male	0.58	0.52	0.35	0.31	0.27	0.24
Female	0.58	0.44	0.27	0.25	0.19	0.18
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.50	0.41	0.26	0.24	0.19	0.18
Black Only	1.14	0.94	0.71	0.53	0.52	0.36
American Indian or Alaska Native Only	5.35	4.82	3.74	3.40	2.03	2.84
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	2.35
Asian Only	1.85	1.98	0.76	0.63	0.56	0.50
More Than One Race	3.93	3.82	1.84	3.66	1.60	3.66
Hispanic	1.26	1.04	0.63	0.51	0.48	0.35

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.26P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.1639		0.0845		0.8949	
AGE						
12-17	0.1413		0.0119		0.8229	
18-25	0.0542		0.0571		0.3377	
26 or Older	0.3184		0.3454		0.6538	
GENDER						
Male	0.8644		0.0572		0.2780	
Female	0.0666		0.6957		0.1220	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.4164		0.5284		0.2544	
Black Only	0.1546		0.0076		0.0680	
American Indian or Alaska Native Only	0.6776		0.7652		0.5101	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.4979		0.3616		0.5604	
More Than One Race	0.2144		0.2150		0.2624	
Hispanic	0.4263		0.2576		0.1955	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.27C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	83	86	75	73	53	56
GENDER						
Male	55	61	54	51	41	38
Female	58	55	52	50	37	38
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	69	68	60	60	42	46
Black Only	31	32	28	24	22	18
American Indian or Alaska Native Only	6	*	6	*	5	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	18	14	15	12	12	9
More Than One Race	10	11	9	8	7	8
Hispanic	33	29	31	26	23	20
GENDER/RACE/HISPANIC ORIGIN						
Male - White	45	47	42	41	29	30
Female - White	47	44	42	40	31	32
Male - Black	23	25	21	19	16	14
Female - Black	21	20	20	16	15	12
Male - Hispanic	23	24	22	19	18	17
Female - Hispanic	22	19	20	17	13	14

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.27D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.36	0.37	0.32	0.31	0.23	0.24
GENDER						
Male	0.47	0.51	0.45	0.42	0.34	0.31
Female	0.51	0.48	0.46	0.44	0.33	0.34
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.45	0.44	0.39	0.39	0.27	0.30
Black Only	0.98	0.97	0.88	0.72	0.68	0.54
American Indian or Alaska Native Only	4.49	*	4.60	*	3.50	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.96	1.64	1.63	1.46	1.35	1.05
More Than One Race	2.70	2.90	2.44	2.19	1.90	2.08
Hispanic	1.02	0.88	0.96	0.77	0.73	0.61
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.58	0.60	0.53	0.53	0.37	0.38
Female - White	0.64	0.59	0.56	0.54	0.41	0.43
Male - Black	1.43	1.45	1.28	1.13	0.97	0.82
Female - Black	1.32	1.23	1.25	0.97	0.95	0.71
Male - Hispanic	1.39	1.37	1.36	1.11	1.06	1.02
Female - Hispanic	1.39	1.19	1.25	1.05	0.84	0.85

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.27P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 to 17, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.1413		0.0119		0.8229	
GENDER						
Male	0.0956		0.0175		0.6434	
Female	0.6600		0.2487		0.8762	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.7039		0.5132		0.4158	
Black Only	0.3115		0.0110		0.2886	
American Indian or Alaska Native Only	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.4583		0.5084		0.4450	
More Than One Race	0.6530		0.1531		0.9749	
Hispanic	0.0707		0.0348		0.3571	
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.3587		0.4609		0.2452	
Female - White	0.7166		0.8303		0.9713	
Male - Black	0.1741		0.0450		0.0951	
Female - Black	0.9968		0.1253		0.8108	
Male - Hispanic	0.3488		0.0877		0.3092	
Female - Hispanic	0.0687		0.1747		0.7968	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.28C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	154	150	137	132	115	105
GENDER						
Male	97	103	92	89	82	75
Female	100	95	83	83	63	62
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	120	112	112	106	93	88
Black Only	52	46	46	41	41	31
American Indian or Alaska Native Only	7	9	5	*	5	7
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	26	24	23	16	16	12
More Than One Race	13	12	13	13	12	12
Hispanic	53	55	45	44	39	36
ADULT EDUCATION						
< High School	65	61	56	54	50	47
High School Graduate	84	74	76	72	61	56
Some College	85	77	82	76	64	61
College Graduate	44	52	42	41	32	27
CURRENT EMPLOYMENT						
Full-Time	96	106	88	92	77	71
Part-Time	70	67	63	59	54	51
Unemployed	35	29	31	28	27	26
Other ¹	61	57	57	52	42	39

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.28D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.54	0.52	0.48	0.46	0.40	0.36
GENDER						
Male	0.69	0.71	0.65	0.62	0.58	0.52
Female	0.70	0.65	0.58	0.57	0.44	0.42
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.64	0.59	0.60	0.55	0.49	0.46
Black Only	1.35	1.16	1.20	1.06	1.05	0.79
American Indian or Alaska Native Only	5.18	4.82	3.95	*	3.44	3.98
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	2.40	2.41	2.10	1.61	1.52	1.22
More Than One Race	3.82	4.00	3.70	4.20	3.42	3.88
Hispanic	1.26	1.26	1.07	1.02	0.92	0.82
ADULT EDUCATION						
< High School	1.11	1.00	0.96	0.89	0.86	0.77
High School Graduate	0.82	0.71	0.74	0.69	0.59	0.54
Some College	0.93	0.85	0.90	0.84	0.71	0.68
College Graduate	1.35	1.52	1.30	1.20	0.97	0.78
CURRENT EMPLOYMENT						
Full-Time	0.65	0.69	0.60	0.60	0.52	0.46
Part-Time	1.02	1.00	0.92	0.88	0.79	0.76
Unemployed	2.12	1.78	1.88	1.72	1.64	1.61
Other ¹	1.16	1.07	1.07	0.97	0.80	0.74

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.28P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 to 25, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0542		0.0571		0.3377	
GENDER						
Male	0.1371		0.0857		0.1197	
Female	0.1587		0.2780		0.6868	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.0945		0.1425		0.9754	
Black Only	0.5561		0.1939		0.3652	
American Indian or Alaska Native Only	0.0093		*		0.9900	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.3628		0.0953		0.6578	
More Than One Race	0.9071		0.1906		0.4620	
Hispanic	0.1880		0.5197		0.0440	
ADULT EDUCATION						
< High School	0.5078		0.5462		0.7625	
High School Graduate	0.3071		0.1393		0.1818	
Some College	0.4222		0.5343		0.9921	
College Graduate	0.0514		0.1636		0.7999	
CURRENT EMPLOYMENT						
Full-Time	0.0877		0.2460		0.4469	
Part-Time	0.2773		0.0231		0.0782	
Unemployed	0.2206		0.3174		0.2760	
Other ¹	0.1435		0.6640		0.6070	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.29C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	898	751	451	398	332	300
GENDER						
Male	600	542	339	301	261	235
Female	631	491	274	256	202	197
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	778	641	385	354	284	264
Black Only	268	230	158	118	119	84
American Indian or Alaska Native Only	*	*	*	26	23	25
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	128	147	43	43	32	36
More Than One Race	73	68	28	*	27	*
Hispanic	268	229	124	106	99	70
ADULT EDUCATION						
< High School	334	289	176	165	139	129
High School Graduate	521	455	223	218	165	185
Some College	413	384	235	175	178	130
College Graduate	448	415	245	216	200	153
CURRENT EMPLOYMENT						
Full-Time	661	587	370	317	269	243
Part-Time	276	229	119	106	90	91
Unemployed	122	103	93	62	79	55
Other ¹	390	339	191	177	152	133

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.29D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.53	0.44	0.27	0.23	0.20	0.18
GENDER						
Male	0.75	0.67	0.42	0.37	0.32	0.29
Female	0.71	0.55	0.31	0.28	0.23	0.22
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.61	0.50	0.30	0.28	0.22	0.21
Black Only	1.51	1.27	0.89	0.65	0.67	0.46
American Indian or Alaska Native Only	*	*	*	3.43	2.58	3.20
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	2.43	2.55	0.81	0.74	0.61	0.62
More Than One Race	5.90	5.73	2.30	*	2.17	*
Hispanic	1.72	1.41	0.79	0.65	0.63	0.43
ADULT EDUCATION						
< High School	1.09	0.99	0.57	0.56	0.45	0.44
High School Graduate	0.90	0.80	0.39	0.38	0.29	0.33
Some College	1.06	0.98	0.60	0.45	0.46	0.33
College Graduate	1.06	0.91	0.58	0.47	0.47	0.34
CURRENT EMPLOYMENT						
Full-Time	0.67	0.58	0.38	0.32	0.27	0.24
Part-Time	1.63	1.37	0.70	0.63	0.53	0.54
Unemployed	3.48	3.38	2.64	2.03	2.25	1.81
Other ¹	0.77	0.67	0.38	0.35	0.30	0.26

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.29P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 26 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3184		0.3454		0.6538	
GENDER						
Male	0.8120		0.2072		0.5269	
Female	0.0951		0.9369		0.1437	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.5576		0.7930		0.3247	
Black Only	0.2122		0.0394		0.1568	
American Indian or Alaska Native Only	*		*		0.7016	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.6958		0.8207		0.9947	
More Than One Race	0.1395		*		*	
Hispanic	0.7513		0.6088		0.7153	
ADULT EDUCATION						
< High School	0.3093		0.2518		0.2744	
High School Graduate	0.6013		0.3935		0.0092	
Some College	0.0524		0.0792		0.4699	
College Graduate	0.5368		0.8840		0.8003	
CURRENT EMPLOYMENT						
Full-Time	0.2790		0.4038		0.5716	
Part-Time	0.8843		0.6871		0.8084	
Unemployed	0.6720		0.1553		0.2851	
Other ¹	0.4581		0.8487		0.7236	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.30C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	927	785	492	442	360	322
GENDER						
Male	610	557	364	331	278	256
Female	657	508	297	276	213	204
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	802	662	415	388	307	284
Black Only	279	236	169	132	126	91
American Indian or Alaska Native Only	*	53	44	31	23	26
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	12
Asian Only	132	149	51	46	37	38
More Than One Race	76	70	34	67	31	*
Hispanic	289	242	141	118	108	79
ADULT EDUCATION						
< High School	349	305	192	178	152	141
High School Graduate	539	460	248	233	177	195
Some College	426	390	271	209	204	150
College Graduate	448	422	248	224	200	155
CURRENT EMPLOYMENT						
Full-Time	669	599	383	337	282	252
Part-Time	283	241	146	131	110	105
Unemployed	128	106	101	73	85	63
Other ¹	407	348	212	190	164	141

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.30D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.47	0.39	0.25	0.22	0.18	0.16
GENDER						
Male	0.65	0.58	0.39	0.35	0.29	0.27
Female	0.63	0.49	0.29	0.26	0.21	0.19
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.55	0.45	0.28	0.26	0.21	0.19
Black Only	1.29	1.07	0.78	0.60	0.58	0.41
American Indian or Alaska Native Only	*	5.55	4.27	3.23	2.27	2.72
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	2.59
Asian Only	2.08	2.20	0.81	0.68	0.58	0.56
More Than One Race	4.80	4.68	2.16	4.49	1.96	*
Hispanic	1.46	1.18	0.71	0.58	0.55	0.39
ADULT EDUCATION						
< High School	0.96	0.86	0.53	0.50	0.42	0.40
High School Graduate	0.79	0.68	0.37	0.35	0.26	0.29
Some College	0.89	0.81	0.56	0.43	0.42	0.31
College Graduate	0.98	0.86	0.54	0.46	0.44	0.32
CURRENT EMPLOYMENT						
Full-Time	0.59	0.52	0.34	0.29	0.25	0.22
Part-Time	1.19	1.03	0.61	0.56	0.46	0.45
Unemployed	2.48	2.26	1.95	1.57	1.65	1.34
Other ¹	0.73	0.62	0.38	0.34	0.29	0.25

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.30P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2051		0.1755		0.8582	
GENDER						
Male	0.9876		0.1131		0.2971	
Female	0.0699		0.8544		0.1177	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.4263		0.5975		0.2951	
Black Only	0.1886		0.0205		0.0964	
American Indian or Alaska Native Only	*		0.9891		0.6142	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.5354		0.4914		0.7436	
More Than One Race	0.1793		0.1388		*	
Hispanic	0.5601		0.4959		0.2653	
ADULT EDUCATION						
< High School	0.4103		0.3955		0.4238	
High School Graduate	0.6782		0.6126		0.0275	
Some College	0.0385		0.0627		0.4785	
College Graduate	0.4015		0.6837		0.7501	
CURRENT EMPLOYMENT						
Full-Time	0.1954		0.3131		0.6829	
Part-Time	0.8802		0.1707		0.5319	
Unemployed	0.9502		0.4673		0.6807	
Other ¹	0.3503		0.9153		0.6152	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.31C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	872	775	428	398	295	281
AGE						
12-17	74	72	66	62	47	49
18-25	149	150	131	126	109	98
26 or Older	841	734	379	339	261	248
GENDER						
Male	582	549	331	297	250	230
Female	609	509	236	219	147	162
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	758	656	357	347	250	248
Black Only	277	231	149	118	111	88
American Indian or Alaska Native Only	45	52	26	32	17	28
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	5
Asian Only	106	99	45	26	30	20
More Than One Race	77	74	34	68	29	68
Hispanic	246	229	115	87	71	68

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.31D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.39	0.35	0.19	0.18	0.13	0.13
AGE						
12-17	0.32	0.31	0.29	0.27	0.20	0.21
18-25	0.52	0.52	0.46	0.43	0.38	0.34
26 or Older	0.50	0.43	0.22	0.20	0.15	0.15
GENDER						
Male	0.55	0.51	0.31	0.28	0.23	0.21
Female	0.53	0.44	0.21	0.19	0.13	0.14
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.47	0.40	0.22	0.21	0.15	0.15
Black Only	1.12	0.91	0.60	0.46	0.45	0.35
American Indian or Alaska Native Only	3.91	4.74	2.23	2.96	1.52	2.53
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	0.99
Asian Only	1.46	1.30	0.62	0.34	0.42	0.26
More Than One Race	3.93	3.97	1.75	3.65	1.49	3.63
Hispanic	1.07	0.96	0.50	0.36	0.31	0.29

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.31P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.4517		0.1936		0.6920	
AGE						
12-17	0.3569		0.0503		0.8917	
18-25	0.1309		0.1937		0.2250	
26 or Older	0.6525		0.4785		0.3787	
GENDER						
Male	0.9721		0.0748		0.2695	
Female	0.2493		0.7753		0.0177	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.8750		0.9213		0.1774	
Black Only	0.3720		0.0578		0.2409	
American Indian or Alaska Native Only	0.6070		0.4227		0.2710	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.1589		0.0083		0.0605	
More Than One Race	0.2853		0.2727		0.3090	
Hispanic	0.4106		0.0638		0.1744	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.32C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	74	72	66	62	47	49
GENDER						
Male	52	51	49	44	37	35
Female	50	48	44	43	32	32
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	59	60	52	53	37	41
Black Only	29	25	26	19	21	15
American Indian or Alaska Native Only	6	*	6	*	4	7
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	16	11	13	10	12	6
More Than One Race	9	9	9	7	7	7
Hispanic	30	27	27	23	18	19
GENDER/RACE/HISPANIC ORIGIN						
Male - White	42	40	37	36	26	28
Female - White	41	40	36	36	27	27
Male - Black	21	19	19	15	15	12
Female - Black	18	16	17	12	14	9
Male - Hispanic	23	21	21	17	15	16
Female - Hispanic	19	17	16	16	10	11

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.32D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.32	0.31	0.29	0.27	0.20	0.21
GENDER						
Male	0.44	0.42	0.41	0.36	0.31	0.29
Female	0.44	0.42	0.39	0.38	0.28	0.28
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.39	0.39	0.34	0.34	0.24	0.27
Black Only	0.90	0.74	0.80	0.58	0.64	0.45
American Indian or Alaska Native Only	4.68	*	4.80	*	3.00	4.54
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.77	1.26	1.47	1.18	1.29	0.78
More Than One Race	2.52	2.53	2.37	2.00	1.80	1.97
Hispanic	0.92	0.80	0.83	0.69	0.57	0.58
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.53	0.50	0.47	0.46	0.33	0.35
Female - White	0.55	0.54	0.49	0.48	0.36	0.37
Male - Black	1.29	1.09	1.17	0.90	0.91	0.72
Female - Black	1.15	0.99	1.05	0.72	0.86	0.53
Male - Hispanic	1.39	1.23	1.27	1.01	0.92	0.93
Female - Hispanic	1.23	1.06	1.03	0.98	0.63	0.72

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.32P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3569		0.0503		0.8917	
GENDER						
Male	0.0980		0.0150		0.9758	
Female	0.7135		0.8123		0.8685	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.6032		0.9409		0.2544	
Black Only	0.1638		0.0056		0.1272	
American Indian or Alaska Native Only	*		*		0.2453	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.0222		0.0640		0.0314	
More Than One Race	0.4408		0.3167		0.9249	
Hispanic	0.1708		0.0820		0.7281	
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.8723		0.9555		0.0897	
Female - White	0.3660		0.8712		0.9498	
Male - Black	0.1410		0.0256		0.2752	
Female - Black	0.6618		0.0930		0.2754	
Male - Hispanic	0.2369		0.0256		0.2829	
Female - Hispanic	0.5138		0.9802		0.3733	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.33C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	149	150	131	126	109	98
GENDER						
Male	95	98	90	84	79	69
Female	98	97	77	78	59	58
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	120	114	107	101	90	82
Black Only	52	44	44	37	39	30
American Indian or Alaska Native Only	7	9	5	*	4	7
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	3
Asian Only	25	21	20	14	15	11
More Than One Race	14	12	12	12	11	10
Hispanic	51	55	40	40	34	31
ADULT EDUCATION						
< High School	64	59	52	52	46	43
High School Graduate	83	75	72	66	58	53
Some College	87	76	81	72	63	56
College Graduate	45	52	39	38	31	24
CURRENT EMPLOYMENT						
Full-Time	95	103	84	86	73	63
Part-Time	69	68	60	56	53	49
Unemployed	33	29	29	26	25	24
Other ¹	59	55	51	48	41	37

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.33D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.52	0.52	0.46	0.43	0.38	0.34
GENDER						
Male	0.67	0.68	0.64	0.58	0.56	0.48
Female	0.69	0.66	0.54	0.53	0.41	0.40
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.64	0.60	0.57	0.53	0.48	0.43
Black Only	1.34	1.12	1.14	0.93	1.00	0.76
American Indian or Alaska Native Only	4.84	5.01	3.71	*	3.19	3.86
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	2.60
Asian Only	2.28	2.05	1.80	1.40	1.41	1.09
More Than One Race	3.96	4.03	3.48	3.89	3.17	3.44
Hispanic	1.22	1.26	0.94	0.92	0.81	0.73
ADULT EDUCATION						
< High School	1.09	0.96	0.89	0.84	0.78	0.71
High School Graduate	0.81	0.72	0.70	0.63	0.56	0.51
Some College	0.95	0.84	0.89	0.79	0.69	0.62
College Graduate	1.39	1.53	1.19	1.11	0.94	0.71
CURRENT EMPLOYMENT						
Full-Time	0.64	0.67	0.57	0.56	0.49	0.41
Part-Time	1.01	1.01	0.89	0.84	0.77	0.73
Unemployed	2.01	1.80	1.72	1.63	1.53	1.50
Other ¹	1.12	1.03	0.96	0.89	0.78	0.69

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.33P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.1309		0.1937		0.2250	
GENDER						
Male	0.1125		0.0998		0.0991	
Female	0.5005		0.8282		0.9474	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.1628		0.4599		0.8712	
Black Only	0.5407		0.1752		0.4083	
American Indian or Alaska Native Only	0.0036		*		0.9963	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.1533		0.0430		0.7171	
More Than One Race	0.8285		0.2629		0.4864	
Hispanic	0.5206		0.6182		0.0156	
ADULT EDUCATION						
< High School	0.5554		0.8185		0.8795	
High School Graduate	0.2642		0.2904		0.2097	
Some College	0.8466		0.7328		0.4399	
College Graduate	0.0307		0.1832		0.6103	
CURRENT EMPLOYMENT						
Full-Time	0.1527		0.5030		0.2842	
Part-Time	0.3109		0.0427		0.0908	
Unemployed	0.2369		0.2683		0.1678	
Other ¹	0.2781		0.8693		0.9421	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.34C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	841	734	379	339	261	248
GENDER						
Male	572	530	302	266	223	206
Female	582	489	207	190	129	150
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	737	630	325	304	226	218
Black Only	267	226	133	102	100	80
American Indian or Alaska Native Only	46	*	24	23	16	22
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	102	97	35	18	22	14
More Than One Race	*	71	28	*	25	*
Hispanic	230	209	96	67	54	57
ADULT EDUCATION						
< High School	292	272	147	143	109	107
High School Graduate	491	439	204	178	134	148
Some College	410	372	201	144	146	110
College Graduate	425	400	195	187	145	134
CURRENT EMPLOYMENT						
Full-Time	641	573	321	277	232	201
Part-Time	248	217	92	84	70	66
Unemployed	120	104	87	57	61	52
Other ¹	333	305	137	131	100	109

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.34D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.50	0.43	0.22	0.20	0.15	0.15
GENDER						
Male	0.71	0.65	0.38	0.33	0.28	0.25
Female	0.65	0.54	0.23	0.21	0.14	0.17
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.58	0.49	0.25	0.24	0.18	0.17
Black Only	1.50	1.24	0.75	0.56	0.56	0.44
American Indian or Alaska Native Only	5.19	*	2.77	3.00	1.82	2.86
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.93	1.67	0.67	0.30	0.41	0.24
More Than One Race	*	5.92	2.26	*	2.06	*
Hispanic	1.48	1.29	0.62	0.41	0.35	0.35
ADULT EDUCATION						
< High School	0.95	0.93	0.48	0.49	0.35	0.36
High School Graduate	0.85	0.77	0.35	0.31	0.23	0.26
Some College	1.06	0.95	0.52	0.37	0.38	0.28
College Graduate	1.00	0.88	0.46	0.41	0.34	0.29
CURRENT EMPLOYMENT						
Full-Time	0.65	0.57	0.33	0.27	0.24	0.20
Part-Time	1.47	1.30	0.54	0.50	0.41	0.40
Unemployed	3.42	3.39	2.47	1.86	1.74	1.71
Other ¹	0.66	0.60	0.27	0.26	0.20	0.22

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.34P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.6525		0.4785		0.3787	
GENDER						
Male	0.6554		0.2697		0.5021	
Female	0.2539		0.6903		0.0096	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.9796		1.0000		0.2246	
Black Only	0.5210		0.2913		0.5352	
American Indian or Alaska Native Only	*		0.8647		0.4387	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.4431		0.3310		0.3951	
More Than One Race	*		*		*	
Hispanic	0.5860		0.1505		0.8847	
ADULT EDUCATION						
< High School	0.7414		0.1507		0.2537	
High School Graduate	0.5703		0.9685		0.0253	
Some College	0.0812		0.2400		0.7911	
College Graduate	0.7082		0.4014		0.6181	
CURRENT EMPLOYMENT						
Full-Time	0.2683		0.5426		0.6458	
Part-Time	0.4962		0.8822		0.6191	
Unemployed	0.5108		0.1477		0.9255	
Other ¹	0.8058		0.8912		0.3575	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.35C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	868	771	419	387	290	273
GENDER						
Male	584	546	327	293	246	225
Female	605	507	227	208	142	157
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	756	651	351	337	248	241
Black Only	275	228	144	116	108	87
American Indian or Alaska Native Only	47	49	25	27	17	23
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	5
Asian Only	105	98	42	23	27	18
More Than One Race	76	73	33	67	29	*
Hispanic	248	226	111	83	67	65
ADULT EDUCATION						
< High School	309	290	162	156	120	117
High School Graduate	506	446	225	198	147	160
Some College	420	379	238	178	174	130
College Graduate	426	409	200	196	148	136
CURRENT EMPLOYMENT						
Full-Time	648	585	336	300	248	210
Part-Time	259	232	125	111	96	88
Unemployed	126	107	94	65	68	58
Other ¹	350	315	155	142	110	118

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.35D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.44	0.39	0.21	0.19	0.15	0.14
GENDER						
Male	0.62	0.57	0.35	0.31	0.26	0.24
Female	0.58	0.49	0.22	0.20	0.14	0.15
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.51	0.44	0.24	0.23	0.17	0.16
Black Only	1.27	1.04	0.67	0.52	0.50	0.39
American Indian or Alaska Native Only	4.62	5.19	2.44	2.80	1.64	2.47
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	0.97
Asian Only	1.65	1.45	0.67	0.34	0.43	0.27
More Than One Race	4.80	4.85	2.07	4.49	1.83	*
Hispanic	1.26	1.10	0.56	0.40	0.34	0.32
ADULT EDUCATION						
< High School	0.85	0.82	0.44	0.44	0.33	0.33
High School Graduate	0.75	0.66	0.33	0.29	0.22	0.24
Some College	0.88	0.79	0.50	0.37	0.36	0.27
College Graduate	0.93	0.83	0.44	0.40	0.32	0.28
CURRENT EMPLOYMENT						
Full-Time	0.57	0.50	0.30	0.26	0.22	0.18
Part-Time	1.09	0.99	0.53	0.47	0.40	0.38
Unemployed	2.43	2.29	1.81	1.40	1.32	1.24
Other ¹	0.63	0.56	0.28	0.25	0.20	0.21

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.35P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.4973		0.3217		0.6625	
GENDER						
Male	0.8658		0.1480		0.2590	
Female	0.2318		0.7256		0.0123	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.8266		0.9197		0.2319	
Black Only	0.4622		0.1531		0.3683	
American Indian or Alaska Native Only	0.7631		0.7318		0.3674	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.2403		0.0360		0.2639	
More Than One Race	0.2275		0.2058		*	
Hispanic	0.5083		0.1485		0.1844	
ADULT EDUCATION						
< High School	0.8765		0.3229		0.5204	
High School Graduate	0.6652		0.8855		0.0783	
Some College	0.0930		0.2205		0.5535	
College Graduate	0.5167		0.6084		0.7132	
CURRENT EMPLOYMENT						
Full-Time	0.2012		0.5065		0.8362	
Part-Time	0.7004		0.3567		0.5968	
Unemployed	0.7689		0.4754		0.4834	
Other ¹	0.9416		0.8998		0.3619	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.36C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	596	519	188	150	130	112
AGE						
12-17	29	31	24	27	14	17
18-25	93	83	60	52	35	32
26 or Older	580	505	170	135	123	102
GENDER						
Male	448	409	148	113	105	78
Female	356	323	106	99	71	74
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	523	461	153	129	108	91
Black Only	192	139	65	55	56	45
American Indian or Alaska Native Only	36	48	13	24	7	8
Native Hawaiian or Other Pacific Islander	*	*	3	*	1	*
Asian Only	36	33	13	6	8	4
More Than One Race	45	60	16	14	6	12
Hispanic	164	158	72	42	41	33

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.36D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.27	0.23	0.08	0.07	0.06	0.05
AGE						
12-17	0.13	0.13	0.10	0.12	0.06	0.07
18-25	0.33	0.29	0.21	0.18	0.12	0.11
26 or Older	0.34	0.30	0.10	0.08	0.07	0.06
GENDER						
Male	0.42	0.38	0.14	0.11	0.10	0.07
Female	0.31	0.28	0.09	0.09	0.06	0.06
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.32	0.28	0.09	0.08	0.07	0.06
Black Only	0.77	0.55	0.26	0.22	0.22	0.18
American Indian or Alaska Native Only	3.14	4.36	1.12	2.23	0.57	0.74
Native Hawaiian or Other Pacific Islander	*	*	0.41	*	0.12	*
Asian Only	0.50	0.44	0.18	0.08	0.11	0.05
More Than One Race	2.32	3.24	0.82	0.77	0.32	0.65
Hispanic	0.71	0.66	0.31	0.18	0.18	0.14

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.36P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3230		0.0660		0.0520	
AGE						
12-17	0.9117		0.7337		0.7400	
18-25	0.0204		0.0022		0.0552	
26 or Older	0.5398		0.3156		0.1003	
GENDER						
Male	0.5455		0.1980		0.0984	
Female	0.4124		0.1333		0.1994	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.8801		0.2093		0.0314	
Black Only	0.0217		0.6794		0.5205	
American Indian or Alaska Native Only	0.6824		0.6839		0.6230	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.7141		0.0839		0.5453	
More Than One Race	0.2950		0.4799		0.5187	
Hispanic	0.6435		0.0967		0.9895	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.37C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	29	31	24	27	14	17
GENDER						
Male	21	20	16	18	10	12
Female	20	22	17	18	10	10
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	23	26	20	23	12	15
Black Only	4	5	2	4	1	2
American Indian or Alaska Native Only	*	*	*	*	1	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	2	2	1	1	1	1
More Than One Race	4	4	2	4	1	1
Hispanic	15	13	12	11	6	5
GENDER/RACE/HISPANIC ORIGIN						
Male - White	16	18	13	16	8	11
Female - White	17	17	16	14	9	9
Male - Black	3	3	2	3	1	2
Female - Black	2	3	0	3	*	1
Male - Hispanic	12	9	10	8	5	4
Female - Hispanic	8	9	6	7	3	3

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.37D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.13	0.13	0.10	0.12	0.06	0.07
GENDER						
Male	0.17	0.17	0.14	0.15	0.08	0.10
Female	0.17	0.19	0.15	0.16	0.09	0.09
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.15	0.17	0.13	0.15	0.08	0.10
Black Only	0.13	0.14	0.07	0.13	0.04	0.06
American Indian or Alaska Native Only	*	*	*	*	0.96	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	0.23	0.21	0.16	0.11	0.06	0.09
More Than One Race	0.96	1.11	0.57	1.09	0.15	0.27
Hispanic	0.46	0.40	0.36	0.33	0.19	0.15
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.20	0.23	0.16	0.21	0.10	0.14
Female - White	0.23	0.23	0.21	0.19	0.12	0.12
Male - Black	0.21	0.20	0.13	0.19	0.07	0.09
Female - Black	0.14	0.18	0.02	0.17	*	0.06
Male - Hispanic	0.70	0.51	0.60	0.46	0.32	0.25
Female - Hispanic	0.52	0.60	0.38	0.46	0.19	0.17

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.37P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 to 17, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.9117		0.7337		0.7400	
GENDER						
Male	0.6278		0.3047		0.4438	
Female	0.5273		0.6083		0.7627	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.8420		0.9012		0.9272	
Black Only	0.6100		0.1006		0.3761	
American Indian or Alaska Native Only	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.9774		0.8803		0.5577	
More Than One Race	0.4534		0.6887		0.8145	
Hispanic	0.8673		0.8844		0.9859	
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.2799		0.2316		0.5662	
Female - White	0.4247		0.3009		0.6355	
Male - Black	0.9581		0.4420		0.6410	
Female - Black	0.4354		0.0845		*	
Male - Hispanic	0.5264		0.7567		0.7556	
Female - Hispanic	0.6645		0.5327		0.5639	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.38C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	93	83	60	52	35	32
GENDER						
Male	71	63	49	39	27	24
Female	50	54	33	34	20	19
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	80	69	52	44	30	23
Black Only	19	17	14	11	10	5
American Indian or Alaska Native Only	4	*	2	*	1	1
Native Hawaiian or Other Pacific Islander	*	3	*	1	*	*
Asian Only	14	8	12	5	8	3
More Than One Race	10	8	9	5	6	2
Hispanic	37	31	24	22	13	19
ADULT EDUCATION						
< High School	45	42	31	26	18	19
High School Graduate	51	50	36	34	22	18
Some College	49	47	33	30	18	14
College Graduate	22	19	16	13	11	8
CURRENT EMPLOYMENT						
Full-Time	66	61	42	40	23	23
Part-Time	41	36	28	25	16	13
Unemployed	21	22	16	16	12	10
Other ¹	33	30	23	17	15	9

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.38D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.33	0.29	0.21	0.18	0.12	0.11
GENDER						
Male	0.50	0.44	0.34	0.27	0.19	0.16
Female	0.35	0.37	0.23	0.23	0.14	0.13
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.42	0.36	0.28	0.23	0.16	0.12
Black Only	0.49	0.44	0.38	0.27	0.26	0.12
American Indian or Alaska Native Only	2.95	*	1.46	*	0.41	0.35
Native Hawaiian or Other Pacific Islander	*	2.65	*	1.15	*	*
Asian Only	1.27	0.78	1.07	0.55	0.75	0.34
More Than One Race	3.01	2.80	2.47	1.73	1.70	0.82
Hispanic	0.89	0.73	0.57	0.51	0.32	0.44
ADULT EDUCATION						
< High School	0.77	0.69	0.53	0.43	0.30	0.30
High School Graduate	0.50	0.48	0.36	0.32	0.21	0.17
Some College	0.54	0.52	0.36	0.33	0.19	0.15
College Graduate	0.67	0.56	0.48	0.37	0.33	0.23
CURRENT EMPLOYMENT						
Full-Time	0.45	0.40	0.29	0.26	0.15	0.15
Part-Time	0.60	0.54	0.42	0.38	0.23	0.20
Unemployed	1.26	1.36	0.96	1.00	0.70	0.62
Other ¹	0.63	0.56	0.43	0.32	0.27	0.17

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.38P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 to 25, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0204		0.0022		0.0552	
GENDER						
Male	0.0344		0.0045		0.0732	
Female	0.2470		0.1979		0.3729	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.2693		0.0732		0.0567	
Black Only	0.3070		0.2596		0.1652	
American Indian or Alaska Native Only	*		*		0.9710	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.1515		0.2126		0.4775	
More Than One Race	0.4733		0.1008		0.1919	
Hispanic	0.0091		0.0275		0.5092	
ADULT EDUCATION						
< High School	0.2482		0.1521		0.6133	
High School Graduate	0.0546		0.0205		0.1193	
Some College	0.3146		0.2377		0.2069	
College Graduate	0.5615		0.4320		0.6866	
CURRENT EMPLOYMENT						
Full-Time	0.0262		0.0128		0.1474	
Part-Time	0.2520		0.2378		0.7022	
Unemployed	0.6623		0.7552		0.8858	
Other ¹	0.2259		0.0412		0.0680	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.39C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	580	505	170	135	123	102
GENDER						
Male	435	396	136	103	101	73
Female	349	321	98	88	70	68
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	507	454	140	118	102	86
Black Only	190	138	61	54	53	44
American Indian or Alaska Native Only	36	*	12	*	6	7
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	33	32	4	1	*	1
More Than One Race	43	58	13	13	*	12
Hispanic	152	149	65	32	38	25
ADULT EDUCATION						
< High School	185	184	94	66	75	60
High School Graduate	297	281	87	78	68	48
Some College	307	287	64	53	46	32
College Graduate	321	258	69	69	42	54
CURRENT EMPLOYMENT						
Full-Time	482	440	130	107	105	73
Part-Time	158	147	53	62	29	54
Unemployed	101	89	58	23	44	16
Other ¹	196	195	51	48	28	42

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.39D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.34	0.30	0.10	0.08	0.07	0.06
GENDER						
Male	0.54	0.49	0.17	0.13	0.13	0.09
Female	0.39	0.36	0.11	0.10	0.08	0.08
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.40	0.35	0.11	0.09	0.08	0.07
Black Only	1.07	0.76	0.34	0.30	0.30	0.25
American Indian or Alaska Native Only	4.11	*	1.36	*	0.70	0.88
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	0.62	0.55	0.08	0.02	*	0.02
More Than One Race	3.52	4.85	1.03	1.05	*	1.00
Hispanic	0.97	0.92	0.42	0.20	0.24	0.16
ADULT EDUCATION						
< High School	0.60	0.63	0.31	0.22	0.24	0.21
High School Graduate	0.52	0.49	0.15	0.14	0.12	0.09
Some College	0.79	0.73	0.17	0.14	0.12	0.08
College Graduate	0.76	0.56	0.16	0.15	0.10	0.12
CURRENT EMPLOYMENT						
Full-Time	0.49	0.44	0.13	0.11	0.11	0.07
Part-Time	0.93	0.88	0.32	0.37	0.17	0.32
Unemployed	2.87	2.93	1.65	0.76	1.25	0.52
Other ¹	0.39	0.39	0.10	0.09	0.06	0.08

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.39P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 26 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.5398		0.3156		0.1003	
GENDER						
Male	0.7802		0.6106		0.1762	
Female	0.5354		0.2886		0.2977	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	1.0000		0.4395		0.0687	
Black Only	0.0270		0.7848		0.6323	
American Indian or Alaska Native Only	*		*		0.8942	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.9044		0.4692		*	
More Than One Race	0.1967		0.9580		*	
Hispanic	0.9118		0.3556		0.7357	
ADULT EDUCATION						
< High School	0.4838		0.1546		0.1495	
High School Graduate	0.2708		0.7892		0.5488	
Some College	0.4901		0.6864		0.1223	
College Graduate	0.1992		0.6050		0.8521	
CURRENT EMPLOYMENT						
Full-Time	0.4830		0.4140		0.0298	
Part-Time	0.7662		0.7240		0.6162	
Unemployed	0.3124		0.0823		0.2895	
Other ¹	0.5998		0.9740		0.5449	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.40C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	594	515	187	146	130	110
GENDER						
Male	445	405	147	111	105	77
Female	355	324	106	96	72	72
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	524	459	154	127	109	90
Black Only	192	139	65	55	56	45
American Indian or Alaska Native Only	36	*	12	19	6	7
Native Hawaiian or Other Pacific Islander	*	*	1	*	1	*
Asian Only	36	34	13	6	8	4
More Than One Race	45	59	16	14	6	12
Hispanic	163	157	71	39	40	33
ADULT EDUCATION						
< High School	196	189	103	71	79	63
High School Graduate	303	282	94	86	71	52
Some College	313	295	74	61	49	36
College Graduate	325	259	70	72	44	56
CURRENT EMPLOYMENT						
Full-Time	492	448	137	115	108	77
Part-Time	167	153	60	66	33	55
Unemployed	105	90	61	29	45	19
Other ¹	200	199	58	51	33	43

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.40D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.30	0.26	0.09	0.07	0.07	0.06
GENDER						
Male	0.47	0.42	0.16	0.12	0.11	0.08
Female	0.34	0.31	0.10	0.09	0.07	0.07
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.36	0.31	0.10	0.09	0.07	0.06
Black Only	0.89	0.63	0.30	0.25	0.26	0.20
American Indian or Alaska Native Only	3.54	*	1.18	2.03	0.61	0.72
Native Hawaiian or Other Pacific Islander	*	*	0.16	*	0.13	*
Asian Only	0.57	0.50	0.20	0.08	0.13	0.05
More Than One Race	2.87	3.96	1.03	0.91	0.39	0.81
Hispanic	0.82	0.76	0.36	0.19	0.20	0.16
ADULT EDUCATION						
< High School	0.54	0.53	0.28	0.20	0.22	0.18
High School Graduate	0.45	0.42	0.14	0.13	0.11	0.08
Some College	0.65	0.61	0.15	0.13	0.10	0.07
College Graduate	0.71	0.53	0.15	0.15	0.10	0.11
CURRENT EMPLOYMENT						
Full-Time	0.44	0.39	0.12	0.10	0.10	0.07
Part-Time	0.70	0.66	0.25	0.28	0.14	0.24
Unemployed	2.03	1.93	1.18	0.62	0.87	0.40
Other ¹	0.36	0.36	0.10	0.09	0.06	0.08

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.40P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3246		0.0559		0.0455	
GENDER						
Male	0.5220		0.1474		0.0806	
Female	0.4353		0.1618		0.2164	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.8620		0.2019		0.0300	
Black Only	0.0222		0.6187		0.5037	
American Indian or Alaska Native Only	*		0.5896		0.9156	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.6821		0.0859		0.5034	
More Than One Race	0.2561		0.4402		0.5258	
Hispanic	0.6480		0.0882		0.9871	
ADULT EDUCATION						
< High School	0.4050		0.1090		0.1592	
High School Graduate	0.4577		0.5730		0.3194	
Some College	0.4093		0.3407		0.0582	
College Graduate	0.1948		0.4947		0.9246	
CURRENT EMPLOYMENT						
Full-Time	0.3184		0.1246		0.0186	
Part-Time	0.5780		0.8625		0.7262	
Unemployed	0.3255		0.1489		0.3131	
Other ¹	0.7381		0.4485		0.9890	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.41C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	557	487	101	131	59	64
AGE						
12-17	45	43	36	36	20	22
18-25	118	118	67	71	36	34
26 or Older	525	453	59	81	42	39
GENDER						
Male	390	380	87	115	51	62
Female	354	304	56	67	30	37
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	511	438	93	115	54	54
Black Only	123	136	18	21	11	10
American Indian or Alaska Native Only	36	47	6	11	3	4
Native Hawaiian or Other Pacific Islander	*	14	6	4	0	4
Asian Only	37	34	15	14	9	5
More Than One Race	39	63	13	11	6	6
Hispanic	128	127	32	31	22	17

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.41D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.25	0.22	0.05	0.06	0.03	0.03
AGE						
12-17	0.19	0.18	0.16	0.16	0.08	0.09
18-25	0.42	0.41	0.24	0.25	0.13	0.12
26 or Older	0.31	0.26	0.03	0.05	0.02	0.02
GENDER						
Male	0.37	0.35	0.08	0.11	0.05	0.06
Female	0.31	0.26	0.05	0.06	0.03	0.03
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.32	0.27	0.06	0.07	0.03	0.03
Black Only	0.49	0.54	0.07	0.08	0.04	0.04
American Indian or Alaska Native Only	3.11	4.30	0.54	0.98	0.24	0.38
Native Hawaiian or Other Pacific Islander	*	2.59	0.72	0.71	0.02	0.65
Asian Only	0.51	0.44	0.20	0.19	0.12	0.07
More Than One Race	1.99	3.37	0.65	0.58	0.30	0.30
Hispanic	0.55	0.53	0.14	0.13	0.09	0.07

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.41P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2561		0.1092		0.6325	
AGE						
12-17	0.7335		0.8101		0.6370	
18-25	0.9899		0.9661		0.8172	
26 or Older	0.2595		0.0157		0.5384	
GENDER						
Male	0.2781		0.2204		0.8697	
Female	0.6251		0.3010		0.5647	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.1135		0.0607		0.4493	
Black Only	0.8554		0.1111		0.2963	
American Indian or Alaska Native Only	0.2816		0.4562		0.7773	
Native Hawaiian or Other Pacific Islander	*		0.9270		0.2402	
Asian Only	0.2012		0.4147		0.5103	
More Than One Race	0.3230		0.9815		0.8280	
Hispanic	0.2574		0.3094		0.2417	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.42C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	45	43	36	36	20	22
GENDER						
Male	31	31	25	26	13	15
Female	30	28	24	24	14	14
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	36	39	31	33	17	21
Black Only	7	8	6	6	2	5
American Indian or Alaska Native Only	5	5	4	4	2	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	6	7	5	6	2	4
More Than One Race	6	4	4	6	1	1
Hispanic	19	14	14	11	8	6
GENDER/RACE/HISPANIC ORIGIN						
Male - White	26	29	22	23	11	15
Female - White	25	25	20	21	12	12
Male - Black	5	6	5	5	2	3
Female - Black	5	6	3	4	2	3
Male - Hispanic	12	10	9	8	6	2
Female - Hispanic	13	10	10	8	6	5

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.42D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.19	0.18	0.16	0.16	0.08	0.09
GENDER						
Male	0.26	0.26	0.21	0.21	0.11	0.13
Female	0.27	0.24	0.21	0.21	0.12	0.12
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.24	0.26	0.20	0.21	0.11	0.14
Black Only	0.22	0.25	0.18	0.19	0.08	0.14
American Indian or Alaska Native Only	3.66	3.61	3.32	3.00	1.69	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	0.66	0.81	0.55	0.68	0.27	0.54
More Than One Race	1.64	1.16	1.19	1.52	0.14	0.36
Hispanic	0.58	0.42	0.43	0.33	0.26	0.17
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.34	0.36	0.29	0.30	0.14	0.19
Female - White	0.34	0.33	0.27	0.28	0.16	0.16
Male - Black	0.33	0.34	0.30	0.32	0.12	0.19
Female - Black	0.29	0.36	0.18	0.26	0.10	0.20
Male - Hispanic	0.72	0.60	0.54	0.47	0.35	0.14
Female - Hispanic	0.82	0.61	0.61	0.49	0.41	0.32

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.42P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 to 17, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.7335		0.8101		0.6370	
GENDER						
Male	0.5470		0.6539		0.1384	
Female	0.8917		0.9081		0.4161	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.6123		0.7177		0.5362	
Black Only	0.1425		0.2199		0.1308	
American Indian or Alaska Native Only	0.4398		0.3785		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.1163		0.0267		0.1052	
More Than One Race	0.7966		0.5491		0.1784	
Hispanic	0.2672		0.1622		0.0979	
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.4359		0.5677		0.0764	
Female - White	0.9332		0.9437		0.3612	
Male - Black	0.5326		0.5602		0.3589	
Female - Black	0.1343		0.2022		0.2232	
Male - Hispanic	0.5063		0.4789		0.1336	
Female - Hispanic	0.3463		0.2325		0.5231	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.43C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	118	118	67	71	36	34
GENDER						
Male	86	78	54	53	27	28
Female	67	75	41	44	22	20
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	103	93	60	62	33	30
Black Only	24	24	15	18	7	8
American Indian or Alaska Native Only	7	8	2	6	1	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	19	12	13	9	9	3
More Than One Race	12	12	10	8	5	5
Hispanic	36	43	20	22	10	8
ADULT EDUCATION						
< High School	46	47	29	31	15	16
High School Graduate	62	62	40	38	21	19
Some College	70	68	41	43	22	21
College Graduate	38	36	21	22	11	7
CURRENT EMPLOYMENT						
Full-Time	80	79	48	50	25	23
Part-Time	58	55	33	37	18	18
Unemployed	24	25	17	19	11	12
Other ¹	44	41	26	26	12	13

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.43D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.42	0.41	0.24	0.25	0.13	0.12
GENDER						
Male	0.61	0.54	0.38	0.37	0.19	0.19
Female	0.47	0.51	0.29	0.30	0.16	0.14
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.55	0.49	0.32	0.32	0.18	0.16
Black Only	0.63	0.60	0.39	0.46	0.19	0.20
American Indian or Alaska Native Only	5.07	4.67	1.76	3.42	0.71	0.46
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.79	1.22	1.23	0.92	0.79	0.28
More Than One Race	3.54	3.87	2.91	2.56	1.57	1.75
Hispanic	0.85	0.98	0.47	0.51	0.25	0.18
ADULT EDUCATION						
< High School	0.79	0.78	0.50	0.50	0.25	0.25
High School Graduate	0.61	0.60	0.39	0.36	0.21	0.19
Some College	0.76	0.75	0.45	0.48	0.24	0.24
College Graduate	1.17	1.06	0.63	0.65	0.35	0.22
CURRENT EMPLOYMENT						
Full-Time	0.54	0.51	0.32	0.32	0.17	0.15
Part-Time	0.85	0.83	0.48	0.56	0.26	0.27
Unemployed	1.46	1.55	1.01	1.15	0.67	0.72
Other ¹	0.82	0.77	0.49	0.49	0.23	0.23

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.43P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 to 25, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.9899		0.9661		0.8172	
GENDER						
Male	0.8178		0.6255		0.8018	
Female	0.7737		0.5674		0.9596	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.5241		0.9544		0.9220	
Black Only	0.2929		0.1279		0.2514	
American Indian or Alaska Native Only	0.9171		0.4667		0.5382	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.0104		0.0986		0.1374	
More Than One Race	0.5587		0.9836		0.8922	
Hispanic	0.1559		0.6745		0.4742	
ADULT EDUCATION						
< High School	0.5628		0.4503		0.9626	
High School Graduate	0.7446		0.2342		0.9262	
Some College	0.9213		0.6649		0.8080	
College Graduate	0.8241		0.7663		0.2453	
CURRENT EMPLOYMENT						
Full-Time	0.4083		0.7307		0.3710	
Part-Time	0.2895		0.9640		0.6372	
Unemployed	0.0078		0.0499		0.4440	
Other ¹	0.1515		0.4752		0.2441	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.44C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	525	453	59	81	42	39
GENDER						
Male	367	366	55	88	41	43
Female	334	288	22	39	13	25
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	482	419	55	73	38	31
Black Only	118	133	9	6	8	3
American Indian or Alaska Native Only	35	*	1	5	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	30	28	4	7	*	1
More Than One Race	35	*	4	4	*	*
Hispanic	118	111	18	16	16	14
ADULT EDUCATION						
< High School	165	176	24	13	23	8
High School Graduate	267	248	35	34	18	10
Some College	295	244	36	37	30	25
College Graduate	294	256	19	63	5	27
CURRENT EMPLOYMENT						
Full-Time	429	398	50	76	40	35
Part-Time	158	136	17	33	5	14
Unemployed	96	87	21	16	*	12
Other ¹	180	170	17	20	13	18

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.44D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.31	0.26	0.03	0.05	0.02	0.02
GENDER						
Male	0.46	0.45	0.07	0.11	0.05	0.05
Female	0.37	0.32	0.03	0.04	0.01	0.03
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.38	0.33	0.04	0.06	0.03	0.02
Black Only	0.67	0.73	0.05	0.04	0.04	0.02
American Indian or Alaska Native Only	3.92	*	0.07	0.70	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	0.57	0.48	0.08	0.13	*	0.01
More Than One Race	2.86	*	0.32	0.31	*	*
Hispanic	0.76	0.69	0.11	0.10	0.10	0.09
ADULT EDUCATION						
< High School	0.54	0.60	0.08	0.04	0.07	0.03
High School Graduate	0.46	0.44	0.06	0.06	0.03	0.02
Some College	0.76	0.62	0.09	0.09	0.08	0.06
College Graduate	0.69	0.56	0.04	0.14	0.01	0.06
CURRENT EMPLOYMENT						
Full-Time	0.44	0.40	0.05	0.08	0.04	0.03
Part-Time	0.93	0.82	0.10	0.20	0.03	0.08
Unemployed	2.73	2.85	0.60	0.53	*	0.39
Other ¹	0.35	0.34	0.03	0.04	0.03	0.04

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.44P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 26 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2595		0.0157		0.5384	
GENDER						
Male	0.3172		0.0610		0.8590	
Female	0.5728		0.2730		0.1141	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.1606		0.0086		0.4270	
Black Only	0.9175		0.8461		0.5699	
American Indian or Alaska Native Only	*		0.4445		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.9385		0.8150		*	
More Than One Race	*		0.7996		*	
Hispanic	0.5326		0.8341		0.8142	
ADULT EDUCATION						
< High School	0.6797		0.4092		0.2239	
High School Graduate	0.1257		0.9345		0.4536	
Some College	0.7584		0.2668		0.7152	
College Graduate	0.7995		0.0014		0.0057	
CURRENT EMPLOYMENT						
Full-Time	0.3898		0.0439		0.8650	
Part-Time	0.9435		0.4519		0.3310	
Unemployed	0.9612		0.4332		*	
Other ¹	0.4065		0.6221		0.7088	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.45C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	555	482	94	122	55	57
GENDER						
Male	389	378	81	111	50	58
Female	350	301	49	59	26	33
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	510	435	86	108	51	47
Black Only	121	137	17	20	11	9
American Indian or Alaska Native Only	35	47	3	9	1	3
Native Hawaiian or Other Pacific Islander	*	14	*	4	0	*
Asian Only	37	32	15	12	9	3
More Than One Race	39	62	12	9	6	6
Hispanic	127	125	27	28	19	16
ADULT EDUCATION						
< High School	177	186	40	36	27	18
High School Graduate	277	259	55	56	27	23
Some College	309	250	61	60	38	33
College Graduate	298	265	29	69	13	29
CURRENT EMPLOYMENT						
Full-Time	447	414	72	96	47	41
Part-Time	172	147	40	50	19	18
Unemployed	99	89	28	27	11	17
Other ¹	187	176	34	35	18	22

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.45D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.28	0.24	0.05	0.06	0.03	0.03
GENDER						
Male	0.41	0.40	0.09	0.12	0.05	0.06
Female	0.34	0.29	0.05	0.06	0.03	0.03
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.35	0.29	0.06	0.07	0.03	0.03
Black Only	0.56	0.62	0.08	0.09	0.05	0.04
American Indian or Alaska Native Only	3.47	4.98	0.33	0.90	0.10	0.37
Native Hawaiian or Other Pacific Islander	*	3.01	*	0.80	0.03	*
Asian Only	0.58	0.48	0.24	0.18	0.14	0.04
More Than One Race	2.47	4.15	0.78	0.63	0.37	0.37
Hispanic	0.64	0.61	0.14	0.14	0.10	0.08
ADULT EDUCATION						
< High School	0.49	0.53	0.11	0.10	0.07	0.05
High School Graduate	0.41	0.39	0.08	0.08	0.04	0.03
Some College	0.64	0.52	0.13	0.13	0.08	0.07
College Graduate	0.65	0.54	0.06	0.14	0.03	0.06
CURRENT EMPLOYMENT						
Full-Time	0.40	0.36	0.06	0.08	0.04	0.04
Part-Time	0.73	0.63	0.17	0.22	0.08	0.08
Unemployed	1.91	1.91	0.53	0.57	0.22	0.37
Other ¹	0.33	0.31	0.06	0.06	0.03	0.04

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.45P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2707		0.0999		0.7252	
GENDER						
Male	0.3035		0.2380		0.8144	
Female	0.6150		0.2174		0.2920	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.1256		0.0580		0.5520	
Black Only	0.9166		0.2010		0.5820	
American Indian or Alaska Native Only	0.2423		0.1656		0.4473	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.1219		0.1145		0.1378	
More Than One Race	0.3242		0.7586		0.9910	
Hispanic	0.3117		0.6417		0.5360	
ADULT EDUCATION						
< High School	0.4665		0.4691		0.5809	
High School Graduate	0.1421		0.5528		0.7230	
Some College	0.7689		0.3527		0.6946	
College Graduate	0.8344		0.0053		0.0846	
CURRENT EMPLOYMENT						
Full-Time	0.4855		0.1473		0.5692	
Part-Time	0.6741		0.6388		0.9109	
Unemployed	0.5023		0.0362		0.0900	
Other ¹	0.6121		0.8202		0.2690	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.46C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	443	381	88	90	52	67
AGE						
12-17	54	49	37	32	18	17
18-25	98	94	47	42	22	19
26 or Older	416	355	61	68	40	59
GENDER						
Male	366	319	78	78	48	61
Female	217	218	40	39	20	26
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	408	351	70	79	45	56
Black Only	115	92	37	29	6	28
American Indian or Alaska Native Only	27	40	5	4	4	1
Native Hawaiian or Other Pacific Islander	*	7	3	1	0	1
Asian Only	39	32	9	10	4	7
More Than One Race	41	32	11	6	4	3
Hispanic	91	95	33	29	21	23

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.46D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.20	0.17	0.04	0.04	0.02	0.03
AGE						
12-17	0.23	0.21	0.16	0.14	0.08	0.07
18-25	0.34	0.32	0.16	0.14	0.08	0.07
26 or Older	0.25	0.21	0.04	0.04	0.02	0.03
GENDER						
Male	0.34	0.30	0.07	0.07	0.05	0.06
Female	0.19	0.19	0.03	0.03	0.02	0.02
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.25	0.22	0.04	0.05	0.03	0.03
Black Only	0.46	0.36	0.15	0.12	0.02	0.11
American Indian or Alaska Native Only	2.35	3.71	0.39	0.34	0.32	0.05
Native Hawaiian or Other Pacific Islander	*	1.35	0.43	0.26	0.04	0.24
Asian Only	0.54	0.41	0.12	0.13	0.05	0.10
More Than One Race	2.11	1.74	0.57	0.34	0.18	0.17
Hispanic	0.39	0.40	0.14	0.12	0.09	0.10

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.46P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2871		0.7619		0.5823	
AGE						
12-17	0.4586		0.0783		0.3584	
18-25	0.0047		0.2930		0.4667	
26 or Older	0.7315		0.2160		0.1969	
GENDER						
Male	0.0701		0.7759		0.7971	
Female	0.3748		0.8987		0.4146	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.4717		0.8063		0.6210	
Black Only	0.6425		0.8913		0.2776	
American Indian or Alaska Native Only	0.6530		0.2611		0.1693	
Native Hawaiian or Other Pacific Islander	*		0.8663		0.1639	
Asian Only	0.5559		0.9529		0.3635	
More Than One Race	0.7390		0.0398		0.9150	
Hispanic	0.5597		0.6459		0.4262	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.47C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	54	49	37	32	18	17
GENDER						
Male	39	38	27	24	13	12
Female	37	32	27	22	13	12
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	45	41	30	27	15	14
Black Only	15	14	9	8	5	5
American Indian or Alaska Native Only	3	4	3	*	1	1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	8	8	6	6	2	3
More Than One Race	7	8	6	2	2	2
Hispanic	20	20	14	12	10	7
GENDER/RACE/HISPANIC ORIGIN						
Male - White	34	30	23	20	10	10
Female - White	30	27	21	18	10	9
Male - Black	12	11	6	6	4	3
Female - Black	10	10	7	6	3	4
Male - Hispanic	14	16	10	10	7	6
Female - Hispanic	15	13	10	7	7	3

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.47D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.23	0.21	0.16	0.14	0.08	0.07
GENDER						
Male	0.33	0.32	0.23	0.20	0.11	0.10
Female	0.33	0.28	0.24	0.20	0.12	0.10
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.30	0.27	0.20	0.17	0.09	0.09
Black Only	0.48	0.42	0.29	0.25	0.15	0.15
American Indian or Alaska Native Only	2.56	2.98	1.92	*	1.12	0.37
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	0.93	0.99	0.65	0.73	0.22	0.32
More Than One Race	1.81	2.16	1.68	0.63	0.55	0.45
Hispanic	0.63	0.60	0.44	0.36	0.31	0.22
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.43	0.39	0.29	0.26	0.13	0.13
Female - White	0.40	0.36	0.29	0.25	0.13	0.12
Male - Black	0.72	0.63	0.39	0.35	0.22	0.18
Female - Black	0.61	0.59	0.43	0.35	0.20	0.23
Male - Hispanic	0.86	0.94	0.60	0.60	0.40	0.37
Female - Hispanic	0.98	0.81	0.61	0.44	0.46	0.21

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.47P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 to 17, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.4586		0.0783		0.3584	
GENDER						
Male	0.7382		0.4633		0.5937	
Female	0.4711		0.0769		0.4232	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.3686		0.1280		0.6225	
Black Only	0.6396		0.8910		0.6425	
American Indian or Alaska Native Only	0.2509		*		0.1948	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.5589		0.6719		0.4185	
More Than One Race	0.9680		0.0119		0.7148	
Hispanic	0.9583		0.7227		0.1752	
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.6193		0.3454		0.6762	
Female - White	0.4283		0.2329		0.7703	
Male - Black	0.5240		0.8413		0.6186	
Female - Black	0.1456		0.6987		0.3044	
Male - Hispanic	0.2799		0.3149		0.8484	
Female - Hispanic	0.1902		0.0498		0.0135	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.48C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	98	94	47	42	22	19
GENDER						
Male	70	69	38	32	18	16
Female	57	55	24	25	14	12
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	88	80	43	38	20	17
Black Only	19	16	5	6	2	4
American Indian or Alaska Native Only	3	6	0	0	*	0
Native Hawaiian or Other Pacific Islander	*	3	*	*	*	*
Asian Only	15	8	6	3	3	1
More Than One Race	12	10	8	6	3	3
Hispanic	33	30	11	14	7	6
ADULT EDUCATION						
< High School	42	37	20	18	13	6
High School Graduate	50	48	26	22	15	10
Some College	55	59	28	27	9	15
College Graduate	33	30	13	12	4	6
CURRENT EMPLOYMENT						
Full-Time	69	65	29	26	16	13
Part-Time	48	43	25	22	10	12
Unemployed	20	22	10	10	3	5
Other ¹	36	35	21	19	10	7

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.48D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.34	0.32	0.16	0.14	0.08	0.07
GENDER						
Male	0.49	0.48	0.27	0.22	0.12	0.11
Female	0.40	0.38	0.17	0.17	0.09	0.08
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.47	0.42	0.23	0.20	0.11	0.09
Black Only	0.48	0.40	0.12	0.17	0.06	0.09
American Indian or Alaska Native Only	2.56	3.57	0.24	0.03	*	0.03
Native Hawaiian or Other Pacific Islander	*	3.09	*	*	*	*
Asian Only	1.35	0.80	0.52	0.26	0.31	0.15
More Than One Race	3.34	3.23	2.38	1.85	0.83	0.91
Hispanic	0.79	0.69	0.27	0.31	0.17	0.14
ADULT EDUCATION						
< High School	0.72	0.61	0.35	0.29	0.22	0.10
High School Graduate	0.49	0.46	0.25	0.21	0.15	0.09
Some College	0.61	0.65	0.31	0.30	0.10	0.17
College Graduate	1.02	0.88	0.39	0.35	0.14	0.17
CURRENT EMPLOYMENT						
Full-Time	0.47	0.42	0.20	0.17	0.11	0.09
Part-Time	0.70	0.65	0.36	0.33	0.15	0.18
Unemployed	1.22	1.34	0.59	0.62	0.18	0.28
Other ¹	0.68	0.66	0.39	0.35	0.20	0.12

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.48P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 to 25, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0047		0.2930		0.4667	
GENDER						
Male	0.0096		0.1207		0.2226	
Female	0.1145		0.6808		0.7097	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.0429		0.2228		0.4786	
Black Only	0.6646		0.1426		0.5482	
American Indian or Alaska Native Only	0.8559		0.1099		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.1440		0.2341		0.3472	
More Than One Race	0.7484		0.3955		0.8628	
Hispanic	0.0207		0.5242		0.8176	
ADULT EDUCATION						
< High School	0.1861		0.9586		0.0934	
High School Graduate	0.0058		0.1107		0.2853	
Some College	0.5663		0.8916		0.3302	
College Graduate	0.1873		0.7974		0.3352	
CURRENT EMPLOYMENT						
Full-Time	0.0071		0.2500		0.4984	
Part-Time	0.1082		0.1791		0.5640	
Unemployed	0.0646		0.3260		0.3913	
Other ¹	0.1035		0.6001		0.8193	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.49C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	416	355	61	68	40	59
GENDER						
Male	348	299	60	64	40	54
Female	199	209	9	23	4	22
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	388	329	41	59	36	48
Black Only	112	89	36	28	2	27
American Indian or Alaska Native Only	27	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	35	27	2	7	*	7
More Than One Race	39	29	*	*	*	*
Hispanic	76	83	26	22	17	21
ADULT EDUCATION						
< High School	133	122	7	25	5	21
High School Graduate	209	191	14	32	12	29
Some College	212	190	43	18	22	14
College Graduate	263	215	40	51	31	45
CURRENT EMPLOYMENT						
Full-Time	363	304	59	57	40	50
Part-Time	108	115	6	7	4	6
Unemployed	63	58	1	17	*	17
Other ¹	126	147	12	31	5	27

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.49D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.25	0.21	0.04	0.04	0.02	0.03
GENDER						
Male	0.43	0.37	0.07	0.08	0.05	0.07
Female	0.22	0.23	0.01	0.03	0.00	0.02
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.30	0.26	0.03	0.05	0.03	0.04
Black Only	0.63	0.49	0.20	0.15	0.01	0.15
American Indian or Alaska Native Only	3.07	*	*	*	*	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	0.66	0.47	0.05	0.11	*	0.11
More Than One Race	3.13	2.41	*	*	*	*
Hispanic	0.49	0.51	0.17	0.13	0.11	0.13
ADULT EDUCATION						
< High School	0.43	0.42	0.02	0.09	0.02	0.07
High School Graduate	0.36	0.34	0.02	0.06	0.02	0.05
Some College	0.55	0.48	0.11	0.05	0.06	0.04
College Graduate	0.62	0.47	0.09	0.11	0.07	0.10
CURRENT EMPLOYMENT						
Full-Time	0.37	0.30	0.06	0.06	0.04	0.05
Part-Time	0.64	0.69	0.03	0.04	0.02	0.03
Unemployed	1.80	1.90	0.02	0.56	*	0.55
Other ¹	0.25	0.29	0.02	0.06	0.01	0.05

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.49P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 26 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.7315		0.2160		0.1969	
GENDER						
Male	0.1848		0.4034		0.3823	
Female	0.1350		0.1104		0.1481	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.8357		0.0308		0.2810	
Black Only	0.6328		0.6516		0.3583	
American Indian or Alaska Native Only	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.9463		0.3500		*	
More Than One Race	0.8557		*		*	
Hispanic	0.8257		0.4736		0.8521	
ADULT EDUCATION						
< High School	0.7093		0.1857		0.3525	
High School Graduate	0.9116		0.0667		0.2709	
Some College	0.3340		0.2388		0.4956	
College Graduate	0.9284		0.3464		0.3514	
CURRENT EMPLOYMENT						
Full-Time	0.1164		0.8243		0.7545	
Part-Time	0.1953		0.4911		0.5969	
Unemployed	0.7942		0.0988		*	
Other ¹	0.1379		0.0984		0.1348	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.50C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	435	378	77	83	47	65
GENDER						
Male	362	316	72	74	46	60
Female	209	217	26	33	14	25
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	401	347	61	74	42	54
Black Only	114	90	36	29	3	28
American Indian or Alaska Native Only	27	*	3	*	*	*
Native Hawaiian or Other Pacific Islander	*	7	*	*	*	*
Asian Only	38	30	6	7	3	7
More Than One Race	41	31	9	6	3	3
Hispanic	88	93	28	26	19	22
ADULT EDUCATION						
< High School	144	132	22	30	14	22
High School Graduate	220	200	29	40	18	31
Some College	220	198	52	35	24	21
College Graduate	265	218	41	52	31	45
CURRENT EMPLOYMENT						
Full-Time	374	313	68	65	45	52
Part-Time	123	124	27	24	11	13
Unemployed	66	63	10	20	3	17
Other ¹	133	154	22	36	11	28

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.50D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.22	0.19	0.04	0.04	0.02	0.03
GENDER						
Male	0.38	0.33	0.08	0.08	0.05	0.06
Female	0.20	0.21	0.02	0.03	0.01	0.02
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.27	0.24	0.04	0.05	0.03	0.04
Black Only	0.52	0.41	0.17	0.13	0.01	0.12
American Indian or Alaska Native Only	2.67	*	0.34	*	*	*
Native Hawaiian or Other Pacific Islander	*	1.48	*	*	*	*
Asian Only	0.59	0.44	0.10	0.10	0.05	0.10
More Than One Race	2.60	2.07	0.56	0.39	0.18	0.19
Hispanic	0.45	0.45	0.14	0.13	0.09	0.11
ADULT EDUCATION						
< High School	0.39	0.37	0.06	0.09	0.04	0.06
High School Graduate	0.32	0.30	0.04	0.06	0.03	0.05
Some College	0.46	0.41	0.11	0.07	0.05	0.04
College Graduate	0.58	0.44	0.09	0.11	0.07	0.09
CURRENT EMPLOYMENT						
Full-Time	0.33	0.27	0.06	0.06	0.04	0.05
Part-Time	0.52	0.53	0.11	0.10	0.05	0.06
Unemployed	1.27	1.35	0.20	0.43	0.06	0.37
Other ¹	0.24	0.28	0.04	0.07	0.02	0.05

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.50P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3249		0.6377		0.3773	
GENDER						
Male	0.0740		0.9838		0.6842	
Female	0.2979		0.2011		0.1657	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.5360		0.3157		0.4943	
Black Only	0.5856		0.8487		0.3177	
American Indian or Alaska Native Only	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.4957		0.8356		0.4804	
More Than One Race	0.7426		0.3527		0.9332	
Hispanic	0.5664		0.7360		0.7997	
ADULT EDUCATION						
< High School	0.6005		0.2572		0.8991	
High School Graduate	0.6143		0.6837		0.6625	
Some College	0.2630		0.2980		0.9659	
College Graduate	0.9151		0.4153		0.2916	
CURRENT EMPLOYMENT						
Full-Time	0.0439		0.8113		0.9291	
Part-Time	0.5348		0.2653		0.7416	
Unemployed	0.3515		0.0417		0.0921	
Other ¹	0.2895		0.0894		0.2001	

*Low precision; no estimate reported.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.51C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	645	582	296	259	209	178
AGE						
12-17	58	61	47	47	30	31
18-25	117	102	86	73	52	50
26 or Older	621	566	272	243	195	170
GENDER						
Male	461	422	197	168	140	128
Female	447	381	211	201	149	124
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	566	527	254	230	180	160
Black Only	183	146	90	62	64	42
American Indian or Alaska Native Only	55	47	*	16	18	15
Native Hawaiian or Other Pacific Islander	*	19	6	10	1	10
Asian Only	93	113	31	44	27	31
More Than One Race	46	52	17	17	12	13
Hispanic	185	162	97	95	66	55

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.51D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.29	0.26	0.13	0.12	0.09	0.08
AGE						
12-17	0.25	0.26	0.20	0.20	0.13	0.13
18-25	0.41	0.35	0.30	0.25	0.18	0.17
26 or Older	0.37	0.33	0.16	0.14	0.11	0.10
GENDER						
Male	0.43	0.39	0.19	0.16	0.13	0.12
Female	0.39	0.33	0.18	0.17	0.13	0.11
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.35	0.32	0.16	0.14	0.11	0.10
Black Only	0.74	0.57	0.36	0.24	0.26	0.16
American Indian or Alaska Native Only	4.78	4.33	*	1.48	1.58	1.37
Native Hawaiian or Other Pacific Islander	*	3.42	0.71	1.90	0.17	1.85
Asian Only	1.28	1.49	0.43	0.58	0.37	0.41
More Than One Race	2.38	2.81	0.86	0.93	0.63	0.67
Hispanic	0.80	0.68	0.42	0.40	0.29	0.23

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.51P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0230		0.1617		0.6054	
AGE						
12-17	0.9950		0.8173		0.6305	
18-25	0.0077		0.0930		0.6614	
26 or Older	0.0614		0.3253		0.6019	
GENDER						
Male	0.1396		0.1696		0.5007	
Female	0.0801		0.5210		0.9470	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.0539		0.2646		0.6318	
Black Only	0.2586		0.1896		0.5264	
American Indian or Alaska Native Only	0.5375		*		0.8442	
Native Hawaiian or Other Pacific Islander	*		0.2019		0.0881	
Asian Only	0.9951		0.5633		0.7964	
More Than One Race	0.5056		0.8238		0.6291	
Hispanic	0.5101		0.8258		0.9905	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.52C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	58	61	47	47	30	31
GENDER						
Male	39	43	32	32	21	21
Female	42	40	35	33	22	23
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	48	49	38	39	24	25
Black Only	21	21	17	15	11	11
American Indian or Alaska Native Only	3	5	3	3	3	2
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	9	10	6	9	2	6
More Than One Race	7	6	6	3	3	2
Hispanic	22	23	18	18	11	13
GENDER/RACE/HISPANIC ORIGIN						
Male - White	32	34	26	27	17	17
Female - White	34	34	29	28	18	19
Male - Black	17	16	12	11	9	7
Female - Black	16	12	13	10	7	7
Male - Hispanic	14	17	11	14	7	11
Female - Hispanic	16	14	13	11	8	8

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.52D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.25	0.26	0.20	0.20	0.13	0.13
GENDER						
Male	0.33	0.36	0.27	0.26	0.18	0.18
Female	0.37	0.35	0.31	0.29	0.19	0.20
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.31	0.32	0.25	0.26	0.16	0.16
Black Only	0.66	0.62	0.52	0.44	0.34	0.32
American Indian or Alaska Native Only	2.33	3.80	2.51	1.96	2.23	1.65
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.05	1.21	0.69	1.04	0.26	0.73
More Than One Race	2.01	1.57	1.69	0.92	0.92	0.64
Hispanic	0.69	0.70	0.56	0.55	0.35	0.40
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.40	0.44	0.33	0.35	0.22	0.22
Female - White	0.46	0.46	0.38	0.38	0.24	0.26
Male - Black	1.04	0.92	0.74	0.62	0.54	0.41
Female - Black	1.00	0.74	0.80	0.62	0.45	0.46
Male - Hispanic	0.87	1.00	0.68	0.83	0.43	0.62
Female - Hispanic	1.00	0.87	0.83	0.67	0.54	0.50

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.52P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.9950		0.8173		0.6305	
GENDER						
Male	0.5485		0.3620		0.8130	
Female	0.5652		0.2441		0.6670	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.3580		0.4444		0.5342	
Black Only	0.2762		0.1417		0.5128	
American Indian or Alaska Native Only	0.6769		0.1636		0.1899	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.1762		0.0617		0.0298	
More Than One Race	0.1973		0.0757		0.7231	
Hispanic	0.3167		0.3848		0.9863	
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.5389		0.2408		0.5797	
Female - White	0.4833		0.9764		0.7688	
Male - Black	0.6405		0.3997		0.0692	
Female - Black	0.2398		0.1798		0.3297	
Male - Hispanic	0.9386		0.7839		0.4739	
Female - Hispanic	0.1342		0.0819		0.4695	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.53C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	117	102	86	73	52	50
GENDER						
Male	81	74	61	54	40	35
Female	73	69	55	52	29	32
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	98	87	73	67	44	45
Black Only	29	26	21	19	12	12
American Indian or Alaska Native Only	4	*	3	*	1	*
Native Hawaiian or Other Pacific Islander	*	*	*	3	*	*
Asian Only	19	18	14	13	7	5
More Than One Race	11	12	10	10	6	8
Hispanic	36	34	25	27	18	16
ADULT EDUCATION						
< High School	51	46	38	34	27	23
High School Graduate	63	62	47	47	31	28
Some College	66	60	45	44	24	30
College Graduate	37	30	27	19	10	13
CURRENT EMPLOYMENT						
Full-Time	78	72	56	50	35	31
Part-Time	58	49	39	36	25	25
Unemployed	27	24	20	20	14	14
Other ¹	47	37	38	29	21	20

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.53D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.41	0.35	0.30	0.25	0.18	0.17
GENDER						
Male	0.57	0.52	0.43	0.37	0.28	0.24
Female	0.51	0.47	0.38	0.35	0.20	0.22
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.52	0.45	0.39	0.35	0.23	0.23
Black Only	0.76	0.67	0.56	0.49	0.31	0.30
American Indian or Alaska Native Only	3.20	*	2.24	*	0.79	*
Native Hawaiian or Other Pacific Islander	*	*	*	2.56	*	*
Asian Only	1.72	1.77	1.29	1.25	0.61	0.47
More Than One Race	3.17	4.01	2.95	3.26	1.87	2.55
Hispanic	0.87	0.78	0.60	0.62	0.42	0.37
ADULT EDUCATION						
< High School	0.87	0.76	0.65	0.56	0.46	0.38
High School Graduate	0.61	0.60	0.46	0.45	0.30	0.27
Some College	0.73	0.66	0.49	0.48	0.27	0.33
College Graduate	1.12	0.87	0.82	0.56	0.29	0.39
CURRENT EMPLOYMENT						
Full-Time	0.53	0.47	0.38	0.32	0.24	0.20
Part-Time	0.85	0.74	0.57	0.54	0.36	0.38
Unemployed	1.60	1.48	1.20	1.24	0.87	0.89
Other ¹	0.90	0.69	0.73	0.55	0.39	0.38

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.53P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0077		0.0930		0.6614	
GENDER						
Male	0.1733		0.5353		0.3909	
Female	0.0055		0.0558		0.7020	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.1169		0.2385		0.8893	
Black Only	0.2055		0.7985		0.5075	
American Indian or Alaska Native Only	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.4980		0.5203		0.3104	
More Than One Race	0.8259		0.9577		0.9369	
Hispanic	0.0022		0.1571		0.1468	
ADULT EDUCATION						
< High School	0.1023		0.2115		0.2296	
High School Graduate	0.2756		0.6654		0.2805	
Some College	0.0318		0.3469		0.4918	
College Graduate	0.6617		0.2869		0.0765	
CURRENT EMPLOYMENT						
Full-Time	0.0132		0.0802		0.5149	
Part-Time	0.5567		0.7765		0.5659	
Unemployed	0.0266		0.1129		0.8754	
Other ¹	0.0038		0.0998		0.5177	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.54C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	621	566	272	243	195	170
GENDER						
Male	446	412	183	156	132	120
Female	426	369	192	185	142	119
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	548	513	235	216	169	153
Black Only	174	146	83	60	62	39
American Indian or Alaska Native Only	*	*	*	14	*	13
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	89	109	26	40	26	30
More Than One Race	43	50	9	13	9	9
Hispanic	174	153	93	93	63	52
ADULT EDUCATION						
< High School	223	190	117	112	97	68
High School Graduate	351	308	138	143	93	108
Some College	318	277	142	106	105	71
College Graduate	345	293	155	111	130	74
CURRENT EMPLOYMENT						
Full-Time	483	455	197	189	132	129
Part-Time	210	179	87	77	63	50
Unemployed	95	99	71	32	53	18
Other ¹	287	245	150	120	116	82

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.54D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.37	0.33	0.16	0.14	0.11	0.10
GENDER						
Male	0.56	0.51	0.23	0.19	0.16	0.15
Female	0.48	0.41	0.22	0.21	0.16	0.13
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.43	0.40	0.18	0.17	0.13	0.12
Black Only	0.98	0.80	0.47	0.33	0.35	0.21
American Indian or Alaska Native Only	*	*	*	1.75	*	1.69
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.69	1.88	0.50	0.70	0.49	0.52
More Than One Race	3.51	4.18	0.73	1.07	0.72	0.77
Hispanic	1.12	0.94	0.60	0.58	0.41	0.32
ADULT EDUCATION						
< High School	0.73	0.65	0.38	0.38	0.31	0.23
High School Graduate	0.61	0.54	0.24	0.25	0.16	0.19
Some College	0.82	0.71	0.36	0.27	0.27	0.18
College Graduate	0.81	0.64	0.36	0.24	0.31	0.16
CURRENT EMPLOYMENT						
Full-Time	0.49	0.45	0.20	0.19	0.13	0.13
Part-Time	1.24	1.07	0.52	0.46	0.37	0.30
Unemployed	2.71	3.23	2.03	1.04	1.51	0.60
Other ¹	0.57	0.49	0.30	0.24	0.23	0.16

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.54P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0614		0.3253		0.6019	
GENDER						
Male	0.1755		0.1435		0.6080	
Female	0.1966		0.9925		0.8124	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.0680		0.3098		0.5560	
Black Only	0.4210		0.3108		0.4888	
American Indian or Alaska Native Only	*		*		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.9694		0.4983		0.8588	
More Than One Race	0.3700		0.4363		0.6369	
Hispanic	0.8973		0.7246		0.6988	
ADULT EDUCATION						
< High School	0.0694		0.7832		0.5166	
High School Graduate	0.6774		0.1657		0.0721	
Some College	0.1080		0.0270		0.3691	
College Graduate	0.1624		0.2842		0.1969	
CURRENT EMPLOYMENT						
Full-Time	0.1156		0.8839		0.7356	
Part-Time	0.1926		0.4023		0.7233	
Unemployed	0.4463		0.0474		0.0886	
Other ¹	0.3980		0.7279		0.6123	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.55C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	639	581	290	258	206	178
GENDER						
Male	458	421	193	166	137	126
Female	441	378	206	199	147	123
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	566	528	252	230	178	161
Black Only	179	145	87	61	63	41
American Indian or Alaska Native Only	54	46	*	15	18	14
Native Hawaiian or Other Pacific Islander	*	18	4	10	1	*
Asian Only	92	112	30	42	27	30
More Than One Race	46	52	15	17	12	12
Hispanic	182	158	96	94	65	54
ADULT EDUCATION						
< High School	236	196	126	119	104	74
High School Graduate	359	318	147	154	99	112
Some College	329	285	152	115	108	76
College Graduate	347	295	160	113	131	73
CURRENT EMPLOYMENT						
Full-Time	492	465	208	196	137	133
Part-Time	217	188	94	86	66	56
Unemployed	100	100	75	40	55	25
Other ¹	292	249	163	123	124	84

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.55D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.32	0.29	0.15	0.13	0.10	0.09
GENDER						
Male	0.49	0.44	0.20	0.17	0.15	0.13
Female	0.43	0.36	0.20	0.19	0.14	0.12
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.39	0.36	0.17	0.16	0.12	0.11
Black Only	0.83	0.66	0.40	0.28	0.29	0.19
American Indian or Alaska Native Only	5.31	4.84	*	1.63	1.77	1.48
Native Hawaiian or Other Pacific Islander	*	3.81	0.57	2.08	0.10	*
Asian Only	1.45	1.65	0.47	0.63	0.42	0.45
More Than One Race	2.93	3.45	0.97	1.14	0.74	0.82
Hispanic	0.92	0.77	0.48	0.46	0.33	0.26
ADULT EDUCATION						
< High School	0.65	0.56	0.35	0.34	0.28	0.21
High School Graduate	0.53	0.47	0.22	0.23	0.15	0.17
Some College	0.69	0.59	0.32	0.24	0.23	0.16
College Graduate	0.76	0.60	0.35	0.23	0.29	0.15
CURRENT EMPLOYMENT						
Full-Time	0.44	0.40	0.18	0.17	0.12	0.11
Part-Time	0.92	0.80	0.40	0.37	0.28	0.24
Unemployed	1.93	2.13	1.44	0.86	1.06	0.53
Other ¹	0.52	0.45	0.29	0.22	0.22	0.15

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.55P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0222		0.1713		0.5515	
GENDER						
Male	0.1234		0.1224		0.4713	
Female	0.0872		0.6556		0.8883	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.0453		0.2187		0.5713	
Black Only	0.3133		0.2947		0.6039	
American Indian or Alaska Native Only	0.5819		*		0.6811	
Native Hawaiian or Other Pacific Islander	*		0.1987		*	
Asian Only	0.8940		0.7903		0.9515	
More Than One Race	0.3890		0.7471		0.6866	
Hispanic	0.5892		0.9638		0.9900	
ADULT EDUCATION						
< High School	0.0548		0.6784		0.4260	
High School Graduate	0.8037		0.2024		0.1409	
Some College	0.0469		0.0157		0.4888	
College Graduate	0.1520		0.2160		0.2675	
CURRENT EMPLOYMENT						
Full-Time	0.0555		0.5861		0.8430	
Part-Time	0.1534		0.3729		0.5757	
Unemployed	0.1983		0.2457		0.1541	
Other ¹	0.2004		0.4837		0.7227	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.56C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	787	704	341	306	242	209
AGE						
12-17	73	74	59	60	38	39
18-25	139	133	104	98	64	62
26 or Older	746	662	308	274	225	195
GENDER						
Male	525	544	242	224	174	163
Female	539	441	229	219	165	143
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	668	608	288	281	209	188
Black Only	227	183	113	83	85	63
American Indian or Alaska Native Only	55	50	41	29	19	15
Native Hawaiian or Other Pacific Islander	*	*	8	13	2	11
Asian Only	106	119	36	46	29	35
More Than One Race	58	70	23	23	14	18
Hispanic	234	216	120	107	84	67

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.56D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.36	0.32	0.15	0.14	0.11	0.09
AGE						
12-17	0.31	0.32	0.26	0.26	0.17	0.17
18-25	0.49	0.46	0.37	0.34	0.22	0.21
26 or Older	0.44	0.39	0.18	0.16	0.13	0.11
GENDER						
Male	0.49	0.51	0.23	0.21	0.16	0.15
Female	0.47	0.38	0.20	0.19	0.14	0.12
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.41	0.37	0.18	0.17	0.13	0.12
Black Only	0.91	0.72	0.45	0.33	0.34	0.25
American Indian or Alaska Native Only	4.80	4.58	3.55	2.69	1.62	1.41
Native Hawaiian or Other Pacific Islander	*	*	1.03	2.31	0.24	1.97
Asian Only	1.46	1.57	0.49	0.60	0.41	0.45
More Than One Race	2.96	3.78	1.18	1.24	0.71	0.94
Hispanic	1.01	0.91	0.52	0.45	0.37	0.28

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.56P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2462		0.2589		0.3178	
AGE						
12-17	0.6289		0.3114		0.6965	
18-25	0.0383		0.0825		0.5779	
26 or Older	0.4316		0.6127		0.3108	
GENDER						
Male	0.9889		0.2486		0.3213	
Female	0.0934		0.6585		0.7048	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.7472		0.5806		0.3741	
Black Only	0.0272		0.1762		0.4990	
American Indian or Alaska Native Only	0.8226		0.7408		0.9816	
Native Hawaiian or Other Pacific Islander	*		0.1689		0.0580	
Asian Only	0.6085		0.8112		0.5750	
More Than One Race	0.2600		0.9444		0.9121	
Hispanic	0.4312		0.4637		0.5966	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.57C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	73	74	59	60	38	39
GENDER						
Male	49	53	42	41	27	26
Female	51	48	42	41	27	28
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	60	60	50	49	32	32
Black Only	24	24	18	17	12	12
American Indian or Alaska Native Only	5	7	5	6	3	4
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	12	12	9	10	4	8
More Than One Race	9	9	8	5	4	3
Hispanic	29	26	24	22	17	16
GENDER/RACE/HISPANIC ORIGIN						
Male - White	42	43	35	34	22	22
Female - White	41	39	34	34	22	23
Male - Black	18	19	13	13	9	8
Female - Black	17	15	14	12	8	9
Male - Hispanic	18	21	16	18	12	13
Female - Hispanic	20	17	17	14	12	10

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.57D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.31	0.32	0.26	0.26	0.17	0.17
GENDER						
Male	0.42	0.45	0.35	0.34	0.23	0.22
Female	0.45	0.42	0.37	0.36	0.24	0.25
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.39	0.39	0.33	0.32	0.21	0.21
Black Only	0.73	0.72	0.56	0.51	0.36	0.37
American Indian or Alaska Native Only	4.08	5.05	3.58	4.36	2.40	2.84
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.30	1.42	1.05	1.23	0.44	0.93
More Than One Race	2.42	2.37	2.05	1.46	1.08	0.79
Hispanic	0.90	0.80	0.75	0.68	0.54	0.48
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.53	0.55	0.44	0.44	0.28	0.27
Female - White	0.55	0.52	0.46	0.46	0.30	0.31
Male - Black	1.10	1.09	0.82	0.76	0.55	0.46
Female - Black	1.06	0.90	0.86	0.73	0.49	0.52
Male - Hispanic	1.11	1.20	0.98	1.04	0.73	0.76
Female - Hispanic	1.29	1.09	1.07	0.86	0.75	0.64

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.57P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.6289		0.3114		0.6965	
GENDER						
Male	0.9837		0.6076		0.5893	
Female	0.4660		0.3424		0.9787	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.7641		0.4086		0.8036	
Black Only	0.9653		0.5293		0.8505	
American Indian or Alaska Native Only	0.6362		0.5262		0.6138	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.1296		0.0801		0.0066	
More Than One Race	0.9299		0.3861		0.6671	
Hispanic	0.3306		0.6205		0.5964	
GENDER/RACE/HISPANIC ORIGIN						
Male - White	0.8316		0.4736		0.5127	
Female - White	0.8242		0.6188		0.7711	
Male - Black	0.7443		0.6500		0.1538	
Female - Black	0.7584		0.6387		0.0802	
Male - Hispanic	0.9022		0.7055		0.6786	
Female - Hispanic	0.0837		0.1966		0.1874	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.58C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	139	133	104	98	64	62
GENDER						
Male	94	93	73	71	48	45
Female	84	87	64	64	37	41
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	115	104	88	83	55	53
Black Only	37	37	28	26	18	15
American Indian or Alaska Native Only	7	*	4	8	1	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	22	21	18	15	10	7
More Than One Race	13	12	12	11	8	8
Hispanic	51	50	35	35	24	26
ADULT EDUCATION						
< High School	57	58	45	44	33	31
High School Graduate	72	69	56	55	38	35
Some College	78	76	58	59	33	37
College Graduate	41	41	33	29	15	16
CURRENT EMPLOYMENT						
Full-Time	91	94	68	66	43	42
Part-Time	67	62	48	47	31	30
Unemployed	32	28	26	24	19	18
Other ¹	54	48	44	37	26	24

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.58D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.49	0.46	0.37	0.34	0.22	0.21
GENDER						
Male	0.67	0.65	0.51	0.49	0.34	0.31
Female	0.59	0.60	0.45	0.44	0.26	0.28
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.61	0.54	0.47	0.43	0.29	0.28
Black Only	0.97	0.93	0.73	0.67	0.48	0.38
American Indian or Alaska Native Only	5.21	*	2.80	4.46	1.04	*
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	2.04	2.07	1.63	1.52	0.93	0.69
More Than One Race	3.73	4.10	3.56	3.59	2.31	2.67
Hispanic	1.22	1.16	0.82	0.81	0.56	0.59
ADULT EDUCATION						
< High School	0.98	0.95	0.77	0.71	0.56	0.50
High School Graduate	0.71	0.66	0.54	0.53	0.37	0.34
Some College	0.86	0.84	0.64	0.65	0.36	0.41
College Graduate	1.25	1.19	1.00	0.83	0.47	0.46
CURRENT EMPLOYMENT						
Full-Time	0.62	0.62	0.47	0.43	0.30	0.28
Part-Time	0.98	0.92	0.70	0.70	0.45	0.45
Unemployed	1.91	1.73	1.56	1.47	1.12	1.13
Other ¹	1.03	0.89	0.83	0.68	0.50	0.46

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.58P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0383		0.0825		0.5779	
GENDER						
Male	0.2168		0.1627		0.2191	
Female	0.0460		0.2501		0.4975	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.2640		0.1820		0.6427	
Black Only	0.4865		0.5441		0.6447	
American Indian or Alaska Native Only	*		0.8501		*	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.0771		0.1964		0.3843	
More Than One Race	0.9027		0.8134		0.7555	
Hispanic	0.0429		0.0825		0.5374	
ADULT EDUCATION						
< High School	0.2868		0.1938		0.1737	
High School Graduate	0.2193		0.1971		0.0947	
Some College	0.1815		0.6774		0.1276	
College Graduate	0.4379		0.6295		0.2980	
CURRENT EMPLOYMENT						
Full-Time	0.0150		0.0929		0.2720	
Part-Time	0.9132		0.4854		0.8687	
Unemployed	0.0221		0.2179		0.7389	
Other ¹	0.0245		0.1946		0.5916	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.59C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	746	662	308	274	225	195
GENDER						
Male	504	519	218	202	163	149
Female	512	424	204	197	157	134
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	640	582	259	249	194	177
Black Only	216	175	103	77	80	59
American Indian or Alaska Native Only	*	*	*	21	18	13
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	99	114	27	41	26	34
More Than One Race	54	*	15	17	9	15
Hispanic	213	196	110	99	80	59
ADULT EDUCATION						
< High School	256	234	134	118	110	89
High School Graduate	407	360	151	160	117	119
Some College	365	331	155	121	116	82
College Graduate	424	348	170	141	138	95
CURRENT EMPLOYMENT						
Full-Time	586	544	234	224	163	152
Part-Time	228	195	98	88	68	71
Unemployed	111	102	78	43	68	30
Other ¹	321	270	153	129	120	91

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.59D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.44	0.39	0.18	0.16	0.13	0.11
GENDER						
Male	0.63	0.64	0.27	0.25	0.20	0.18
Female	0.57	0.47	0.23	0.22	0.18	0.15
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.50	0.45	0.20	0.19	0.15	0.14
Black Only	1.22	0.97	0.58	0.42	0.45	0.32
American Indian or Alaska Native Only	*	*	*	2.72	2.05	1.69
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian Only	1.87	1.98	0.51	0.71	0.49	0.58
More Than One Race	4.38	*	1.25	1.45	0.72	1.26
Hispanic	1.37	1.21	0.71	0.61	0.52	0.37
ADULT EDUCATION						
< High School	0.84	0.80	0.44	0.40	0.36	0.31
High School Graduate	0.71	0.64	0.26	0.28	0.20	0.21
Some College	0.94	0.84	0.40	0.31	0.30	0.21
College Graduate	1.00	0.76	0.40	0.31	0.32	0.21
CURRENT EMPLOYMENT						
Full-Time	0.60	0.54	0.24	0.22	0.17	0.15
Part-Time	1.35	1.17	0.58	0.53	0.40	0.43
Unemployed	3.16	3.35	2.23	1.41	1.94	0.99
Other ¹	0.63	0.54	0.30	0.25	0.24	0.18

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.59P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.4316		0.6127		0.3108	
GENDER						
Male	0.8361		0.4621		0.4138	
Female	0.1842		0.9714		0.5580	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.8803		0.9267		0.3679	
Black Only	0.0300		0.1187		0.3757	
American Indian or Alaska Native Only	*		*		0.8886	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian Only	0.8221		0.4401		0.6710	
More Than One Race	*		0.7665		0.5360	
Hispanic	0.8593		0.9272		0.8113	
ADULT EDUCATION						
< High School	0.2722		0.3135		0.2639	
High School Graduate	0.4356		0.0721		0.2943	
Some College	0.3469		0.0487		0.1551	
College Graduate	0.3172		0.8575		0.5104	
CURRENT EMPLOYMENT						
Full-Time	0.7062		1.0000		0.4861	
Part-Time	0.4510		0.5933		0.9392	
Unemployed	0.6700		0.1492		0.1303	
Other ¹	0.5099		0.9802		0.8698	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.60C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	776	693	330	300	239	208
GENDER						
Male	522	535	235	218	172	160
Female	532	437	220	214	162	141
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	662	603	283	277	207	188
Black Only	224	180	108	80	84	61
American Indian or Alaska Native Only	55	49	*	24	18	14
Native Hawaiian or Other Pacific Islander	*	*	7	12	1	10
Asian Only	104	118	33	44	28	34
More Than One Race	58	69	22	22	13	17
Hispanic	232	210	117	105	83	65
ADULT EDUCATION						
< High School	272	245	146	129	120	98
High School Graduate	418	370	165	174	124	125
Some College	381	342	172	141	123	92
College Graduate	427	355	175	144	139	96
CURRENT EMPLOYMENT						
Full-Time	601	563	245	237	170	158
Part-Time	243	206	109	104	74	76
Unemployed	117	104	85	54	71	37
Other ¹	332	276	168	136	129	94

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.60D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 1999 and 2000

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.39	0.35	0.17	0.15	0.12	0.10
GENDER						
Male	0.55	0.56	0.25	0.23	0.18	0.17
Female	0.51	0.42	0.21	0.21	0.16	0.13
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.45	0.41	0.19	0.19	0.14	0.13
Black Only	1.03	0.81	0.50	0.36	0.39	0.28
American Indian or Alaska Native Only	5.39	5.16	*	2.58	1.81	1.47
Native Hawaiian or Other Pacific Islander	*	*	0.89	2.52	0.18	2.19
Asian Only	1.63	1.74	0.52	0.65	0.43	0.50
More Than One Race	3.66	4.62	1.39	1.49	0.83	1.16
Hispanic	1.17	1.02	0.59	0.51	0.42	0.32
ADULT EDUCATION						
< High School	0.75	0.69	0.40	0.36	0.33	0.28
High School Graduate	0.62	0.55	0.24	0.26	0.18	0.19
Some College	0.79	0.71	0.36	0.29	0.26	0.19
College Graduate	0.93	0.72	0.38	0.29	0.31	0.20
CURRENT EMPLOYMENT						
Full-Time	0.53	0.48	0.22	0.20	0.15	0.14
Part-Time	1.02	0.88	0.46	0.45	0.31	0.32
Unemployed	2.26	2.22	1.64	1.16	1.38	0.80
Other ¹	0.59	0.50	0.30	0.24	0.23	0.17

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.60P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Demographic Characteristics: 2000 Versus 1999

Demographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2597		0.3386		0.2835	
GENDER						
Male	0.9824		0.2752		0.2780	
Female	0.1059		0.7991		0.6999	
HISPANIC ORIGIN AND RACE						
Not Hispanic						
White Only	0.7633		0.6969		0.3563	
Black Only	0.0252		0.1920		0.4653	
American Indian or Alaska Native Only	0.7711		*		0.9153	
Native Hawaiian or Other Pacific Islander	*		0.1307		0.0748	
Asian Only	0.4911		0.9340		0.9156	
More Than One Race	0.2471		0.8721		0.8361	
Hispanic	0.5031		0.5177		0.6774	
ADULT EDUCATION						
< High School	0.3011		0.2928		0.2156	
High School Graduate	0.5520		0.1691		0.5639	
Some College	0.2308		0.0429		0.3718	
College Graduate	0.2825		0.7660		0.5936	
CURRENT EMPLOYMENT						
Full-Time	0.4541		0.6918		0.3702	
Part-Time	0.4405		0.4190		0.8858	
Unemployed	0.8385		0.4716		0.2327	
Other ¹	0.3079		0.7347		0.9850	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

¹ Retired, disabled, homemaker, student, or "other."

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.61C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	548	308	222
Black	179	116	83
American Indian or Alaska Native	42	29	19
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	44	10	3
Filipino	50	19	8
Japanese	26	13	12
Asian Indian	43	19	13
Korean	33	16	14
Vietnamese	23	12	9
HISPANIC			
Mexican	146	71	50
Puerto Rican	69	47	37
Central or South American	58	23	15
Cuban	47	20	10

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.61D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.34	0.19	0.14
Black	0.71	0.46	0.33
American Indian or Alaska Native	3.73	2.59	1.74
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	3.08	0.68	0.24
Filipino	3.32	1.27	0.55
Japanese	4.08	2.05	1.98
Asian Indian	2.75	1.21	0.86
Korean	5.29	2.48	2.28
Vietnamese	3.49	1.80	1.35
HISPANIC			
Mexican	0.96	0.47	0.33
Puerto Rican	2.68	1.85	1.45
Central or South American	1.80	0.72	0.48
Cuban	4.14	1.79	0.85

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.62C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	49	42	31
Black	23	18	13
American Indian or Alaska Native	7	7	5
Native Hawaiian	*	*	*
Other Pacific Islander	2	1	1
Chinese	4	3	3
Filipino	6	5	3
Japanese	1	1	0
Asian Indian	4	3	2
Korean	3	3	3
Vietnamese	6	*	4
HISPANIC			
Mexican	17	16	12
Puerto Rican	8	7	5
Central or South American	7	6	5
Cuban	4	4	3

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.62D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.32	0.28	0.20
Black	0.70	0.56	0.40
American Indian or Alaska Native	5.08	5.15	3.58
Native Hawaiian	*	*	*
Other Pacific Islander	4.57	2.92	1.96
Chinese	2.78	2.24	1.94
Filipino	3.53	3.16	2.01
Japanese	3.51	3.17	0.98
Asian Indian	2.86	2.31	1.55
Korean	3.68	3.42	3.04
Vietnamese	4.54	*	2.80
HISPANIC			
Mexican	0.75	0.70	0.52
Puerto Rican	2.11	1.80	1.25
Central or South American	1.88	1.55	1.30
Cuban	4.13	3.83	3.46

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.63C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	84	79	66
Black	35	30	26
American Indian or Alaska Native	6	6	4
Native Hawaiian	*	*	*
Other Pacific Islander	*	3	*
Chinese	7	6	2
Filipino	8	7	5
Japanese	*	*	3
Asian Indian	8	6	5
Korean	5	5	*
Vietnamese	5	3	3
HISPANIC			
Mexican	34	27	21
Puerto Rican	12	11	10
Central or South American	12	10	8
Cuban	5	5	*

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.63D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.44	0.42	0.35
Black	0.90	0.78	0.68
American Indian or Alaska Native	3.67	3.71	2.64
Native Hawaiian	*	*	*
Other Pacific Islander	*	4.81	*
Chinese	4.10	3.69	1.09
Filipino	4.47	3.68	2.69
Japanese	*	*	3.81
Asian Indian	3.83	3.02	2.26
Korean	5.39	5.23	*
Vietnamese	3.87	2.47	2.33
HISPANIC			
Mexican	1.12	0.91	0.69
Puerto Rican	2.69	2.47	2.22
Central or South American	2.27	1.89	1.55
Cuban	5.11	5.38	*

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.64C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	523	272	198
Black	171	103	75
American Indian or Alaska Native	41	26	17
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	43	4	1
Filipino	48	15	5
Japanese	24	12	11
Asian Indian	41	17	*
Korean	*	*	*
Vietnamese	*	8	7
HISPANIC			
Mexican	132	61	45
Puerto Rican	66	46	35
Central or South American	56	20	12
Cuban	43	18	6

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.64D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.41	0.21	0.15
Black	0.95	0.57	0.42
American Indian or Alaska Native	4.97	3.19	2.07
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	3.78	0.33	0.07
Filipino	4.14	1.27	0.41
Japanese	4.70	2.36	2.21
Asian Indian	3.43	1.42	*
Korean	*	*	*
Vietnamese	*	1.98	1.73
HISPANIC			
Mexican	1.33	0.62	0.46
Puerto Rican	3.79	2.68	2.01
Central or South American	2.39	0.85	0.50
Cuban	4.58	1.93	0.64

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.65C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	541	300	219
Black	177	113	81
American Indian or Alaska Native	43	27	18
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	43	9	2
Filipino	49	17	7
Japanese	25	13	12
Asian Indian	42	18	13
Korean	*	15	14
Vietnamese	22	9	8
HISPANIC			
Mexican	144	69	50
Puerto Rican	68	47	36
Central or South American	58	23	14
Cuban	46	19	8

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.65D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.37	0.20	0.15
Black	0.81	0.52	0.37
American Indian or Alaska Native	4.35	2.77	1.79
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	3.31	0.70	0.16
Filipino	3.67	1.25	0.51
Japanese	4.30	2.19	2.10
Asian Indian	3.03	1.32	0.94
Korean	*	2.83	2.61
Vietnamese	4.14	1.68	1.48
HISPANIC			
Mexican	1.11	0.54	0.39
Puerto Rican	3.15	2.16	1.67
Central or South American	2.04	0.80	0.50
Cuban	4.42	1.88	0.79

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.66C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	520	267	188
Black	178	100	74
American Indian or Alaska Native	32	21	17
Native Hawaiian	*	*	4
Other Pacific Islander	*	*	*
Chinese	29	8	3
Filipino	37	18	8
Japanese	25	9	7
Asian Indian	29	6	5
Korean	29	10	9
Vietnamese	22	10	9
HISPANIC			
Mexican	134	58	38
Puerto Rican	61	29	24
Central or South American	53	19	11
Cuban	39	12	8

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.66D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.32	0.16	0.12
Black	0.71	0.40	0.30
American Indian or Alaska Native	2.84	1.84	1.49
Native Hawaiian	*	*	3.02
Other Pacific Islander	*	*	*
Chinese	2.04	0.54	0.18
Filipino	2.48	1.17	0.52
Japanese	3.92	1.39	1.18
Asian Indian	1.89	0.42	0.35
Korean	4.61	1.52	1.37
Vietnamese	3.29	1.50	1.33
HISPANIC			
Mexican	0.88	0.38	0.25
Puerto Rican	2.39	1.13	0.95
Central or South American	1.63	0.59	0.32
Cuban	3.46	1.06	0.72

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.67C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	43	38	28
Black	20	16	13
American Indian or Alaska Native	7	7	4
Native Hawaiian	*	*	*
Other Pacific Islander	1	1	1
Chinese	3	3	2
Filipino	4	4	3
Japanese	1	1	0
Asian Indian	2	2	2
Korean	3	3	*
Vietnamese	*	*	*
HISPANIC			
Mexican	16	14	10
Puerto Rican	8	7	5
Central or South American	6	5	4
Cuban	4	3	3

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.67D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 to 17, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.28	0.25	0.18
Black	0.60	0.48	0.38
American Indian or Alaska Native	5.02	5.26	3.15
Native Hawaiian	*	*	*
Other Pacific Islander	2.35	2.11	1.70
Chinese	2.04	2.03	1.35
Filipino	2.44	2.35	1.73
Japanese	2.65	1.71	0.94
Asian Indian	1.70	1.32	1.11
Korean	3.34	3.14	*
Vietnamese	*	*	*
HISPANIC			
Mexican	0.73	0.62	0.45
Puerto Rican	1.98	1.84	1.26
Central or South American	1.48	1.28	0.91
Cuban	4.00	3.52	3.21

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.68C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	84	76	63
Black	34	28	25
American Indian or Alaska Native	6	6	4
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	7	6	2
Filipino	8	6	5
Japanese	*	3	3
Asian Indian	6	5	5
Korean	4	4	*
Vietnamese	4	3	2
HISPANIC			
Mexican	33	23	19
Puerto Rican	12	11	10
Central or South American	12	9	8
Cuban	5	*	*

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.68D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 to 25, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.44	0.40	0.33
Black	0.88	0.73	0.65
American Indian or Alaska Native	3.87	3.62	2.58
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	3.94	3.29	1.00
Filipino	4.20	3.15	2.56
Japanese	*	4.47	3.43
Asian Indian	2.87	2.46	2.18
Korean	4.84	4.66	*
Vietnamese	3.46	2.20	2.05
HISPANIC			
Mexican	1.10	0.77	0.62
Puerto Rican	2.64	2.57	2.25
Central or South American	2.30	1.78	1.46
Cuban	5.38	*	*

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.69C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	501	236	164
Black	175	87	64
American Indian or Alaska Native	31	17	14
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	29	3	1
Filipino	37	15	5
Japanese	23	8	7
Asian Indian	28	3	1
Korean	*	*	*
Vietnamese	*	7	7
HISPANIC			
Mexican	123	47	30
Puerto Rican	59	26	21
Central or South American	50	14	7
Cuban	35	7	5

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.69D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 26 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.39	0.18	0.13
Black	0.97	0.49	0.36
American Indian or Alaska Native	3.71	2.05	1.72
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	2.58	0.30	0.07
Filipino	3.20	1.26	0.41
Japanese	4.48	1.52	1.28
Asian Indian	2.38	0.23	0.10
Korean	*	*	*
Vietnamese	*	1.73	1.73
HISPANIC			
Mexican	1.24	0.48	0.31
Puerto Rican	3.39	1.51	1.21
Central or South American	2.15	0.62	0.29
Cuban	3.72	0.75	0.49

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.70C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	516	260	186
Black	176	98	72
American Indian or Alaska Native	32	18	15
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	29	8	2
Filipino	37	16	7
Japanese	25	9	7
Asian Indian	29	6	5
Korean	*	9	8
Vietnamese	21	8	8
HISPANIC			
Mexican	133	55	36
Puerto Rican	61	28	23
Central or South American	53	18	10
Cuban	38	10	7

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.70D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 or Older, by Racial/Ethnic Subgroups: Annual Averages Based on 1999 and 2000 NHSDAs

Racial/Ethnic Subgroup ¹	TIME PERIOD		
	Lifetime	Past Year	Past Month
NOT HISPANIC			
White	0.35	0.18	0.13
Black	0.81	0.45	0.33
American Indian or Alaska Native	3.25	1.79	1.48
Native Hawaiian	*	*	*
Other Pacific Islander	*	*	*
Chinese	2.22	0.59	0.15
Filipino	2.76	1.20	0.49
Japanese	4.14	1.47	1.25
Asian Indian	2.09	0.45	0.36
Korean	*	1.69	1.53
Vietnamese	3.97	1.46	1.44
HISPANIC			
Mexican	1.03	0.42	0.28
Puerto Rican	2.83	1.27	1.06
Central or South American	1.85	0.64	0.34
Cuban	3.70	0.99	0.68

*Low precision; no estimate reported.

¹ This table is not an exhaustive summary of all racial/ethnic subgroups. Respondents who reported a racial/ethnic subgroup that is rare in the United States, or who reported more than one racial/ethnic subgroup, are not included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.71C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	942	795	508	455	366	332
GEOGRAPHIC DIVISION						
Northeast	458	292	212	196	137	148
New England	216	143	100	132	87	102
Middle Atlantic	405	257	188	146	109	107
Midwest	370	356	179	200	128	144
East North Central	309	294	150	174	106	134
West North Central	203	196	97	99	72	53
South	572	504	277	250	212	185
South Atlantic	467	362	227	188	173	147
East South Central	154	203	96	106	83	69
West South Central	295	283	128	127	90	87
West	464	402	323	257	232	183
Mountain	165	156	126	94	86	60
Pacific	429	370	298	240	216	173
COUNTY TYPE						
Large Metro	736	614	408	336	291	238
Small Metro	475	412	247	256	192	197
250K - 1 Mil. Pop.	422	359	207	206	165	146
<250K Pop.	212	194	127	145	93	121
Nonmetro	340	327	149	161	116	130
Urbanized	185	178	87	92	65	85
Less Urbanized	260	256	124	116	89	83
Completely Rural	102	122	50	54	39	42

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.71D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.43	0.36	0.23	0.20	0.17	0.15
GEOGRAPHIC DIVISION						
Northeast	1.07	0.68	0.50	0.46	0.32	0.35
New England	1.94	1.27	0.89	1.18	0.78	0.91
Middle Atlantic	1.29	0.81	0.60	0.46	0.35	0.34
Midwest	0.72	0.69	0.35	0.39	0.25	0.28
East North Central	0.85	0.81	0.42	0.48	0.29	0.37
West North Central	1.32	1.26	0.63	0.64	0.47	0.34
South	0.73	0.63	0.35	0.31	0.27	0.23
South Atlantic	1.15	0.88	0.56	0.46	0.43	0.36
East South Central	1.12	1.46	0.70	0.76	0.61	0.50
West South Central	1.22	1.15	0.53	0.52	0.37	0.35
West	0.96	0.82	0.67	0.52	0.48	0.37
Mountain	1.18	1.09	0.90	0.66	0.62	0.42
Pacific	1.24	1.06	0.86	0.69	0.63	0.50
COUNTY TYPE						
Large Metro	0.67	0.56	0.37	0.31	0.26	0.22
Small Metro	0.71	0.60	0.37	0.37	0.29	0.29
250K - 1 Mil. Pop.	0.84	0.70	0.41	0.40	0.33	0.29
<250K Pop.	1.27	1.10	0.76	0.82	0.56	0.69
Nonmetro	0.77	0.72	0.34	0.35	0.26	0.28
Urbanized	1.40	1.34	0.66	0.69	0.49	0.64
Less Urbanized	1.04	0.95	0.50	0.43	0.36	0.31
Completely Rural	1.78	2.31	0.87	1.03	0.69	0.81

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.71P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.1639		0.0845		0.8949	
GEOGRAPHIC DIVISION						
Northeast	0.0435		0.6430		0.4852	
New England	0.4202		0.2196		0.8481	
Middle Atlantic	0.0645		0.1260		0.2730	
Midwest	0.0096		0.0255		0.2508	
East North Central	0.2419		0.3015		0.9598	
West North Central	0.0063		0.0166		0.0262	
South	0.7565		0.3469		0.5287	
South Atlantic	0.6488		0.8047		0.7578	
East South Central	0.4940		0.4224		0.3093	
West South Central	0.7335		0.3305		0.9876	
West	0.1004		0.8482		0.3653	
Mountain	0.5763		0.7952		0.9404	
Pacific	0.0459		0.9366		0.3407	
COUNTY TYPE						
Large Metro	0.5333		0.1699		0.6842	
Small Metro	0.2154		0.5711		0.6559	
250K - 1 Mil. Pop.	0.2929		0.3452		0.6649	
<250K Pop.	0.5371		0.6131		0.1233	
Nonmetro	0.5921		0.3757		0.4865	
Urbanized	0.7092		0.7491		0.3402	
Less Urbanized	0.6431		0.3006		0.8108	
Completely Rural	0.8771		0.7291		0.9410	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.72C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	83	86	75	73	53	56
GEOGRAPHIC DIVISION						
Northeast	40	35	39	29	24	19
New England	17	18	18	17	13	10
Middle Atlantic	36	30	34	24	20	16
Midwest	40	38	34	33	25	25
East North Central	30	31	26	28	20	21
West North Central	27	22	21	17	16	15
South	50	55	42	48	30	36
South Atlantic	36	34	27	30	20	23
East South Central	19	29	17	24	12	15
West South Central	29	33	27	29	19	23
West	35	41	35	35	27	29
Mountain	19	20	19	19	14	16
Pacific	30	36	29	29	22	24
COUNTY TYPE						
Large Metro	65	60	59	51	41	41
Small Metro	45	50	40	43	31	33
250K - 1 Mil. Pop.	39	42	36	35	28	27
<250K Pop.	21	28	17	25	13	20
Nonmetro	32	37	27	31	20	24
Urbanized	18	23	16	19	12	14
Less Urbanized	25	27	20	23	14	17
Completely Rural	10	14	9	13	7	10

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.72D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.36	0.37	0.32	0.31	0.23	0.24
GEOGRAPHIC DIVISION						
Northeast	0.96	0.83	0.93	0.68	0.59	0.45
New England	1.51	1.67	1.65	1.52	1.18	0.92
Middle Atlantic	1.18	0.96	1.10	0.76	0.66	0.52
Midwest	0.73	0.70	0.61	0.60	0.46	0.47
East North Central	0.79	0.83	0.69	0.74	0.52	0.56
West North Central	1.56	1.30	1.27	1.01	0.94	0.86
South	0.60	0.66	0.50	0.58	0.37	0.43
South Atlantic	0.89	0.83	0.67	0.73	0.50	0.56
East South Central	1.33	2.01	1.20	1.68	0.87	1.08
West South Central	1.04	1.17	0.95	1.03	0.67	0.80
West	0.66	0.76	0.65	0.64	0.50	0.54
Mountain	1.18	1.26	1.21	1.19	0.90	1.00
Pacific	0.79	0.95	0.77	0.76	0.60	0.64
COUNTY TYPE						
Large Metro	0.56	0.53	0.51	0.45	0.35	0.36
Small Metro	0.65	0.70	0.58	0.60	0.44	0.46
250K - 1 Mil. Pop.	0.75	0.78	0.67	0.66	0.53	0.50
<250K Pop.	1.22	1.53	0.99	1.37	0.76	1.09
Nonmetro	0.68	0.78	0.57	0.65	0.43	0.49
Urbanized	1.22	1.72	1.04	1.42	0.81	1.05
Less Urbanized	0.95	0.95	0.75	0.80	0.53	0.58
Completely Rural	1.83	2.28	1.69	2.24	1.31	1.63

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.72P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 12 to 17, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.1413		0.0119		0.8229	
GEOGRAPHIC DIVISION						
Northeast	0.7556		0.5887		0.7048	
New England	0.6009		0.8554		0.8852	
Middle Atlantic	0.5054		0.5918		0.7141	
Midwest	0.2051		0.0597		0.1934	
East North Central	0.1917		0.0766		0.1218	
West North Central	0.6996		0.4457		0.9734	
South	0.9269		0.6601		0.3531	
South Atlantic	0.4376		0.7996		0.4566	
East South Central	0.5519		0.5011		0.0559	
West South Central	0.1710		0.1640		0.5947	
West	0.0239		0.0076		0.9612	
Mountain	0.2542		0.6262		0.7985	
Pacific	0.0494		0.0040		0.8285	
COUNTY TYPE						
Large Metro	0.0999		0.0145		0.7918	
Small Metro	0.7309		0.4504		0.9451	
250K - 1 Mil. Pop.	0.2132		0.2039		0.4311	
<250K Pop.	0.1695		0.5193		0.2324	
Nonmetro	0.8652		0.4938		1.0000	
Urbanized	0.5736		0.2805		0.1596	
Less Urbanized	0.3203		0.0828		0.3620	
Completely Rural	0.2445		0.7399		0.9165	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.73C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	154	150	137	132	115	105
GEOGRAPHIC DIVISION						
Northeast	69	54	66	55	53	43
New England	38	27	39	26	35	22
Middle Atlantic	58	47	53	49	39	38
Midwest	68	63	65	55	52	46
East North Central	56	49	53	46	43	40
West North Central	38	41	37	30	29	22
South	93	83	73	71	62	57
South Atlantic	57	61	53	48	46	39
East South Central	38	34	31	32	28	27
West South Central	62	45	40	42	30	30
West	73	93	67	80	61	62
Mountain	29	30	30	30	31	28
Pacific	67	88	60	73	52	56
COUNTY TYPE						
Large Metro	111	118	98	101	87	78
Small Metro	87	66	81	69	65	57
250K - 1 Mil. Pop.	65	55	58	57	46	48
<250K Pop.	58	37	55	38	45	32
Nonmetro	57	61	51	47	39	39
Urbanized	35	30	30	27	23	22
Less Urbanized	47	49	40	37	29	29
Completely Rural	18	15	14	13	12	9

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.73D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.54	0.52	0.48	0.46	0.40	0.36
GEOGRAPHIC DIVISION						
Northeast	1.38	1.06	1.31	1.08	1.05	0.85
New England	2.95	2.05	3.02	1.96	2.68	1.64
Middle Atlantic	1.54	1.25	1.41	1.28	1.03	0.99
Midwest	1.00	0.92	0.95	0.80	0.77	0.66
East North Central	1.18	1.02	1.11	0.96	0.91	0.84
West North Central	1.86	1.94	1.78	1.45	1.44	1.02
South	0.92	0.81	0.72	0.69	0.61	0.55
South Atlantic	1.17	1.23	1.09	0.97	0.94	0.80
East South Central	2.10	1.83	1.68	1.75	1.55	1.48
West South Central	1.79	1.27	1.16	1.18	0.87	0.85
West	1.12	1.40	1.04	1.19	0.93	0.93
Mountain	1.50	1.51	1.55	1.51	1.57	1.37
Pacific	1.47	1.89	1.32	1.58	1.15	1.20
COUNTY TYPE						
Large Metro	0.80	0.85	0.70	0.73	0.62	0.56
Small Metro	0.95	0.70	0.88	0.73	0.71	0.60
250K - 1 Mil. Pop.	1.02	0.83	0.90	0.86	0.71	0.72
<250K Pop.	2.10	1.28	2.00	1.32	1.63	1.12
Nonmetro	1.07	1.08	0.95	0.85	0.72	0.70
Urbanized	1.79	1.61	1.54	1.43	1.17	1.17
Less Urbanized	1.60	1.52	1.36	1.14	1.01	0.90
Completely Rural	3.34	3.06	2.62	2.71	2.26	1.94

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.73P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 to 25, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0542		0.0571		0.3377	
GEOGRAPHIC DIVISION						
Northeast	0.4216		0.4504		0.6588	
New England	0.8068		0.6180		0.7972	
Middle Atlantic	0.2546		0.2360		0.7106	
Midwest	0.2788		0.0189		0.1679	
East North Central	0.9260		0.0377		0.5719	
West North Central	0.0738		0.2531		0.1167	
South	0.0892		0.4739		0.7315	
South Atlantic	0.2002		0.5517		0.9334	
East South Central	0.5698		0.7539		0.1841	
West South Central	0.3271		0.7706		0.6545	
West	0.7675		0.8371		0.5510	
Mountain	0.7802		0.8309		0.8331	
Pacific	0.6656		0.7455		0.4367	
COUNTY TYPE						
Large Metro	0.4458		0.2888		0.2770	
Small Metro	0.3710		0.2959		0.8957	
250K - 1 Mil. Pop.	0.2035		0.2281		0.9578	
<250K Pop.	0.7833		0.9318		0.7562	
Nonmetro	0.0211		0.1996		0.9855	
Urbanized	0.2527		0.4224		0.5782	
Less Urbanized	0.0736		0.2651		0.7965	
Completely Rural	0.7588		0.8252		0.7001	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.74C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	898	751	451	398	332	300
GEOGRAPHIC DIVISION						
Northeast	440	274	192	173	132	133
New England	225	127	91	117	85	90
Middle Atlantic	379	242	170	127	103	98
Midwest	347	329	151	167	112	122
East North Central	289	271	125	143	94	113
West North Central	191	185	85	86	60	46
South	542	480	251	227	192	168
South Atlantic	449	342	210	180	161	138
East South Central	142	206	86	92	71	55
West South Central	269	264	106	102	74	76
West	451	380	285	220	211	171
Mountain	160	141	115	79	77	58
Pacific	417	353	261	206	196	161
COUNTY TYPE						
Large Metro	711	575	371	295	271	222
Small Metro	453	390	216	215	171	170
250K - 1 Mil. Pop.	406	345	192	175	157	126
<250K Pop.	203	177	101	120	73	105
Nonmetro	322	318	133	140	103	114
Urbanized	172	175	73	81	56	74
Less Urbanized	245	246	108	105	80	77
Completely Rural	100	114	45	43	37	38

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.74D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.53	0.44	0.27	0.23	0.20	0.18
GEOGRAPHIC DIVISION						
Northeast	1.31	0.82	0.57	0.52	0.39	0.40
New England	2.57	1.45	1.04	1.33	0.97	1.03
Middle Atlantic	1.53	0.98	0.69	0.51	0.42	0.40
Midwest	0.88	0.83	0.38	0.42	0.28	0.31
East North Central	1.05	0.98	0.45	0.52	0.34	0.41
West North Central	1.63	1.57	0.73	0.73	0.52	0.39
South	0.90	0.79	0.42	0.37	0.32	0.28
South Atlantic	1.42	1.07	0.66	0.56	0.51	0.43
East South Central	1.35	1.94	0.82	0.87	0.68	0.52
West South Central	1.50	1.45	0.59	0.56	0.41	0.42
West	1.23	1.02	0.78	0.60	0.58	0.46
Mountain	1.54	1.32	1.11	0.74	0.74	0.55
Pacific	1.59	1.34	1.00	0.78	0.75	0.61
COUNTY TYPE						
Large Metro	0.84	0.69	0.44	0.35	0.32	0.26
Small Metro	0.90	0.75	0.43	0.41	0.34	0.33
250K - 1 Mil. Pop.	1.06	0.88	0.50	0.45	0.41	0.32
<250K Pop.	1.66	1.36	0.82	0.92	0.60	0.81
Nonmetro	0.95	0.90	0.39	0.40	0.30	0.32
Urbanized	1.76	1.73	0.75	0.80	0.58	0.74
Less Urbanized	1.25	1.18	0.55	0.50	0.41	0.37
Completely Rural	2.15	2.72	0.97	1.03	0.80	0.90

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.74P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 26 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3184		0.3454		0.6538	
GEOGRAPHIC DIVISION						
Northeast	0.0505		0.8211		0.5833	
New England	0.4496		0.2687		0.8074	
Middle Atlantic	0.0683		0.2387		0.3426	
Midwest	0.0151		0.1895		0.6136	
East North Central	0.2680		0.9704		0.6399	
West North Central	0.0091		0.0370		0.0608	
South	0.9525		0.4486		0.7023	
South Atlantic	0.7035		0.8570		0.8240	
East South Central	0.4717		0.3412		0.8543	
West South Central	0.9819		0.5090		0.7452	
West	0.0459		0.8586		0.2495	
Mountain	0.5970		0.7851		0.9578	
Pacific	0.0175		0.7182		0.1988	
COUNTY TYPE						
Large Metro	0.7021		0.3926		0.9414	
Small Metro	0.2708		0.8782		0.5993	
250K - 1 Mil. Pop.	0.4406		0.6376		0.7059	
<250K Pop.	0.3718		0.5965		0.0958	
Nonmetro	0.8838		0.6587		0.4335	
Urbanized	0.7941		0.9485		0.3893	
Less Urbanized	0.9340		0.5218		0.6975	
Completely Rural	0.9289		0.9369		0.9567	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.75C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	927	785	492	442	360	322
GEOGRAPHIC DIVISION						
Northeast	455	290	208	191	139	147
New England	219	138	104	126	94	100
Middle Atlantic	399	257	182	146	105	109
Midwest	360	357	174	188	126	136
East North Central	302	294	145	162	104	126
West North Central	196	198	96	95	71	51
South	563	493	270	248	211	181
South Atlantic	464	357	225	189	174	146
East South Central	148	198	91	103	82	67
West South Central	287	273	117	123	85	83
West	457	396	310	249	225	176
Mountain	162	155	116	85	82	55
Pacific	423	364	288	234	210	167
COUNTY TYPE						
Large Metro	726	607	397	327	287	232
Small Metro	470	402	242	248	191	191
250K - 1 Mil. Pop.	420	352	203	199	165	141
<250K Pop.	210	190	125	142	92	118
Nonmetro	338	321	142	155	113	125
Urbanized	184	177	86	89	63	81
Less Urbanized	255	251	117	114	86	83
Completely Rural	101	117	46	48	37	38

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.75D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.47	0.39	0.25	0.22	0.18	0.16
GEOGRAPHIC DIVISION						
Northeast	1.18	0.75	0.54	0.50	0.36	0.38
New England	2.18	1.37	1.03	1.25	0.94	0.99
Middle Atlantic	1.40	0.90	0.64	0.51	0.37	0.38
Midwest	0.78	0.77	0.38	0.41	0.27	0.29
East North Central	0.93	0.91	0.45	0.50	0.32	0.39
West North Central	1.43	1.42	0.70	0.69	0.52	0.37
South	0.80	0.69	0.38	0.35	0.30	0.25
South Atlantic	1.27	0.97	0.62	0.51	0.48	0.39
East South Central	1.20	1.59	0.74	0.83	0.66	0.54
West South Central	1.34	1.26	0.55	0.56	0.40	0.38
West	1.06	0.91	0.72	0.57	0.52	0.40
Mountain	1.31	1.22	0.94	0.67	0.66	0.44
Pacific	1.38	1.17	0.94	0.75	0.68	0.54
COUNTY TYPE						
Large Metro	0.73	0.62	0.40	0.34	0.29	0.24
Small Metro	0.79	0.65	0.41	0.40	0.32	0.31
250K - 1 Mil. Pop.	0.94	0.77	0.45	0.44	0.37	0.31
<250K Pop.	1.41	1.20	0.84	0.90	0.61	0.74
Nonmetro	0.86	0.79	0.36	0.38	0.29	0.31
Urbanized	1.57	1.48	0.73	0.74	0.54	0.68
Less Urbanized	1.14	1.04	0.52	0.47	0.38	0.34
Completely Rural	1.96	2.50	0.89	1.03	0.71	0.82

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.75P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Among Persons Aged 18 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2051		0.1755		0.8582	
GEOGRAPHIC DIVISION						
Northeast	0.0448		0.7008		0.5172	
New England	0.4730		0.2059		0.8362	
Middle Atlantic	0.0592		0.1444		0.2876	
Midwest	0.0122		0.0503		0.3482	
East North Central	0.2979		0.4671		0.8017	
West North Central	0.0049		0.0204		0.0202	
South	0.7541		0.3833		0.6293	
South Atlantic	0.5980		0.7786		0.8331	
East South Central	0.5465		0.3278		0.5117	
West South Central	0.8433		0.5091		0.8917	
West	0.0632		0.8891		0.3408	
Mountain	0.6565		0.8499		0.9686	
Pacific	0.0286		0.7988		0.3095	
COUNTY TYPE						
Large Metro	0.6322		0.3077		0.7105	
Small Metro	0.2251		0.6535		0.6349	
250K - 1 Mil. Pop.	0.3513		0.4608		0.7690	
<250K Pop.	0.4202		0.6906		0.1657	
Nonmetro	0.5979		0.4305		0.4657	
Urbanized	0.6122		0.7108		0.4038	
Less Urbanized	0.7196		0.4116		0.7146	
Completely Rural	0.9645		0.8837		0.8784	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug indicates use at least once of marijuana/hashish, cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.76C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	872	775	428	398	295	281
GEOGRAPHIC DIVISION						
Northeast	445	286	187	163	107	123
New England	245	131	96	106	70	86
Middle Atlantic	377	257	161	127	81	89
Midwest	355	342	144	157	107	113
East North Central	291	289	122	140	86	105
West North Central	202	180	77	73	64	44
South	523	492	229	227	172	162
South Atlantic	433	336	195	189	144	139
East South Central	146	207	77	85	60	52
West South Central	259	289	94	91	70	64
West	403	393	274	236	188	157
Mountain	153	140	99	78	81	59
Pacific	369	367	256	223	170	146
COUNTY TYPE						
Large Metro	677	593	341	311	231	202
Small Metro	458	402	208	219	154	168
250K - 1 Mil. Pop.	405	351	176	176	126	130
<250K Pop.	215	195	113	119	88	101
Nonmetro	311	339	128	136	93	117
Urbanized	171	177	75	78	53	75
Less Urbanized	243	254	95	94	68	73
Completely Rural	90	117	47	51	36	42

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.76D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.39	0.35	0.19	0.18	0.13	0.13
GEOGRAPHIC DIVISION						
Northeast	1.04	0.67	0.44	0.38	0.25	0.29
New England	2.20	1.17	0.87	0.95	0.63	0.77
Middle Atlantic	1.19	0.81	0.51	0.40	0.26	0.28
Midwest	0.69	0.66	0.28	0.30	0.21	0.22
East North Central	0.81	0.80	0.34	0.38	0.24	0.29
West North Central	1.31	1.16	0.50	0.47	0.41	0.28
South	0.67	0.62	0.29	0.29	0.22	0.20
South Atlantic	1.07	0.82	0.48	0.46	0.36	0.34
East South Central	1.06	1.50	0.56	0.61	0.43	0.37
West South Central	1.07	1.18	0.39	0.37	0.29	0.26
West	0.83	0.80	0.57	0.48	0.39	0.32
Mountain	1.10	0.98	0.71	0.54	0.58	0.41
Pacific	1.07	1.06	0.74	0.64	0.49	0.42
COUNTY TYPE						
Large Metro	0.61	0.54	0.31	0.28	0.21	0.18
Small Metro	0.69	0.59	0.31	0.32	0.23	0.24
250K - 1 Mil. Pop.	0.81	0.69	0.35	0.35	0.25	0.25
<250K Pop.	1.29	1.10	0.68	0.68	0.53	0.57
Nonmetro	0.71	0.74	0.29	0.30	0.21	0.26
Urbanized	1.30	1.33	0.57	0.58	0.40	0.56
Less Urbanized	0.97	0.94	0.38	0.35	0.27	0.27
Completely Rural	1.58	2.22	0.83	0.96	0.63	0.80

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.76P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.4517		0.1936		0.6920	
GEOGRAPHIC DIVISION						
Northeast	0.1037		0.9188		0.3921	
New England	0.4273		0.0494		0.1141	
Middle Atlantic	0.1548		0.1099		0.7292	
Midwest	0.0746		0.3407		0.5009	
East North Central	0.3539		0.5825		0.9815	
West North Central	0.0840		0.3782		0.2198	
South	0.9575		0.2477		0.7731	
South Atlantic	0.6868		0.7547		0.7921	
East South Central	0.5863		0.1943		0.0999	
West South Central	0.8963		0.2203		0.7793	
West	0.1034		0.6994		0.7954	
Mountain	0.1849		1.0000		0.6658	
Pacific	0.0248		0.6732		0.6280	
COUNTY TYPE						
Large Metro	0.8458		0.3986		0.8025	
Small Metro	0.2310		0.3498		0.9602	
250K - 1 Mil. Pop.	0.2716		0.2494		0.8135	
<250K Pop.	0.6736		0.8923		0.6591	
Nonmetro	0.8384		0.8274		0.1134	
Urbanized	0.8703		0.7863		0.0758	
Less Urbanized	0.8828		0.6259		0.5936	
Completely Rural	0.9838		0.9788		0.6622	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.77C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	74	72	66	62	47	49
GEOGRAPHIC DIVISION						
Northeast	33	28	31	24	20	17
New England	15	17	16	15	12	10
Middle Atlantic	29	23	27	20	16	14
Midwest	35	32	30	29	24	24
East North Central	26	26	24	24	17	19
West North Central	24	20	19	17	16	15
South	43	45	38	38	26	29
South Atlantic	30	27	26	25	19	19
East South Central	18	24	16	19	11	14
West South Central	26	27	22	22	15	17
West	35	37	32	32	24	26
Mountain	21	20	18	18	14	13
Pacific	28	31	27	27	19	22
COUNTY TYPE						
Large Metro	57	51	51	44	36	36
Small Metro	41	43	36	37	26	29
250K - 1 Mil. Pop.	36	36	32	30	24	24
<250K Pop.	18	25	15	22	10	17
Nonmetro	26	31	22	28	17	22
Urbanized	17	18	14	17	10	13
Less Urbanized	19	22	16	19	11	15
Completely Rural	10	12	8	11	7	10

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.77D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.32	0.31	0.29	0.27	0.20	0.21
GEOGRAPHIC DIVISION						
Northeast	0.79	0.67	0.75	0.58	0.48	0.40
New England	1.39	1.51	1.45	1.31	1.06	0.86
Middle Atlantic	0.94	0.75	0.87	0.64	0.52	0.45
Midwest	0.64	0.59	0.56	0.53	0.43	0.44
East North Central	0.69	0.69	0.62	0.63	0.46	0.50
West North Central	1.40	1.16	1.13	1.01	0.94	0.88
South	0.53	0.54	0.46	0.45	0.32	0.35
South Atlantic	0.75	0.66	0.65	0.60	0.46	0.48
East South Central	1.25	1.66	1.10	1.32	0.77	0.95
West South Central	0.91	0.96	0.78	0.77	0.55	0.62
West	0.65	0.68	0.60	0.59	0.45	0.48
Mountain	1.29	1.20	1.12	1.09	0.87	0.83
Pacific	0.75	0.83	0.71	0.71	0.52	0.59
COUNTY TYPE						
Large Metro	0.50	0.45	0.44	0.39	0.31	0.32
Small Metro	0.59	0.59	0.52	0.51	0.38	0.40
250K - 1 Mil. Pop.	0.68	0.67	0.61	0.56	0.46	0.44
<250K Pop.	1.07	1.37	0.89	1.21	0.60	0.96
Nonmetro	0.56	0.65	0.48	0.58	0.37	0.46
Urbanized	1.15	1.39	0.93	1.29	0.65	0.99
Less Urbanized	0.74	0.76	0.62	0.67	0.43	0.52
Completely Rural	1.72	2.02	1.51	1.86	1.22	1.64

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.77P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3569		0.0503		0.8917	
GEOGRAPHIC DIVISION						
Northeast	0.2188		0.6907		0.5817	
New England	0.8343		0.4227		0.5882	
Middle Atlantic	0.1898		0.9493		0.7870	
Midwest	0.2348		0.2574		0.4427	
East North Central	0.1661		0.1766		0.2512	
West North Central	0.8025		0.9069		0.8565	
South	0.6992		0.1640		0.8139	
South Atlantic	0.6186		0.6256		0.8109	
East South Central	0.3568		0.5926		0.0395	
West South Central	0.0657		0.0345		0.4726	
West	0.1699		0.1005		0.7868	
Mountain	0.7790		0.8942		0.7806	
Pacific	0.1490		0.0631		0.8856	
COUNTY TYPE						
Large Metro	0.1440		0.0223		0.4679	
Small Metro	0.8486		0.6554		0.9125	
250K - 1 Mil. Pop.	0.2322		0.2385		0.3335	
<250K Pop.	0.0846		0.2910		0.1852	
Nonmetro	0.4454		0.9178		0.2541	
Urbanized	0.1202		0.0493		0.0251	
Less Urbanized	0.9061		0.1475		0.5938	
Completely Rural	0.5767		0.5268		0.4175	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.78C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	149	150	131	126	109	98
GEOGRAPHIC DIVISION						
Northeast	69	57	59	53	51	44
New England	38	27	37	26	33	21
Middle Atlantic	57	50	46	45	38	39
Midwest	63	65	63	53	50	43
East North Central	53	53	52	46	41	38
West North Central	34	38	35	26	29	20
South	92	81	67	66	58	51
South Atlantic	58	58	51	46	44	36
East South Central	36	31	25	27	24	22
West South Central	61	46	37	38	29	27
West	68	92	69	77	58	57
Mountain	27	31	27	28	29	23
Pacific	63	87	63	72	50	52
COUNTY TYPE						
Large Metro	108	117	93	97	83	74
Small Metro	87	67	81	64	61	51
250K - 1 Mil. Pop.	64	54	56	52	42	43
<250K Pop.	59	40	57	36	43	30
Nonmetro	54	59	44	45	35	37
Urbanized	36	30	27	25	21	22
Less Urbanized	43	46	36	35	29	27
Completely Rural	16	15	12	12	7	9

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.78D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.52	0.52	0.46	0.43	0.38	0.34
GEOGRAPHIC DIVISION						
Northeast	1.37	1.12	1.18	1.03	1.02	0.86
New England	2.98	2.05	2.84	1.95	2.59	1.62
Middle Atlantic	1.53	1.33	1.23	1.20	1.00	1.01
Midwest	0.93	0.94	0.93	0.77	0.73	0.62
East North Central	1.12	1.10	1.10	0.97	0.85	0.80
West North Central	1.66	1.82	1.68	1.24	1.40	0.94
South	0.91	0.79	0.67	0.64	0.57	0.49
South Atlantic	1.21	1.18	1.05	0.94	0.91	0.74
East South Central	1.96	1.70	1.35	1.50	1.33	1.22
West South Central	1.77	1.31	1.05	1.08	0.83	0.77
West	1.05	1.38	1.06	1.15	0.89	0.85
Mountain	1.38	1.51	1.38	1.37	1.50	1.16
Pacific	1.38	1.87	1.38	1.54	1.10	1.12
COUNTY TYPE						
Large Metro	0.78	0.84	0.67	0.70	0.60	0.53
Small Metro	0.95	0.70	0.88	0.68	0.67	0.54
250K - 1 Mil. Pop.	0.99	0.81	0.87	0.78	0.66	0.65
<250K Pop.	2.15	1.41	2.07	1.28	1.56	1.05
Nonmetro	1.00	1.05	0.83	0.80	0.65	0.66
Urbanized	1.88	1.58	1.38	1.34	1.07	1.15
Less Urbanized	1.49	1.44	1.22	1.09	0.99	0.83
Completely Rural	3.04	3.13	2.34	2.44	1.32	1.77

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.78P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.1309		0.1937		0.2250	
GEOGRAPHIC DIVISION						
Northeast	0.5316		0.6372		0.5194	
New England	0.5879		0.4283		0.8354	
Middle Atlantic	0.2730		0.3008		0.5262	
Midwest	0.2814		0.1242		0.2824	
East North Central	0.6928		0.0994		0.6725	
West North Central	0.2083		0.7311		0.2118	
South	0.1965		0.4949		0.9776	
South Atlantic	0.4413		0.4487		0.6165	
East South Central	0.7201		0.5479		0.1210	
West South Central	0.3233		0.4899		0.5063	
West	0.9381		0.9177		0.4194	
Mountain	0.8180		0.9779		0.9014	
Pacific	0.8616		0.9063		0.4067	
COUNTY TYPE						
Large Metro	0.7560		0.3806		0.1767	
Small Metro	0.2559		0.3654		0.4242	
250K - 1 Mil. Pop.	0.0604		0.1900		0.4938	
<250K Pop.	0.5299		0.7685		0.6767	
Nonmetro	0.0568		0.8345		0.3448	
Urbanized	0.1848		0.7292		0.2575	
Less Urbanized	0.1751		0.9324		0.9161	
Completely Rural	0.7125		0.7384		0.4454	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.79C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	841	734	379	339	261	248
GEOGRAPHIC DIVISION						
Northeast	434	268	165	141	84	111
New England	248	117	80	93	51	78
Middle Atlantic	358	242	145	107	67	79
Midwest	335	316	112	120	86	89
East North Central	275	266	92	107	73	83
West North Central	190	171	65	56	47	34
South	506	470	214	198	158	144
South Atlantic	428	320	184	172	140	126
East South Central	142	209	70	70	48	41
West South Central	234	269	81	65	54	54
West	393	371	242	202	170	143
Mountain	150	125	88	67	70	52
Pacific	359	350	226	190	155	133
COUNTY TYPE						
Large Metro	665	557	312	269	209	182
Small Metro	437	380	178	178	134	143
250K - 1 Mil. Pop.	387	336	159	145	116	111
<250K Pop.	210	178	82	93	69	81
Nonmetro	296	330	111	113	82	102
Urbanized	157	173	63	71	47	67
Less Urbanized	232	249	82	76	58	66
Completely Rural	90	106	45	41	36	37

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.79D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.50	0.43	0.22	0.20	0.15	0.15
GEOGRAPHIC DIVISION						
Northeast	1.30	0.80	0.49	0.42	0.25	0.33
New England	2.82	1.34	0.91	1.06	0.58	0.89
Middle Atlantic	1.45	0.98	0.59	0.43	0.27	0.32
Midwest	0.85	0.80	0.29	0.30	0.22	0.22
East North Central	1.00	0.96	0.33	0.39	0.26	0.30
West North Central	1.63	1.45	0.56	0.48	0.40	0.29
South	0.84	0.77	0.36	0.32	0.26	0.24
South Atlantic	1.35	1.00	0.58	0.54	0.44	0.39
East South Central	1.36	1.97	0.67	0.66	0.45	0.39
West South Central	1.30	1.48	0.45	0.36	0.30	0.29
West	1.07	1.00	0.66	0.55	0.47	0.39
Mountain	1.44	1.17	0.84	0.62	0.67	0.48
Pacific	1.37	1.33	0.86	0.72	0.59	0.50
COUNTY TYPE						
Large Metro	0.78	0.66	0.37	0.32	0.25	0.22
Small Metro	0.87	0.73	0.35	0.34	0.27	0.27
250K - 1 Mil. Pop.	1.01	0.86	0.42	0.37	0.30	0.29
<250K Pop.	1.72	1.37	0.67	0.72	0.57	0.63
Nonmetro	0.87	0.94	0.33	0.32	0.24	0.29
Urbanized	1.61	1.72	0.65	0.71	0.48	0.66
Less Urbanized	1.19	1.19	0.42	0.36	0.30	0.31
Completely Rural	1.93	2.53	0.97	0.98	0.78	0.89

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.79P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.6525		0.4785		0.3787	
GEOGRAPHIC DIVISION						
Northeast	0.1038		0.9314		0.2967	
New England	0.3775		0.1003		0.0989	
Middle Atlantic	0.1679		0.1695		0.8912	
Midwest	0.1100		0.8658		0.9092	
East North Central	0.4306		0.6679		0.6397	
West North Central	0.1026		0.3603		0.2853	
South	0.7622		0.4149		0.8168	
South Atlantic	0.6225		0.9140		0.9205	
East South Central	0.5489		0.1056		0.4956	
West South Central	0.7976		0.5918		0.7973	
West	0.0601		0.8386		0.5468	
Mountain	0.1366		0.9907		0.7138	
Pacific	0.0103		0.8197		0.3923	
COUNTY TYPE						
Large Metro	0.9665		0.7423		0.7354	
Small Metro	0.3030		0.5239		0.6996	
250K - 1 Mil. Pop.	0.4674		0.5283		0.8710	
<250K Pop.	0.4418		0.8711		0.6355	
Nonmetro	0.6604		0.8275		0.2329	
Urbanized	0.8293		0.7254		0.1805	
Less Urbanized	0.7025		0.6901		0.5885	
Completely Rural	0.9287		0.7491		0.9578	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.80C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	868	771	419	387	290	273
GEOGRAPHIC DIVISION						
Northeast	446	294	188	161	108	123
New England	244	132	101	104	75	84
Middle Atlantic	376	264	159	125	78	91
Midwest	350	341	142	146	107	105
East North Central	289	290	118	129	85	97
West North Central	197	178	80	69	64	41
South	522	485	224	224	171	158
South Atlantic	434	337	192	189	146	137
East South Central	144	202	73	81	57	48
West South Central	256	279	89	86	68	62
West	401	387	266	229	181	152
Mountain	150	137	93	69	75	52
Pacific	367	362	250	219	165	143
COUNTY TYPE						
Large Metro	677	592	337	303	226	196
Small Metro	454	394	204	210	154	161
250K - 1 Mil. Pop.	402	347	172	171	125	124
<250K Pop.	212	190	110	113	88	97
Nonmetro	311	334	123	129	92	112
Urbanized	170	176	74	75	52	72
Less Urbanized	242	251	91	92	67	73
Completely Rural	90	111	46	46	36	38

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.80D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.44	0.39	0.21	0.19	0.15	0.14
GEOGRAPHIC DIVISION						
Northeast	1.16	0.76	0.49	0.42	0.28	0.32
New England	2.43	1.30	1.00	1.03	0.74	0.83
Middle Atlantic	1.32	0.92	0.56	0.44	0.27	0.32
Midwest	0.76	0.74	0.31	0.31	0.23	0.23
East North Central	0.89	0.89	0.36	0.40	0.26	0.30
West North Central	1.44	1.28	0.58	0.50	0.47	0.30
South	0.74	0.68	0.32	0.31	0.24	0.22
South Atlantic	1.19	0.91	0.53	0.51	0.40	0.37
East South Central	1.17	1.62	0.60	0.65	0.46	0.38
West South Central	1.20	1.29	0.42	0.40	0.32	0.29
West	0.93	0.89	0.62	0.52	0.42	0.35
Mountain	1.21	1.08	0.75	0.54	0.61	0.41
Pacific	1.19	1.17	0.81	0.70	0.54	0.46
COUNTY TYPE						
Large Metro	0.68	0.61	0.34	0.31	0.23	0.20
Small Metro	0.76	0.64	0.34	0.34	0.26	0.26
250K - 1 Mil. Pop.	0.90	0.76	0.38	0.37	0.28	0.27
<250K Pop.	1.42	1.20	0.74	0.71	0.59	0.61
Nonmetro	0.79	0.82	0.31	0.32	0.23	0.27
Urbanized	1.46	1.47	0.63	0.63	0.44	0.60
Less Urbanized	1.08	1.04	0.40	0.38	0.30	0.30
Completely Rural	1.75	2.37	0.88	0.98	0.69	0.80

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.80P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.4973		0.3217		0.6625	
GEOGRAPHIC DIVISION						
Northeast	0.0942		0.8620		0.4534	
New England	0.4278		0.0679		0.1368	
Middle Atlantic	0.1387		0.1141		0.6885	
Midwest	0.0897		0.4680		0.5996	
East North Central	0.4178		0.7746		0.7916	
West North Central	0.0769		0.4001		0.1799	
South	0.9358		0.3538		0.7994	
South Atlantic	0.7163		0.8038		0.8167	
East South Central	0.5152		0.1502		0.1931	
West South Central	0.9643		0.4797		0.9441	
West	0.0807		0.8606		0.7550	
Mountain	0.1793		0.9681		0.6798	
Pacific	0.0182		0.8390		0.6037	
COUNTY TYPE						
Large Metro	0.9376		0.6163		0.8947	
Small Metro	0.2371		0.3762		0.9415	
250K - 1 Mil. Pop.	0.3239		0.3388		0.9595	
<250K Pop.	0.5411		0.9283		0.8430	
Nonmetro	0.8926		0.8104		0.1501	
Urbanized	0.9375		0.9737		0.1318	
Less Urbanized	0.8529		0.8017		0.5398	
Completely Rural	0.9790		0.8403		0.8177	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.81C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	596	519	188	150	130	112
GEOGRAPHIC DIVISION						
Northeast	284	211	86	66	73	39
New England	158	93	64	38	54	23
Middle Atlantic	239	192	59	54	49	31
Midwest	213	191	62	58	44	36
East North Central	174	167	49	50	29	30
West North Central	124	93	37	30	33	20
South	360	300	123	71	78	44
South Atlantic	297	226	88	58	58	35
East South Central	119	119	61	25	35	15
West South Central	164	155	60	34	39	23
West	319	302	95	98	59	88
Mountain	103	105	30	40	23	20
Pacific	302	283	90	89	55	86
COUNTY TYPE						
Large Metro	478	406	137	113	101	95
Small Metro	317	259	113	90	77	53
250K - 1 Mil. Pop.	288	225	105	82	75	49
<250K Pop.	127	127	41	33	19	16
Nonmetro	179	192	57	44	36	24
Urbanized	95	120	41	20	27	11
Less Urbanized	137	131	36	32	25	18
Completely Rural	59	65	14	20	2	11

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.81D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.27	0.23	0.08	0.07	0.06	0.05
GEOGRAPHIC DIVISION						
Northeast	0.66	0.49	0.20	0.15	0.17	0.09
New England	1.41	0.83	0.57	0.34	0.48	0.21
Middle Atlantic	0.76	0.61	0.19	0.17	0.15	0.10
Midwest	0.41	0.37	0.12	0.11	0.09	0.07
East North Central	0.48	0.46	0.14	0.14	0.08	0.08
West North Central	0.80	0.60	0.24	0.19	0.22	0.13
South	0.46	0.38	0.16	0.09	0.10	0.06
South Atlantic	0.73	0.55	0.22	0.14	0.14	0.08
East South Central	0.87	0.86	0.45	0.18	0.25	0.11
West South Central	0.68	0.63	0.25	0.14	0.16	0.09
West	0.66	0.61	0.20	0.20	0.12	0.18
Mountain	0.74	0.73	0.21	0.28	0.16	0.14
Pacific	0.88	0.81	0.26	0.26	0.16	0.25
COUNTY TYPE						
Large Metro	0.43	0.37	0.12	0.10	0.09	0.09
Small Metro	0.47	0.38	0.17	0.13	0.12	0.08
250K - 1 Mil. Pop.	0.58	0.44	0.21	0.16	0.15	0.10
<250K Pop.	0.76	0.72	0.25	0.19	0.11	0.09
Nonmetro	0.41	0.42	0.13	0.10	0.08	0.05
Urbanized	0.72	0.90	0.31	0.15	0.21	0.08
Less Urbanized	0.54	0.48	0.14	0.12	0.10	0.07
Completely Rural	1.03	1.23	0.25	0.37	0.04	0.21

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.81P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3230		0.0660		0.0520	
GEOGRAPHIC DIVISION						
Northeast	0.1139		0.9842		0.0959	
New England	0.9556		0.5388		0.3722	
Middle Atlantic	0.0794		0.5802		0.1455	
Midwest	0.8288		0.4170		0.3553	
East North Central	0.8897		0.6322		0.6336	
West North Central	0.8510		0.4737		0.3984	
South	0.0933		0.0041		0.0146	
South Atlantic	0.5399		0.1089		0.0317	
East South Central	0.2908		0.0706		0.4878	
West South Central	0.0644		0.0705		0.2937	
West	0.1072		0.7808		0.4956	
Mountain	0.0718		0.1496		0.8491	
Pacific	0.3221		0.8126		0.5127	
COUNTY TYPE						
Large Metro	0.3920		0.2542		0.2330	
Small Metro	0.1009		0.1358		0.0948	
250K - 1 Mil. Pop.	0.0731		0.1638		0.0876	
<250K Pop.	0.9571		0.5958		0.9103	
Nonmetro	0.0721		0.8177		0.8664	
Urbanized	0.0101		0.3813		0.5838	
Less Urbanized	0.4933		0.9363		0.8924	
Completely Rural	0.6348		0.4192		0.3097	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.82C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	29	31	24	27	14	17
GEOGRAPHIC DIVISION						
Northeast	11	10	11	8	5	4
New England	5	5	5	4	2	2
Middle Atlantic	9	8	10	7	5	3
Midwest	10	13	8	12	5	10
East North Central	8	12	6	11	4	10
West North Central	6	6	4	6	3	4
South	19	20	15	18	10	10
South Atlantic	12	13	10	12	6	8
East South Central	6	11	6	9	5	4
West South Central	14	11	10	10	6	4
West	16	17	13	14	6	8
Mountain	10	11	8	10	3	7
Pacific	13	14	11	10	5	5
COUNTY TYPE						
Large Metro	22	19	17	17	10	9
Small Metro	17	17	14	15	9	9
250K - 1 Mil. Pop.	14	14	12	11	8	7
<250K Pop.	9	10	8	9	4	5
Nonmetro	11	16	9	15	3	11
Urbanized	6	7	5	5	2	3
Less Urbanized	8	11	7	10	3	5
Completely Rural	4	9	3	9	0	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.82D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.13	0.13	0.10	0.12	0.06	0.07
GEOGRAPHIC DIVISION						
Northeast	0.26	0.23	0.26	0.19	0.13	0.09
New England	0.46	0.49	0.44	0.38	0.15	0.22
Middle Atlantic	0.31	0.26	0.31	0.22	0.17	0.09
Midwest	0.18	0.24	0.14	0.23	0.10	0.19
East North Central	0.22	0.31	0.17	0.29	0.11	0.26
West North Central	0.33	0.35	0.26	0.33	0.19	0.21
South	0.23	0.24	0.18	0.22	0.12	0.12
South Atlantic	0.29	0.32	0.24	0.30	0.15	0.20
East South Central	0.45	0.78	0.39	0.65	0.32	0.30
West South Central	0.49	0.38	0.34	0.34	0.22	0.14
West	0.31	0.32	0.25	0.26	0.12	0.15
Mountain	0.59	0.66	0.52	0.62	0.19	0.41
Pacific	0.36	0.36	0.28	0.26	0.14	0.13
COUNTY TYPE						
Large Metro	0.19	0.17	0.15	0.15	0.09	0.08
Small Metro	0.24	0.24	0.21	0.20	0.12	0.12
250K - 1 Mil. Pop.	0.27	0.25	0.23	0.21	0.14	0.13
<250K Pop.	0.55	0.57	0.47	0.51	0.23	0.27
Nonmetro	0.23	0.34	0.20	0.31	0.07	0.23
Urbanized	0.42	0.50	0.35	0.39	0.11	0.23
Less Urbanized	0.30	0.38	0.27	0.36	0.11	0.16
Completely Rural	0.72	1.58	0.50	1.55	0.07	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.82P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 12 to 17, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.9117		0.7337		0.7400	
GEOGRAPHIC DIVISION						
Northeast	0.7221		0.3766		0.2580	
New England	0.6315		0.9634		0.9260	
Middle Atlantic	0.4833		0.2978		0.2223	
Midwest	0.4175		0.1773		0.4463	
East North Central	0.2707		0.2620		0.5647	
West North Central	0.7932		0.4510		0.5697	
South	0.6182		0.8138		0.5799	
South Atlantic	0.6608		0.5769		0.8535	
East South Central	0.3082		0.4088		0.7281	
West South Central	0.2903		0.6903		0.2956	
West	0.9509		0.5934		0.1920	
Mountain	0.9021		0.8570		0.1587	
Pacific	0.8671		0.5731		0.6930	
COUNTY TYPE						
Large Metro	0.3042		0.6295		0.3322	
Small Metro	0.9401		0.7036		0.9650	
250K - 1 Mil. Pop.	0.7100		0.6878		0.8352	
<250K Pop.	0.7591		0.8905		0.6668	
Nonmetro	0.1584		0.0634		0.0757	
Urbanized	0.3789		0.2758		0.0693	
Less Urbanized	0.1917		0.1556		0.5198	
Completely Rural	0.8353		0.4711		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.83C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	93	83	60	52	35	32
GEOGRAPHIC DIVISION						
Northeast	36	34	27	21	11	11
New England	21	18	15	10	6	8
Middle Atlantic	30	29	22	19	9	8
Midwest	44	38	29	22	15	11
East North Central	40	33	24	18	12	10
West North Central	19	19	17	12	10	6
South	57	43	36	29	22	18
South Atlantic	41	25	26	19	15	11
East South Central	20	21	11	12	6	6
West South Central	34	28	22	18	15	12
West	46	49	27	30	19	20
Mountain	27	22	15	15	10	9
Pacific	38	44	23	26	17	18
COUNTY TYPE						
Large Metro	68	63	46	38	26	20
Small Metro	55	43	32	29	19	20
250K - 1 Mil. Pop.	42	36	25	24	16	17
<250K Pop.	35	23	19	16	11	10
Nonmetro	30	28	20	20	12	9
Urbanized	20	17	13	12	7	5
Less Urbanized	22	23	15	15	10	8
Completely Rural	10	7	9	4	2	1

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.83D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.33	0.29	0.21	0.18	0.12	0.11
GEOGRAPHIC DIVISION						
Northeast	0.72	0.66	0.54	0.42	0.22	0.22
New England	1.59	1.35	1.20	0.80	0.49	0.60
Middle Atlantic	0.80	0.76	0.59	0.49	0.25	0.22
Midwest	0.65	0.55	0.43	0.32	0.22	0.17
East North Central	0.84	0.69	0.50	0.38	0.25	0.20
West North Central	0.94	0.88	0.82	0.59	0.47	0.31
South	0.57	0.42	0.35	0.28	0.21	0.17
South Atlantic	0.85	0.51	0.53	0.39	0.31	0.23
East South Central	1.12	1.13	0.62	0.65	0.31	0.34
West South Central	0.99	0.80	0.64	0.51	0.42	0.35
West	0.71	0.73	0.42	0.45	0.30	0.30
Mountain	1.36	1.07	0.76	0.74	0.52	0.45
Pacific	0.83	0.94	0.50	0.56	0.36	0.38
COUNTY TYPE						
Large Metro	0.49	0.45	0.33	0.27	0.19	0.15
Small Metro	0.60	0.46	0.35	0.31	0.21	0.21
250K - 1 Mil. Pop.	0.65	0.55	0.40	0.37	0.24	0.26
<250K Pop.	1.29	0.82	0.71	0.56	0.41	0.36
Nonmetro	0.57	0.51	0.38	0.36	0.22	0.17
Urbanized	1.02	0.89	0.65	0.62	0.36	0.28
Less Urbanized	0.76	0.71	0.52	0.47	0.36	0.23
Completely Rural	1.89	1.49	1.69	0.75	0.41	0.22

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.83P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 to 25, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0204		0.0022		0.0552	
GEOGRAPHIC DIVISION						
Northeast	0.7470		0.9812		0.6508	
New England	0.9814		0.6930		0.4006	
Middle Atlantic	0.7073		0.8365		1.0000	
Midwest	0.3787		0.0469		0.1516	
East North Central	0.8472		0.0934		0.3063	
West North Central	0.0978		0.2898		0.3014	
South	0.0199		0.0166		0.0287	
South Atlantic	0.2183		0.0907		0.0431	
East South Central	0.9007		0.9275		0.5297	
West South Central	0.0108		0.0494		0.1360	
West	0.1980		0.1793		0.8493	
Mountain	0.9520		0.9887		0.7109	
Pacific	0.1183		0.1262		1.0000	
COUNTY TYPE						
Large Metro	0.5039		0.1066		0.1039	
Small Metro	0.0028		0.0083		0.5313	
250K - 1 Mil. Pop.	0.0017		0.0045		0.6412	
<250K Pop.	0.3917		0.5608		0.6533	
Nonmetro	0.7874		0.3612		0.2383	
Urbanized	0.9773		0.7489		0.4648	
Less Urbanized	0.7760		0.6375		0.5268	
Completely Rural	0.8560		0.2483		0.1312	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.84C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	580	505	170	135	123	102
GEOGRAPHIC DIVISION						
Northeast	278	205	77	60	72	35
New England	153	91	55	39	54	19
Middle Atlantic	235	186	53	47	48	30
Midwest	205	181	53	50	39	33
East North Central	166	158	43	45	26	27
West North Central	120	89	32	22	29	19
South	353	294	112	63	73	39
South Atlantic	293	222	78	53	53	32
East South Central	115	122	63	19	38	13
West South Central	157	146	51	29	32	19
West	309	291	89	90	56	81
Mountain	98	99	22	33	20	15
Pacific	294	273	87	84	52	80
COUNTY TYPE						
Large Metro	467	397	124	109	97	91
Small Metro	305	253	108	76	75	41
250K - 1 Mil. Pop.	280	222	104	72	74	41
<250K Pop.	120	123	29	25	11	9
Nonmetro	173	184	46	35	30	19
Urbanized	89	117	33	16	21	8
Less Urbanized	131	126	30	25	21	16
Completely Rural	60	60	10	16	1	6

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.84D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.34	0.30	0.10	0.08	0.07	0.06
GEOGRAPHIC DIVISION						
Northeast	0.83	0.61	0.23	0.18	0.21	0.11
New England	1.75	1.04	0.63	0.44	0.61	0.22
Middle Atlantic	0.95	0.75	0.21	0.19	0.19	0.12
Midwest	0.52	0.46	0.14	0.13	0.10	0.08
East North Central	0.60	0.57	0.16	0.16	0.09	0.10
West North Central	1.03	0.76	0.27	0.18	0.25	0.16
South	0.59	0.48	0.19	0.10	0.12	0.06
South Atlantic	0.93	0.69	0.25	0.17	0.17	0.10
East South Central	1.10	1.15	0.60	0.18	0.36	0.12
West South Central	0.87	0.80	0.29	0.16	0.18	0.10
West	0.84	0.79	0.24	0.24	0.15	0.22
Mountain	0.94	0.93	0.21	0.31	0.19	0.14
Pacific	1.12	1.04	0.33	0.32	0.20	0.30
COUNTY TYPE						
Large Metro	0.55	0.47	0.15	0.13	0.11	0.11
Small Metro	0.60	0.49	0.21	0.15	0.15	0.08
250K - 1 Mil. Pop.	0.73	0.57	0.27	0.18	0.19	0.10
<250K Pop.	0.98	0.95	0.24	0.19	0.09	0.07
Nonmetro	0.51	0.52	0.14	0.10	0.09	0.05
Urbanized	0.91	1.16	0.34	0.16	0.22	0.08
Less Urbanized	0.67	0.60	0.15	0.12	0.11	0.08
Completely Rural	1.29	1.44	0.22	0.39	0.02	0.14

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.84P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 26 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.5398		0.3156		0.1003	
GEOGRAPHIC DIVISION						
Northeast	0.1055		0.9058		0.0966	
New England	0.9448		0.5862		0.3235	
Middle Atlantic	0.0749		0.7127		0.1731	
Midwest	0.9238		0.7800		0.4736	
East North Central	0.8592		0.9670		0.7641	
West North Central	0.9093		0.5937		0.4804	
South	0.1967		0.0223		0.0467	
South Atlantic	0.6655		0.2523		0.0573	
East South Central	0.2422		0.0537		0.4334	
West South Central	0.2273		0.2685		0.7059	
West	0.0600		0.5497		0.5098	
Mountain	0.0589		0.0764		1.0000	
Pacific	0.2049		0.9029		0.4976	
COUNTY TYPE						
Large Metro	0.4690		0.5592		0.4265	
Small Metro	0.2804		0.4775		0.0912	
250K - 1 Mil. Pop.	0.2090		0.5124		0.0862	
<250K Pop.	0.8838		0.8112		0.8947	
Nonmetro	0.0685		0.6607		0.7872	
Urbanized	0.0113		0.2793		0.5066	
Less Urbanized	0.5000		0.8208		0.9822	
Completely Rural	0.6552		0.2362		0.3716	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.85C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Cocaine Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	594	515	187	146	130	110
GEOGRAPHIC DIVISION						
Northeast	285	212	85	65	72	39
New England	157	94	62	37	54	23
Middle Atlantic	241	193	59	53	49	31
Midwest	213	191	62	56	44	35
East North Central	174	166	50	49	28	29
West North Central	124	94	38	28	33	20
South	360	296	123	70	78	44
South Atlantic	298	226	87	58	57	35
East South Central	118	116	63	22	38	15
West South Central	164	151	59	33	38	23
West	315	295	94	94	60	86
Mountain	102	104	27	34	23	17
Pacific	299	276	90	88	55	85
COUNTY TYPE						
Large Metro	478	404	137	112	101	95
Small Metro	314	257	114	86	78	50
250K - 1 Mil. Pop.	286	223	107	78	76	47
<250K Pop.	125	126	39	31	16	14
Nonmetro	178	190	56	41	36	21
Urbanized	93	119	40	20	27	10
Less Urbanized	136	131	36	30	25	18
Completely Rural	59	62	14	17	2	6

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.85D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.30	0.26	0.09	0.07	0.07	0.06
GEOGRAPHIC DIVISION						
Northeast	0.74	0.55	0.22	0.17	0.19	0.10
New England	1.56	0.93	0.62	0.36	0.53	0.23
Middle Atlantic	0.84	0.68	0.21	0.19	0.17	0.11
Midwest	0.46	0.41	0.14	0.12	0.10	0.08
East North Central	0.54	0.51	0.15	0.15	0.09	0.09
West North Central	0.90	0.68	0.28	0.20	0.24	0.14
South	0.51	0.42	0.17	0.10	0.11	0.06
South Atlantic	0.82	0.61	0.24	0.16	0.16	0.09
East South Central	0.96	0.93	0.51	0.18	0.31	0.12
West South Central	0.77	0.70	0.28	0.15	0.18	0.10
West	0.73	0.67	0.22	0.22	0.14	0.20
Mountain	0.83	0.82	0.22	0.27	0.18	0.14
Pacific	0.97	0.89	0.29	0.28	0.18	0.27
COUNTY TYPE						
Large Metro	0.48	0.41	0.14	0.11	0.10	0.10
Small Metro	0.53	0.42	0.19	0.14	0.13	0.08
250K - 1 Mil. Pop.	0.64	0.49	0.24	0.17	0.17	0.10
<250K Pop.	0.83	0.79	0.26	0.20	0.11	0.09
Nonmetro	0.45	0.47	0.14	0.10	0.09	0.05
Urbanized	0.80	0.99	0.34	0.17	0.23	0.09
Less Urbanized	0.60	0.54	0.16	0.13	0.11	0.07
Completely Rural	1.14	1.34	0.27	0.37	0.04	0.13

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.85P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Cocaine Among Persons Aged 18 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3246		0.0559		0.0455	
GEOGRAPHIC DIVISION						
Northeast	0.1215		0.8911		0.1135	
New England	0.9438		0.5220		0.3715	
Middle Atlantic	0.0878		0.6913		0.1802	
Midwest	0.7814		0.2802		0.2629	
East North Central	0.8331		0.4787		0.5091	
West North Central	0.8524		0.3917		0.3534	
South	0.1006		0.0050		0.0180	
South Atlantic	0.5592		0.1296		0.0281	
East South Central	0.2505		0.0648		0.5477	
West South Central	0.0776		0.0771		0.3564	
West	0.1106		0.8425		0.5665	
Mountain	0.0753		0.1169		0.8556	
Pacific	0.3266		0.7594		0.5242	
COUNTY TYPE						
Large Metro	0.4195		0.2820		0.2672	
Small Metro	0.1020		0.1480		0.0924	
250K - 1 Mil. Pop.	0.0764		0.1827		0.0936	
<250K Pop.	0.9372		0.5984		0.8096	
Nonmetro	0.0884		0.4843		0.5216	
Urbanized	0.0141		0.2886		0.4452	
Less Urbanized	0.5373		0.7515		0.8007	
Completely Rural	0.6433		0.5807		0.7643	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.86C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	557	487	101	131	59	64
GEOGRAPHIC DIVISION						
Northeast	250	182	41	56	20	29
New England	150	87	22	39	12	25
Middle Atlantic	205	160	34	40	16	15
Midwest	198	200	46	47	23	18
East North Central	157	168	39	41	20	15
West North Central	120	108	23	24	10	9
South	306	270	66	64	40	33
South Atlantic	242	207	55	55	38	27
East South Central	120	97	25	20	6	13
West South Central	139	141	27	25	12	12
West	336	289	46	88	31	43
Mountain	126	106	26	30	14	17
Pacific	312	269	38	83	28	40
COUNTY TYPE						
Large Metro	437	391	77	109	46	56
Small Metro	300	266	60	63	35	27
250K - 1 Mil. Pop.	262	212	48	50	28	20
<250K Pop.	140	147	37	39	21	17
Nonmetro	171	189	39	36	14	17
Urbanized	95	117	19	20	9	11
Less Urbanized	135	122	32	26	10	11
Completely Rural	61	69	16	12	5	4

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.86D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.25	0.22	0.05	0.06	0.03	0.03
GEOGRAPHIC DIVISION						
Northeast	0.59	0.42	0.10	0.13	0.05	0.07
New England	1.34	0.78	0.20	0.35	0.11	0.23
Middle Atlantic	0.65	0.51	0.11	0.13	0.05	0.05
Midwest	0.38	0.39	0.09	0.09	0.04	0.03
East North Central	0.44	0.46	0.11	0.11	0.06	0.04
West North Central	0.77	0.70	0.15	0.15	0.06	0.06
South	0.39	0.34	0.08	0.08	0.05	0.04
South Atlantic	0.60	0.50	0.14	0.13	0.09	0.07
East South Central	0.88	0.70	0.18	0.14	0.05	0.10
West South Central	0.57	0.57	0.11	0.10	0.05	0.05
West	0.69	0.59	0.10	0.18	0.06	0.09
Mountain	0.90	0.74	0.19	0.21	0.10	0.12
Pacific	0.90	0.77	0.11	0.24	0.08	0.11
COUNTY TYPE						
Large Metro	0.40	0.36	0.07	0.10	0.04	0.05
Small Metro	0.45	0.39	0.09	0.09	0.05	0.04
250K - 1 Mil. Pop.	0.52	0.42	0.10	0.10	0.06	0.04
<250K Pop.	0.84	0.83	0.22	0.22	0.13	0.10
Nonmetro	0.39	0.42	0.09	0.08	0.03	0.04
Urbanized	0.72	0.88	0.15	0.15	0.07	0.08
Less Urbanized	0.54	0.45	0.13	0.10	0.04	0.04
Completely Rural	1.06	1.31	0.27	0.22	0.09	0.09

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.86P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2561		0.1092		0.6325	
GEOGRAPHIC DIVISION						
Northeast	0.3421		0.0387		0.4301	
New England	0.0787		0.0408		0.2555	
Middle Atlantic	0.9687		0.3148		0.9262	
Midwest	0.6090		0.5909		0.1351	
East North Central	0.4577		0.7379		0.0683	
West North Central	0.8629		0.6393		0.8436	
South	0.8197		0.4197		0.5012	
South Atlantic	0.9943		0.4974		0.8360	
East South Central	0.8173		0.7728		0.0688	
West South Central	0.5147		0.7436		0.9463	
West	0.2382		0.3502		0.6458	
Mountain	0.3491		0.8404		0.2617	
Pacific	0.3859		0.3583		0.9787	
COUNTY TYPE						
Large Metro	0.5914		0.0508		0.3653	
Small Metro	0.4524		0.5339		0.6047	
250K - 1 Mil. Pop.	0.9772		0.3907		0.2949	
<250K Pop.	0.1403		0.8022		0.5884	
Nonmetro	0.2975		0.3248		0.8069	
Urbanized	0.0312		0.2701		0.5442	
Less Urbanized	0.8836		0.1038		0.8934	
Completely Rural	0.9255		0.5473		0.9604	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.87C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	45	43	36	36	20	22
GEOGRAPHIC DIVISION						
Northeast	18	19	16	16	8	7
New England	10	13	8	11	6	6
Middle Atlantic	15	14	14	12	6	5
Midwest	18	20	15	17	10	8
East North Central	14	17	13	14	8	8
West North Central	10	10	8	10	5	3
South	26	28	20	23	10	16
South Atlantic	18	21	15	19	7	14
East South Central	10	11	8	8	4	7
West South Central	16	15	11	10	6	5
West	27	17	20	15	11	8
Mountain	14	9	10	8	6	4
Pacific	23	14	18	13	9	7
COUNTY TYPE						
Large Metro	32	28	27	25	15	15
Small Metro	27	25	20	21	11	11
250K - 1 Mil. Pop.	24	20	17	16	10	8
<250K Pop.	12	16	9	13	5	7
Nonmetro	16	19	14	17	6	10
Urbanized	9	10	7	8	3	4
Less Urbanized	12	13	9	11	5	9
Completely Rural	6	9	6	8	1	4

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.87D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.19	0.18	0.16	0.16	0.08	0.09
GEOGRAPHIC DIVISION						
Northeast	0.44	0.45	0.38	0.38	0.20	0.17
New England	0.96	1.21	0.70	0.97	0.56	0.52
Middle Atlantic	0.49	0.44	0.44	0.40	0.19	0.15
Midwest	0.32	0.37	0.28	0.31	0.18	0.15
East North Central	0.37	0.46	0.33	0.38	0.22	0.21
West North Central	0.60	0.60	0.48	0.56	0.31	0.19
South	0.31	0.34	0.24	0.28	0.12	0.20
South Atlantic	0.44	0.51	0.36	0.47	0.17	0.34
East South Central	0.70	0.81	0.56	0.60	0.31	0.48
West South Central	0.55	0.53	0.41	0.37	0.22	0.18
West	0.50	0.31	0.38	0.28	0.20	0.16
Mountain	0.87	0.57	0.63	0.51	0.35	0.26
Pacific	0.61	0.37	0.47	0.34	0.24	0.19
COUNTY TYPE						
Large Metro	0.27	0.25	0.23	0.22	0.13	0.13
Small Metro	0.39	0.35	0.28	0.29	0.16	0.16
250K - 1 Mil. Pop.	0.46	0.37	0.32	0.30	0.19	0.15
<250K Pop.	0.71	0.86	0.55	0.70	0.29	0.41
Nonmetro	0.35	0.39	0.29	0.35	0.12	0.21
Urbanized	0.59	0.75	0.47	0.63	0.23	0.28
Less Urbanized	0.45	0.45	0.36	0.40	0.17	0.30
Completely Rural	1.05	1.53	1.06	1.40	0.10	0.67

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.87P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 12 to 17, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.7335		0.8101		0.6370	
GEOGRAPHIC DIVISION						
Northeast	0.8349		0.5032		0.3777	
New England	0.6386		0.7267		0.7241	
Middle Atlantic	0.4391		0.2224		0.3501	
Midwest	0.2138		0.5531		0.7081	
East North Central	0.3728		0.9276		0.8826	
West North Central	0.3596		0.2467		0.6158	
South	0.4913		0.2768		0.1025	
South Atlantic	0.4435		0.1910		0.1346	
East South Central	0.3891		0.6532		0.2411	
West South Central	0.6826		0.8383		0.8196	
West	0.3435		0.4021		0.8074	
Mountain	0.6810		0.7960		0.9386	
Pacific	0.3862		0.2332		0.8122	
COUNTY TYPE						
Large Metro	0.8660		0.9703		0.9060	
Small Metro	0.9472		0.7406		0.9710	
250K - 1 Mil. Pop.	0.3456		0.9091		0.5296	
<250K Pop.	0.1458		0.4822		0.2759	
Nonmetro	0.6607		0.8693		0.3353	
Urbanized	0.0664		0.2785		0.6250	
Less Urbanized	0.3288		0.5298		0.6086	
Completely Rural	0.5835		0.8055		0.2573	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.88C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	118	118	67	71	36	34
GEOGRAPHIC DIVISION						
Northeast	53	50	30	34	16	13
New England	27	26	19	18	9	7
Middle Atlantic	46	42	23	29	13	11
Midwest	54	51	33	29	17	14
East North Central	45	39	28	25	15	12
West North Central	29	33	18	13	7	7
South	70	59	40	40	20	22
South Atlantic	55	41	32	33	16	18
East South Central	19	26	12	13	5	8
West South Central	40	34	20	19	11	9
West	56	73	31	38	18	17
Mountain	28	25	16	19	10	12
Pacific	48	68	27	33	15	13
COUNTY TYPE						
Large Metro	87	94	50	52	27	25
Small Metro	71	60	42	41	21	19
250K - 1 Mil. Pop.	51	47	34	31	17	14
<250K Pop.	50	39	23	29	11	12
Nonmetro	42	38	24	24	10	12
Urbanized	24	19	14	15	7	10
Less Urbanized	33	30	19	18	6	7
Completely Rural	12	11	11	7	5	1

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.88D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.42	0.41	0.24	0.25	0.13	0.12
GEOGRAPHIC DIVISION						
Northeast	1.06	0.97	0.59	0.67	0.32	0.26
New England	2.12	1.97	1.49	1.36	0.72	0.50
Middle Atlantic	1.22	1.12	0.60	0.77	0.36	0.30
Midwest	0.79	0.74	0.49	0.41	0.25	0.20
East North Central	0.95	0.82	0.58	0.53	0.32	0.24
West North Central	1.42	1.56	0.88	0.63	0.35	0.34
South	0.70	0.58	0.39	0.39	0.20	0.21
South Atlantic	1.13	0.82	0.66	0.67	0.34	0.36
East South Central	1.04	1.42	0.64	0.69	0.27	0.41
West South Central	1.14	0.98	0.59	0.53	0.30	0.27
West	0.86	1.10	0.48	0.57	0.27	0.26
Mountain	1.45	1.24	0.80	0.94	0.53	0.58
Pacific	1.05	1.47	0.59	0.71	0.32	0.27
COUNTY TYPE						
Large Metro	0.62	0.68	0.36	0.38	0.19	0.18
Small Metro	0.78	0.63	0.46	0.44	0.23	0.20
250K - 1 Mil. Pop.	0.79	0.71	0.52	0.46	0.27	0.22
<250K Pop.	1.82	1.35	0.86	1.01	0.40	0.42
Nonmetro	0.78	0.67	0.44	0.44	0.19	0.22
Urbanized	1.24	1.00	0.74	0.80	0.36	0.52
Less Urbanized	1.14	0.94	0.64	0.56	0.19	0.22
Completely Rural	2.29	2.33	2.09	1.40	0.98	0.30

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.88P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 to 25, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.9899		0.9661		0.8172	
GEOGRAPHIC DIVISION						
Northeast	0.2574		0.3022		0.8637	
New England	0.2820		0.6222		0.8968	
Middle Atlantic	0.5390		0.3715		0.8985	
Midwest	0.9033		0.1464		0.2474	
East North Central	0.5071		0.5663		0.1769	
West North Central	0.3007		0.0677		0.9440	
South	0.7219		0.5797		0.2855	
South Atlantic	0.2866		0.8070		0.2416	
East South Central	0.0161		0.3426		0.1990	
West South Central	0.3856		0.9285		0.5943	
West	0.7318		0.8118		0.5575	
Mountain	0.2950		0.9175		0.4902	
Pacific	0.4023		0.8258		0.1519	
COUNTY TYPE						
Large Metro	0.9611		0.8935		0.5397	
Small Metro	0.9932		0.7333		0.8991	
250K - 1 Mil. Pop.	0.9779		0.8814		0.2961	
<250K Pop.	0.9574		0.6957		0.0735	
Nonmetro	0.9778		0.3049		0.6016	
Urbanized	0.4045		0.0371		0.2787	
Less Urbanized	0.4526		0.0294		0.7892	
Completely Rural	0.5822		0.2356		0.3171	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.89C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	525	453	59	81	42	39
GEOGRAPHIC DIVISION						
Northeast	239	168	10	38	7	23
New England	146	76	3	34	3	22
Middle Atlantic	193	148	9	18	6	8
Midwest	182	179	27	25	15	8
East North Central	147	151	26	21	15	6
West North Central	108	97	8	14	*	5
South	286	246	43	33	33	10
South Atlantic	227	189	36	29	33	5
East South Central	115	91	21	9	*	6
West South Central	127	126	10	12	*	6
West	319	274	28	58	21	29
Mountain	110	99	14	15	5	9
Pacific	300	255	24	56	20	27
COUNTY TYPE						
Large Metro	412	365	43	75	33	38
Small Metro	283	240	32	29	25	9
250K - 1 Mil. Pop.	253	194	24	27	19	7
<250K Pop.	124	128	21	11	17	5
Nonmetro	162	181	24	16	8	2
Urbanized	89	112	7	6	4	2
Less Urbanized	126	118	22	15	7	*
Completely Rural	59	63	3	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.89D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.31	0.26	0.03	0.05	0.02	0.02
GEOGRAPHIC DIVISION						
Northeast	0.71	0.50	0.03	0.11	0.02	0.07
New England	1.66	0.87	0.04	0.38	0.04	0.25
Middle Atlantic	0.78	0.60	0.04	0.07	0.02	0.03
Midwest	0.46	0.45	0.07	0.06	0.04	0.02
East North Central	0.53	0.54	0.09	0.08	0.06	0.02
West North Central	0.92	0.82	0.07	0.12	*	0.04
South	0.48	0.40	0.07	0.05	0.05	0.02
South Atlantic	0.72	0.59	0.12	0.09	0.10	0.01
East South Central	1.09	0.86	0.20	0.08	*	0.06
West South Central	0.71	0.69	0.06	0.07	*	0.03
West	0.87	0.74	0.08	0.16	0.06	0.08
Mountain	1.05	0.93	0.13	0.14	0.05	0.08
Pacific	1.15	0.97	0.09	0.21	0.08	0.10
COUNTY TYPE						
Large Metro	0.48	0.43	0.05	0.09	0.04	0.05
Small Metro	0.56	0.46	0.06	0.06	0.05	0.02
250K - 1 Mil. Pop.	0.66	0.50	0.06	0.07	0.05	0.02
<250K Pop.	1.02	0.99	0.17	0.08	0.14	0.04
Nonmetro	0.48	0.51	0.07	0.05	0.02	0.01
Urbanized	0.91	1.11	0.07	0.06	0.04	0.02
Less Urbanized	0.64	0.57	0.11	0.07	0.03	*
Completely Rural	1.26	1.49	0.06	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.89P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 26 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2595		0.0157		0.5384	
GEOGRAPHIC DIVISION						
Northeast	0.4614		0.0108		0.1097	
New England	0.1197		0.0614		0.1456	
Middle Atlantic	0.8958		0.0687		0.5268	
Midwest	0.6625		0.7670		0.4176	
East North Central	0.6229		0.9424		0.2516	
West North Central	0.9695		0.6308		*	
South	0.8046		0.9438		0.4059	
South Atlantic	0.9013		0.9250		0.1919	
East South Central	0.6255		0.5709		*	
West South Central	0.6639		0.4405		*	
West	0.1691		0.0452		0.2317	
Mountain	0.4320		0.7578		0.1410	
Pacific	0.2452		0.0452		0.3667	
COUNTY TYPE						
Large Metro	0.6044		0.0037		0.0975	
Small Metro	0.4350		0.6932		0.2943	
250K - 1 Mil. Pop.	0.9554		0.1699		0.6794	
<250K Pop.	0.1464		0.1737		0.2642	
Nonmetro	0.2994		0.4603		0.0766	
Urbanized	0.0608		0.1977		0.4181	
Less Urbanized	0.9377		0.7868		*	
Completely Rural	0.7666		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.90C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Hallucinogens Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	555	482	94	122	55	57
GEOGRAPHIC DIVISION						
Northeast	251	182	36	54	18	28
New England	152	82	23	38	10	24
Middle Atlantic	204	162	28	38	15	14
Midwest	198	197	44	41	21	16
East North Central	157	165	38	36	19	13
West North Central	120	107	22	20	8	9
South	305	266	63	57	39	24
South Atlantic	242	207	53	50	37	20
East South Central	120	92	23	16	5	10
West South Central	137	136	24	23	11	10
West	333	286	40	85	28	41
Mountain	120	104	24	28	13	17
Pacific	310	266	32	80	24	37
COUNTY TYPE						
Large Metro	435	390	70	104	42	51
Small Metro	298	258	57	58	33	22
250K - 1 Mil. Pop.	261	206	44	46	26	17
<250K Pop.	136	143	36	36	20	14
Nonmetro	170	187	35	31	13	13
Urbanized	93	115	18	18	8	10
Less Urbanized	133	123	29	24	9	7
Completely Rural	60	68	12	7	5	1

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.90D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.28	0.24	0.05	0.06	0.03	0.03
GEOGRAPHIC DIVISION						
Northeast	0.65	0.47	0.09	0.14	0.05	0.07
New England	1.51	0.81	0.23	0.38	0.10	0.24
Middle Atlantic	0.72	0.57	0.10	0.13	0.05	0.05
Midwest	0.43	0.42	0.10	0.09	0.04	0.03
East North Central	0.49	0.51	0.12	0.11	0.06	0.04
West North Central	0.88	0.77	0.16	0.15	0.06	0.06
South	0.43	0.37	0.09	0.08	0.06	0.03
South Atlantic	0.66	0.56	0.15	0.13	0.10	0.05
East South Central	0.98	0.74	0.19	0.13	0.04	0.08
West South Central	0.64	0.63	0.11	0.10	0.05	0.04
West	0.77	0.65	0.09	0.19	0.06	0.09
Mountain	0.97	0.82	0.19	0.22	0.11	0.13
Pacific	1.01	0.86	0.10	0.26	0.08	0.12
COUNTY TYPE						
Large Metro	0.44	0.40	0.07	0.11	0.04	0.05
Small Metro	0.50	0.42	0.10	0.09	0.06	0.04
250K - 1 Mil. Pop.	0.58	0.45	0.10	0.10	0.06	0.04
<250K Pop.	0.91	0.90	0.24	0.23	0.14	0.09
Nonmetro	0.43	0.46	0.09	0.08	0.03	0.03
Urbanized	0.80	0.96	0.15	0.15	0.07	0.09
Less Urbanized	0.59	0.51	0.13	0.10	0.04	0.03
Completely Rural	1.16	1.45	0.23	0.14	0.10	0.03

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.90P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Hallucinogens Among Persons Aged 18 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2707		0.0999		0.7252	
GEOGRAPHIC DIVISION						
Northeast	0.3316		0.0135		0.2307	
New England	0.0784		0.0358		0.1417	
Middle Atlantic	0.9805		0.1349		0.8218	
Midwest	0.6953		0.4282		0.1347	
East North Central	0.5125		0.7466		0.0518	
West North Central	0.7937		0.3298		0.6039	
South	0.7671		0.6496		0.9330	
South Atlantic	0.9442		0.8060		0.7479	
East South Central	0.8940		0.8972		0.1144	
West South Central	0.5333		0.6027		0.9548	
West	0.2098		0.2334		0.5665	
Mountain	0.3187		0.8999		0.2196	
Pacific	0.3527		0.2209		0.9227	
COUNTY TYPE						
Large Metro	0.6060		0.0366		0.3370	
Small Metro	0.4488		0.5961		0.5628	
250K - 1 Mil. Pop.	0.9449		0.3374		0.3757	
<250K Pop.	0.1776		0.5405		0.8264	
Nonmetro	0.3142		0.2330		0.7345	
Urbanized	0.0493		0.3439		0.5843	
Less Urbanized	0.9627		0.1181		0.5330	
Completely Rural	0.8739		0.1777		0.3063	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.91C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	443	381	88	90	52	67
GEOGRAPHIC DIVISION						
Northeast	207	151	32	43	20	36
New England	117	92	15	23	5	19
Middle Atlantic	173	120	28	36	19	30
Midwest	181	156	39	35	23	21
East North Central	152	129	37	30	23	20
West North Central	98	91	11	18	4	5
South	255	225	49	50	30	37
South Atlantic	216	181	39	38	21	30
East South Central	83	90	12	24	5	19
West South Central	107	97	27	21	22	11
West	232	219	53	50	29	38
Mountain	81	82	26	15	22	5
Pacific	218	203	47	48	19	37
COUNTY TYPE						
Large Metro	357	282	69	74	47	58
Small Metro	231	207	47	45	16	32
250K - 1 Mil. Pop.	200	172	35	38	15	24
<250K Pop.	113	114	31	26	7	21
Nonmetro	142	141	24	25	13	11
Urbanized	77	78	13	15	5	7
Less Urbanized	109	101	17	20	8	8
Completely Rural	35	52	12	7	9	3

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.91D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.20	0.17	0.04	0.04	0.02	0.03
GEOGRAPHIC DIVISION						
Northeast	0.48	0.35	0.08	0.10	0.05	0.08
New England	1.05	0.82	0.13	0.21	0.04	0.17
Middle Atlantic	0.55	0.38	0.09	0.11	0.06	0.09
Midwest	0.35	0.30	0.08	0.07	0.05	0.04
East North Central	0.42	0.35	0.10	0.08	0.06	0.06
West North Central	0.64	0.58	0.07	0.12	0.02	0.03
South	0.33	0.28	0.06	0.06	0.04	0.05
South Atlantic	0.53	0.44	0.10	0.09	0.05	0.07
East South Central	0.61	0.65	0.09	0.18	0.03	0.13
West South Central	0.44	0.39	0.11	0.09	0.09	0.05
West	0.48	0.45	0.11	0.10	0.06	0.08
Mountain	0.58	0.57	0.18	0.11	0.16	0.04
Pacific	0.63	0.58	0.14	0.14	0.05	0.11
COUNTY TYPE						
Large Metro	0.32	0.26	0.06	0.07	0.04	0.05
Small Metro	0.35	0.30	0.07	0.07	0.02	0.05
250K - 1 Mil. Pop.	0.40	0.34	0.07	0.07	0.03	0.05
<250K Pop.	0.68	0.64	0.19	0.15	0.04	0.12
Nonmetro	0.32	0.31	0.06	0.06	0.03	0.02
Urbanized	0.58	0.59	0.10	0.11	0.04	0.06
Less Urbanized	0.43	0.37	0.07	0.07	0.03	0.03
Completely Rural	0.61	0.99	0.21	0.13	0.16	0.05

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.91P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2871		0.7619		0.5823	
GEOGRAPHIC DIVISION						
Northeast	0.9351		0.3924		0.1486	
New England	0.6307		0.2883		0.1760	
Middle Atlantic	0.6651		0.7017		0.3895	
Midwest	0.0415		0.5588		0.7452	
East North Central	0.0658		0.4822		0.8389	
West North Central	0.3446		0.8966		0.4949	
South	0.9236		0.9561		0.6472	
South Atlantic	0.5814		0.8947		0.8596	
East South Central	0.4352		0.2401		0.1962	
West South Central	0.7702		0.5385		0.6943	
West	0.7019		0.4581		0.6420	
Mountain	0.4446		0.2158		0.0956	
Pacific	0.9560		0.7780		0.6674	
COUNTY TYPE						
Large Metro	0.2989		0.6612		0.4930	
Small Metro	0.8172		0.2578		0.6061	
250K - 1 Mil. Pop.	0.6412		0.6962		0.6577	
<250K Pop.	0.7553		0.1907		0.7823	
Nonmetro	0.8035		0.8606		0.2447	
Urbanized	0.2949		0.5956		0.4084	
Less Urbanized	0.8186		0.6501		0.2162	
Completely Rural	0.1476		0.7219		0.1503	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.92C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	54	49	37	32	18	17
GEOGRAPHIC DIVISION						
Northeast	24	21	19	14	8	6
New England	12	11	9	9	2	4
Middle Atlantic	21	18	16	11	8	5
Midwest	25	25	17	15	8	7
East North Central	21	21	15	12	7	6
West North Central	12	14	8	8	4	4
South	33	28	20	20	10	11
South Atlantic	24	17	13	12	6	7
East South Central	11	13	8	9	4	5
West South Central	20	18	13	13	7	7
West	24	24	19	15	10	8
Mountain	15	11	12	8	6	5
Pacific	19	21	15	13	8	6
COUNTY TYPE						
Large Metro	39	35	27	22	13	12
Small Metro	30	29	19	18	10	8
250K - 1 Mil. Pop.	26	24	17	16	9	7
<250K Pop.	13	16	9	9	4	4
Nonmetro	22	23	15	15	8	8
Urbanized	12	12	8	8	3	6
Less Urbanized	17	17	13	12	7	6
Completely Rural	7	11	6	5	2	3

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.92D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.23	0.21	0.16	0.14	0.08	0.07
GEOGRAPHIC DIVISION						
Northeast	0.58	0.50	0.45	0.34	0.19	0.15
New England	1.08	0.96	0.85	0.78	0.18	0.33
Middle Atlantic	0.69	0.58	0.53	0.37	0.25	0.17
Midwest	0.45	0.46	0.31	0.27	0.15	0.13
East North Central	0.56	0.55	0.39	0.32	0.20	0.16
West North Central	0.73	0.83	0.48	0.50	0.22	0.24
South	0.40	0.33	0.24	0.24	0.12	0.14
South Atlantic	0.59	0.42	0.32	0.31	0.15	0.18
East South Central	0.80	0.91	0.59	0.61	0.30	0.34
West South Central	0.71	0.62	0.45	0.46	0.24	0.26
West	0.46	0.45	0.36	0.28	0.19	0.14
Mountain	0.94	0.71	0.74	0.50	0.35	0.29
Pacific	0.52	0.56	0.40	0.34	0.22	0.16
COUNTY TYPE						
Large Metro	0.34	0.31	0.23	0.19	0.11	0.11
Small Metro	0.43	0.40	0.28	0.25	0.14	0.11
250K - 1 Mil. Pop.	0.50	0.45	0.32	0.29	0.17	0.13
<250K Pop.	0.79	0.87	0.53	0.50	0.23	0.24
Nonmetro	0.47	0.47	0.33	0.31	0.16	0.17
Urbanized	0.80	0.93	0.50	0.63	0.19	0.46
Less Urbanized	0.65	0.61	0.48	0.42	0.25	0.20
Completely Rural	1.29	1.90	1.02	0.81	0.43	0.42

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.92P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 12 to 17, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.4586		0.0783		0.3584	
GEOGRAPHIC DIVISION						
Northeast	0.8998		0.6097		0.4693	
New England	0.9736		0.8176		0.2522	
Middle Atlantic	0.8667		0.4684		0.1830	
Midwest	0.9463		0.1680		0.5589	
East North Central	0.6932		0.3634		0.5884	
West North Central	0.3809		0.2268		0.8194	
South	0.8237		0.7505		0.6865	
South Atlantic	0.6087		0.8517		0.6710	
East South Central	0.8939		0.9136		0.5201	
West South Central	0.9224		0.4123		0.8499	
West	0.0839		0.1744		0.2271	
Mountain	0.1092		0.4259		0.5679	
Pacific	0.2911		0.2703		0.2793	
COUNTY TYPE						
Large Metro	0.7358		0.7884		1.0000	
Small Metro	0.4662		0.0141		0.1383	
250K - 1 Mil. Pop.	0.4272		0.1072		0.2551	
<250K Pop.	0.9015		0.0276		0.3380	
Nonmetro	0.7453		0.4568		0.6676	
Urbanized	0.4520		0.6155		0.0796	
Less Urbanized	0.5746		0.2188		0.0731	
Completely Rural	0.2706		0.7455		0.7017	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.93C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	98	94	47	42	22	19
GEOGRAPHIC DIVISION						
Northeast	45	39	22	20	11	10
New England	30	19	10	10	4	5
Middle Atlantic	34	34	19	17	10	8
Midwest	47	38	24	18	9	8
East North Central	42	33	23	16	8	7
West North Central	20	19	7	8	3	2
South	55	50	26	21	12	9
South Atlantic	40	37	21	15	10	7
East South Central	21	17	9	10	2	2
West South Central	30	29	12	10	6	5
West	46	57	22	24	12	12
Mountain	21	21	9	8	1	2
Pacific	42	54	20	23	12	12
COUNTY TYPE						
Large Metro	71	68	34	31	16	15
Small Metro	54	52	27	23	12	11
250K - 1 Mil. Pop.	42	43	19	21	11	11
<250K Pop.	33	29	18	11	5	3
Nonmetro	37	33	14	15	9	6
Urbanized	18	18	9	11	4	4
Less Urbanized	28	26	8	10	3	3
Completely Rural	14	10	8	4	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.93D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.34	0.32	0.16	0.14	0.08	0.07
GEOGRAPHIC DIVISION						
Northeast	0.90	0.76	0.43	0.40	0.22	0.19
New England	2.32	1.46	0.79	0.79	0.30	0.37
Middle Atlantic	0.90	0.89	0.51	0.46	0.28	0.22
Midwest	0.70	0.55	0.36	0.25	0.13	0.11
East North Central	0.89	0.68	0.48	0.32	0.18	0.16
West North Central	0.99	0.92	0.36	0.39	0.13	0.11
South	0.54	0.49	0.25	0.20	0.12	0.09
South Atlantic	0.83	0.74	0.44	0.30	0.21	0.14
East South Central	1.17	0.94	0.47	0.52	0.12	0.13
West South Central	0.88	0.82	0.33	0.30	0.16	0.14
West	0.71	0.86	0.34	0.36	0.18	0.18
Mountain	1.06	1.03	0.46	0.42	0.07	0.10
Pacific	0.91	1.16	0.44	0.49	0.26	0.25
COUNTY TYPE						
Large Metro	0.51	0.49	0.25	0.22	0.11	0.11
Small Metro	0.59	0.55	0.29	0.24	0.13	0.12
250K - 1 Mil. Pop.	0.65	0.65	0.29	0.32	0.16	0.16
<250K Pop.	1.19	1.00	0.67	0.39	0.19	0.12
Nonmetro	0.70	0.60	0.27	0.27	0.18	0.10
Urbanized	0.95	0.98	0.46	0.58	0.19	0.23
Less Urbanized	0.96	0.80	0.28	0.31	0.10	0.11
Completely Rural	2.69	1.99	1.51	0.85	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.93P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 to 25, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0047		0.2930		0.4667	
GEOGRAPHIC DIVISION						
Northeast	0.2435		0.6949		1.0000	
New England	0.6554		0.7349		0.5084	
Middle Atlantic	0.0585		0.5234		0.7218	
Midwest	0.0382		0.2542		0.3418	
East North Central	0.2278		0.3461		0.4420	
West North Central	0.0203		0.4654		0.4928	
South	0.3939		0.3520		0.7189	
South Atlantic	0.6661		0.3573		0.6639	
East South Central	0.8361		0.9373		0.8651	
West South Central	0.2651		0.7123		0.9194	
West	0.0997		0.7099		0.7655	
Mountain	0.5970		0.8408		0.7047	
Pacific	0.0437		0.7438		0.7331	
COUNTY TYPE						
Large Metro	0.0158		0.3528		0.8692	
Small Metro	0.2149		0.3506		0.6293	
250K - 1 Mil. Pop.	0.2523		0.8457		0.9591	
<250K Pop.	0.5818		0.1839		0.1281	
Nonmetro	0.2957		0.6439		0.3475	
Urbanized	0.7851		0.2141		0.8396	
Less Urbanized	0.1205		0.9130		0.7724	
Completely Rural	0.9203		0.7094		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.94C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	416	355	61	68	40	59
GEOGRAPHIC DIVISION						
Northeast	191	144	18	33	15	32
New England	106	88	4	15	3	15
Middle Atlantic	160	113	18	29	14	28
Midwest	168	142	21	26	18	18
East North Central	140	113	21	22	18	18
West North Central	91	87	*	15	*	*
South	236	210	34	40	24	33
South Atlantic	198	170	29	31	17	28
East South Central	79	86	4	23	*	17
West South Central	101	87	18	12	16	6
West	226	204	41	36	23	32
Mountain	75	76	21	11	20	2
Pacific	213	189	36	34	10	32
COUNTY TYPE						
Large Metro	337	269	50	58	39	51
Small Metro	208	190	33	34	5	28
250K - 1 Mil. Pop.	182	160	23	27	5	20
<250K Pop.	99	102	23	21	*	20
Nonmetro	133	132	9	12	6	5
Urbanized	75	71	5	4	3	*
Less Urbanized	102	98	7	12	3	5
Completely Rural	30	45	3	*	3	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.94D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.25	0.21	0.04	0.04	0.02	0.03
GEOGRAPHIC DIVISION						
Northeast	0.57	0.43	0.05	0.10	0.04	0.10
New England	1.21	1.00	0.05	0.18	0.04	0.17
Middle Atlantic	0.65	0.46	0.07	0.12	0.06	0.11
Midwest	0.43	0.36	0.05	0.07	0.05	0.04
East North Central	0.51	0.41	0.08	0.08	0.07	0.06
West North Central	0.78	0.74	*	0.13	*	*
South	0.39	0.35	0.06	0.07	0.04	0.05
South Atlantic	0.63	0.53	0.09	0.10	0.05	0.09
East South Central	0.75	0.81	0.04	0.22	*	0.16
West South Central	0.56	0.48	0.10	0.07	0.09	0.03
West	0.62	0.55	0.11	0.10	0.06	0.09
Mountain	0.72	0.71	0.20	0.10	0.19	0.02
Pacific	0.81	0.72	0.14	0.13	0.04	0.12
COUNTY TYPE						
Large Metro	0.40	0.32	0.06	0.07	0.05	0.06
Small Metro	0.41	0.37	0.06	0.07	0.01	0.05
250K - 1 Mil. Pop.	0.48	0.41	0.06	0.07	0.01	0.05
<250K Pop.	0.81	0.79	0.19	0.16	*	0.16
Nonmetro	0.39	0.38	0.03	0.04	0.02	0.01
Urbanized	0.77	0.71	0.05	0.04	0.03	*
Less Urbanized	0.52	0.47	0.04	0.06	0.02	0.03
Completely Rural	0.64	1.07	0.07	*	0.07	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.94P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 26 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.7315		0.2160		0.1969	
GEOGRAPHIC DIVISION						
Northeast	0.8780		0.0874		0.0775	
New England	0.6896		0.2474		0.2474	
Middle Atlantic	0.9350		0.1782		0.1610	
Midwest	0.0981		0.3079		0.7596	
East North Central	0.0886		0.7215		0.7592	
West North Central	0.6323		*		*	
South	0.7872		0.3893		0.5988	
South Atlantic	0.5329		0.7508		0.8066	
East South Central	0.3980		0.1107		*	
West South Central	0.9758		1.0000		0.6497	
West	0.7759		0.5414		0.9893	
Mountain	0.4681		0.2651		0.0891	
Pacific	0.5212		0.8928		0.2795	
COUNTY TYPE						
Large Metro	0.6101		0.2271		0.4092	
Small Metro	0.8339		0.6331		0.1074	
250K - 1 Mil. Pop.	0.9287		0.5341		0.2294	
<250K Pop.	0.5688		0.9739		*	
Nonmetro	0.9641		0.9765		0.5142	
Urbanized	0.3650		0.7114		*	
Less Urbanized	0.8853		0.6580		0.8021	
Completely Rural	0.1812		*		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.95C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Inhalants Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	435	378	77	83	47	65
GEOGRAPHIC DIVISION						
Northeast	204	151	26	40	18	35
New England	115	91	12	22	5	19
Middle Atlantic	170	119	23	34	18	29
Midwest	178	155	35	33	22	19
East North Central	148	127	33	27	21	19
West North Central	97	90	8	18	3	2
South	249	221	44	45	29	34
South Atlantic	211	179	36	35	20	29
East South Central	80	87	10	24	2	17
West South Central	105	94	24	16	21	8
West	229	218	46	47	25	38
Mountain	80	83	23	14	20	3
Pacific	215	201	40	44	14	38
COUNTY TYPE						
Large Metro	351	282	61	70	44	57
Small Metro	224	204	43	41	13	30
250K - 1 Mil. Pop.	193	171	30	34	12	22
<250K Pop.	110	111	30	24	5	21
Nonmetro	139	139	18	20	11	8
Urbanized	76	77	11	12	5	4
Less Urbanized	106	101	11	15	5	6
Completely Rural	34	49	9	4	9	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.95D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.22	0.19	0.04	0.04	0.02	0.03
GEOGRAPHIC DIVISION						
Northeast	0.53	0.39	0.07	0.10	0.05	0.09
New England	1.15	0.90	0.12	0.22	0.05	0.18
Middle Atlantic	0.60	0.42	0.08	0.12	0.06	0.10
Midwest	0.39	0.33	0.07	0.07	0.05	0.04
East North Central	0.46	0.39	0.10	0.08	0.07	0.06
West North Central	0.71	0.65	0.06	0.13	0.02	0.02
South	0.35	0.31	0.06	0.06	0.04	0.05
South Atlantic	0.58	0.48	0.10	0.09	0.06	0.08
East South Central	0.65	0.70	0.08	0.19	0.02	0.14
West South Central	0.49	0.43	0.11	0.07	0.10	0.04
West	0.53	0.50	0.11	0.11	0.06	0.09
Mountain	0.65	0.65	0.19	0.11	0.16	0.02
Pacific	0.70	0.65	0.13	0.14	0.05	0.12
COUNTY TYPE						
Large Metro	0.36	0.29	0.06	0.07	0.04	0.06
Small Metro	0.38	0.33	0.07	0.07	0.02	0.05
250K - 1 Mil. Pop.	0.43	0.37	0.07	0.07	0.03	0.05
<250K Pop.	0.73	0.70	0.20	0.15	0.04	0.13
Nonmetro	0.35	0.34	0.04	0.05	0.03	0.02
Urbanized	0.65	0.64	0.09	0.10	0.04	0.04
Less Urbanized	0.47	0.42	0.05	0.06	0.02	0.03
Completely Rural	0.65	1.05	0.17	0.09	0.17	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.95P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Inhalants Among Persons Aged 18 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.3249		0.6377		0.3773	
GEOGRAPHIC DIVISION						
Northeast	0.9202		0.2139		0.0928	
New England	0.6265		0.2806		0.2435	
Middle Atlantic	0.6429		0.4158		0.2086	
Midwest	0.0382		0.9305		0.9062	
East North Central	0.0520		0.7386		0.9798	
West North Central	0.3883		0.3634		0.5098	
South	0.9415		0.8287		0.7268	
South Atlantic	0.6069		0.8295		0.9530	
East South Central	0.4244		0.1911		0.2255	
West South Central	0.7850		0.8605		0.7470	
West	0.8751		0.7708		0.8947	
Mountain	0.6002		0.3866		0.0999	
Pacific	0.9382		0.9458		0.4646	
COUNTY TYPE						
Large Metro	0.3114		0.5463		0.4715	
Small Metro	0.8962		0.8745		0.2700	
250K - 1 Mil. Pop.	0.7203		0.7089		0.3316	
<250K Pop.	0.7315		0.4841		0.5747	
Nonmetro	0.8422		0.6774		0.2518	
Urbanized	0.3767		0.4616		0.6648	
Less Urbanized	0.8645		0.7577		0.9469	
Completely Rural	0.2352		0.5020		*	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.96C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	645	582	296	259	209	178
GEOGRAPHIC DIVISION						
Northeast	292	218	125	122	102	68
New England	136	121	62	71	49	24
Middle Atlantic	258	184	110	99	90	63
Midwest	225	224	122	100	82	80
East North Central	187	201	97	89	70	76
West North Central	125	96	74	44	43	26
South	420	332	181	145	129	98
South Atlantic	325	246	127	108	101	68
East South Central	153	119	102	56	59	44
West South Central	217	188	81	79	54	55
West	323	354	155	146	100	106
Mountain	129	125	82	61	40	35
Pacific	296	331	132	132	92	100
COUNTY TYPE						
Large Metro	495	448	228	180	173	129
Small Metro	344	309	158	150	96	105
250K - 1 Mil. Pop.	316	266	148	127	90	85
<250K Pop.	145	145	49	78	30	64
Nonmetro	230	228	102	92	72	56
Urbanized	124	134	50	44	32	35
Less Urbanized	170	156	87	74	60	41
Completely Rural	76	93	31	34	30	16

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.96D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.29	0.26	0.13	0.12	0.09	0.08
GEOGRAPHIC DIVISION						
Northeast	0.69	0.51	0.29	0.29	0.24	0.16
New England	1.22	1.08	0.55	0.64	0.44	0.21
Middle Atlantic	0.82	0.58	0.35	0.31	0.29	0.20
Midwest	0.44	0.43	0.24	0.19	0.16	0.15
East North Central	0.52	0.55	0.27	0.25	0.19	0.21
West North Central	0.81	0.62	0.48	0.28	0.28	0.17
South	0.54	0.42	0.23	0.18	0.16	0.12
South Atlantic	0.80	0.60	0.31	0.26	0.25	0.17
East South Central	1.11	0.86	0.74	0.40	0.43	0.32
West South Central	0.90	0.77	0.33	0.32	0.22	0.22
West	0.67	0.72	0.32	0.30	0.21	0.22
Mountain	0.93	0.87	0.59	0.42	0.29	0.24
Pacific	0.86	0.95	0.38	0.38	0.27	0.29
COUNTY TYPE						
Large Metro	0.45	0.41	0.21	0.17	0.16	0.12
Small Metro	0.52	0.45	0.24	0.22	0.14	0.15
250K - 1 Mil. Pop.	0.63	0.52	0.30	0.25	0.18	0.17
<250K Pop.	0.87	0.82	0.29	0.44	0.18	0.36
Nonmetro	0.52	0.50	0.23	0.20	0.16	0.12
Urbanized	0.94	1.01	0.38	0.33	0.24	0.26
Less Urbanized	0.68	0.58	0.35	0.27	0.24	0.15
Completely Rural	1.33	1.76	0.54	0.64	0.52	0.30

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.96P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0230		0.1617		0.6054	
GEOGRAPHIC DIVISION						
Northeast	0.0302		0.4018		0.3950	
New England	0.5443		0.6890		0.0258	
Middle Atlantic	0.0291		0.4617		0.9738	
Midwest	0.0011		0.0006		0.1353	
East North Central	0.1762		0.1063		0.7762	
West North Central	0.0001		0.0002		0.0077	
South	0.1585		0.2548		0.9907	
South Atlantic	0.2852		0.9188		0.6140	
East South Central	0.8101		0.2291		0.7300	
West South Central	0.3534		0.3044		0.5777	
West	0.1463		0.7110		0.3635	
Mountain	0.7216		0.2666		0.9228	
Pacific	0.0842		0.2815		0.3532	
COUNTY TYPE						
Large Metro	0.2200		0.4395		0.5321	
Small Metro	0.0249		0.7024		0.5326	
250K - 1 Mil. Pop.	0.0280		0.1980		0.3850	
<250K Pop.	0.6092		0.0451		0.0049	
Nonmetro	0.7105		0.0697		0.2920	
Urbanized	0.6901		0.1187		0.5882	
Less Urbanized	0.4623		0.1724		0.4803	
Completely Rural	0.2321		0.8948		0.5715	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.97C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	58	61	47	47	30	31
GEOGRAPHIC DIVISION						
Northeast	24	23	20	17	12	12
New England	14	13	13	10	6	5
Middle Atlantic	20	18	16	13	10	11
Midwest	26	25	20	20	13	12
East North Central	21	21	16	18	11	10
West North Central	15	13	12	10	6	6
South	35	43	29	32	19	22
South Atlantic	25	27	19	18	12	13
East South Central	14	19	12	15	9	8
West South Central	20	28	17	22	12	15
West	30	27	23	23	14	15
Mountain	12	12	11	10	6	7
Pacific	27	25	20	20	13	13
COUNTY TYPE						
Large Metro	44	42	35	32	20	20
Small Metro	31	32	26	27	19	19
250K - 1 Mil. Pop.	27	26	22	23	17	16
<250K Pop.	17	19	13	15	8	11
Nonmetro	24	27	20	21	13	13
Urbanized	13	13	11	10	8	6
Less Urbanized	18	21	14	16	9	11
Completely Rural	8	11	7	10	4	5

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.97D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.25	0.26	0.20	0.20	0.13	0.13
GEOGRAPHIC DIVISION						
Northeast	0.57	0.54	0.49	0.40	0.29	0.28
New England	1.27	1.22	1.16	0.94	0.56	0.46
Middle Atlantic	0.64	0.59	0.53	0.43	0.34	0.34
Midwest	0.47	0.46	0.37	0.37	0.23	0.21
East North Central	0.56	0.56	0.43	0.47	0.30	0.26
West North Central	0.86	0.78	0.70	0.57	0.33	0.36
South	0.42	0.52	0.35	0.38	0.23	0.26
South Atlantic	0.62	0.66	0.48	0.45	0.30	0.32
East South Central	0.98	1.35	0.86	1.06	0.62	0.57
West South Central	0.72	0.98	0.62	0.77	0.44	0.55
West	0.56	0.51	0.43	0.42	0.26	0.28
Mountain	0.73	0.71	0.70	0.61	0.35	0.45
Pacific	0.73	0.66	0.54	0.54	0.34	0.35
COUNTY TYPE						
Large Metro	0.38	0.37	0.30	0.28	0.18	0.18
Small Metro	0.45	0.44	0.37	0.38	0.26	0.26
250K - 1 Mil. Pop.	0.50	0.48	0.43	0.42	0.31	0.29
<250K Pop.	1.02	1.03	0.76	0.82	0.49	0.60
Nonmetro	0.52	0.57	0.44	0.43	0.28	0.27
Urbanized	0.84	0.95	0.76	0.73	0.54	0.48
Less Urbanized	0.71	0.75	0.56	0.54	0.36	0.37
Completely Rural	1.36	1.88	1.20	1.72	0.78	0.82

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.97P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 12 to 17, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.9950		0.8173		0.6305	
GEOGRAPHIC DIVISION						
Northeast	0.6251		0.7980		0.6369	
New England	0.7810		0.5642		0.9168	
Middle Atlantic	0.6942		0.4557		0.5684	
Midwest	0.6614		0.7023		0.4824	
East North Central	0.8864		0.6203		0.5674	
West North Central	0.2604		0.1156		0.6624	
South	0.2828		0.4879		0.8714	
South Atlantic	0.3486		0.2945		0.2514	
East South Central	0.0729		0.3338		0.6954	
West South Central	0.6510		0.5434		0.6344	
West	0.1008		0.2094		0.3185	
Mountain	0.0578		0.0510		0.5142	
Pacific	0.3637		0.7032		0.4336	
COUNTY TYPE						
Large Metro	0.5332		0.4237		0.2765	
Small Metro	0.7430		0.2301		0.9185	
250K - 1 Mil. Pop.	0.7474		0.6444		0.7074	
<250K Pop.	0.3228		0.1018		0.6326	
Nonmetro	0.6189		0.3370		0.5567	
Urbanized	0.5788		0.9872		0.7029	
Less Urbanized	0.9740		0.2949		0.7928	
Completely Rural	0.7731		0.6124		0.6050	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.98C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	117	102	86	73	52	50
GEOGRAPHIC DIVISION						
Northeast	63	42	42	36	22	26
New England	29	18	21	14	12	9
Middle Atlantic	56	39	36	34	18	24
Midwest	53	46	38	30	22	18
East North Central	43	41	30	24	18	15
West North Central	31	23	24	18	12	10
South	64	58	52	43	32	29
South Atlantic	44	41	36	27	23	19
East South Central	27	25	24	21	14	14
West South Central	38	33	29	26	18	17
West	55	56	37	37	26	26
Mountain	26	18	17	17	11	11
Pacific	48	53	33	32	24	24
COUNTY TYPE						
Large Metro	85	76	63	54	38	35
Small Metro	67	57	47	44	30	32
250K - 1 Mil. Pop.	51	47	38	37	25	24
<250K Pop.	43	33	26	24	16	19
Nonmetro	42	40	32	29	18	19
Urbanized	27	21	18	17	10	11
Less Urbanized	31	32	24	23	13	14
Completely Rural	12	12	9	7	4	5

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.98D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.41	0.35	0.30	0.25	0.18	0.17
GEOGRAPHIC DIVISION						
Northeast	1.25	0.82	0.83	0.70	0.43	0.50
New England	2.22	1.33	1.64	1.03	0.90	0.66
Middle Atlantic	1.50	1.03	0.96	0.88	0.49	0.64
Midwest	0.77	0.67	0.56	0.44	0.32	0.26
East North Central	0.90	0.85	0.63	0.51	0.39	0.31
West North Central	1.50	1.08	1.15	0.86	0.57	0.46
South	0.63	0.56	0.51	0.41	0.32	0.28
South Atlantic	0.90	0.84	0.75	0.54	0.47	0.38
East South Central	1.47	1.35	1.31	1.13	0.79	0.76
West South Central	1.10	0.92	0.83	0.73	0.51	0.49
West	0.84	0.84	0.58	0.55	0.40	0.40
Mountain	1.31	0.91	0.88	0.86	0.57	0.55
Pacific	1.06	1.14	0.73	0.70	0.52	0.52
COUNTY TYPE						
Large Metro	0.61	0.54	0.45	0.39	0.27	0.25
Small Metro	0.74	0.60	0.52	0.47	0.32	0.34
250K - 1 Mil. Pop.	0.79	0.70	0.59	0.56	0.38	0.36
<250K Pop.	1.56	1.15	0.97	0.86	0.59	0.67
Nonmetro	0.79	0.71	0.59	0.53	0.34	0.34
Urbanized	1.40	1.14	0.95	0.90	0.54	0.58
Less Urbanized	1.07	0.99	0.82	0.72	0.46	0.44
Completely Rural	2.18	2.46	1.76	1.50	0.84	1.11

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.98P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 to 25, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0077		0.0930		0.6614	
GEOGRAPHIC DIVISION						
Northeast	0.7355		0.4222		0.1018	
New England	0.1431		0.7906		0.4864	
Middle Atlantic	0.6728		0.3200		0.1397	
Midwest	0.0313		0.0174		0.1210	
East North Central	0.9172		0.1713		0.4812	
West North Central	0.0000		0.0319		0.0807	
South	0.0152		0.3261		0.8387	
South Atlantic	0.0136		0.1560		0.7525	
East South Central	0.5703		0.7224		0.3505	
West South Central	0.0767		0.7682		0.4983	
West	0.2522		0.3571		0.3554	
Mountain	0.8744		0.8905		0.7991	
Pacific	0.2392		0.2488		0.3736	
COUNTY TYPE						
Large Metro	0.0172		0.3324		0.7216	
Small Metro	0.2123		0.3244		0.9841	
250K - 1 Mil. Pop.	0.3404		0.5699		0.9556	
<250K Pop.	0.4079		0.3406		0.9574	
Nonmetro	0.6024		0.3088		0.5842	
Urbanized	0.8830		0.4817		0.7316	
Less Urbanized	0.3726		0.3845		0.2929	
Completely Rural	0.9815		0.9364		0.8581	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.99C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	621	566	272	243	195	170
GEOGRAPHIC DIVISION						
Northeast	284	213	112	109	100	60
New England	133	122	54	67	51	21
Middle Atlantic	250	176	99	86	87	56
Midwest	214	203	115	89	77	73
East North Central	182	182	89	77	66	68
West North Central	113	89	72	45	39	26
South	406	328	171	136	117	93
South Atlantic	314	248	118	107	88	65
East South Central	156	116	102	49	59	42
West South Central	205	181	74	68	51	51
West	307	346	139	144	91	106
Mountain	125	122	76	57	37	36
Pacific	281	322	116	132	83	100
COUNTY TYPE						
Large Metro	474	432	208	169	161	124
Small Metro	334	295	146	140	86	100
250K - 1 Mil. Pop.	309	257	143	120	84	78
<250K Pop.	138	131	35	72	23	63
Nonmetro	220	221	98	86	68	51
Urbanized	119	126	45	35	30	29
Less Urbanized	162	153	82	71	58	40
Completely Rural	74	90	28	31	27	14

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.99D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.37	0.33	0.16	0.14	0.11	0.10
GEOGRAPHIC DIVISION						
Northeast	0.85	0.64	0.33	0.33	0.30	0.18
New England	1.51	1.38	0.62	0.76	0.58	0.24
Middle Atlantic	1.01	0.71	0.40	0.35	0.35	0.23
Midwest	0.54	0.51	0.29	0.23	0.19	0.19
East North Central	0.66	0.66	0.32	0.28	0.24	0.25
West North Central	0.97	0.76	0.62	0.38	0.34	0.22
South	0.68	0.54	0.28	0.22	0.20	0.15
South Atlantic	0.99	0.77	0.37	0.33	0.28	0.20
East South Central	1.48	1.10	0.97	0.46	0.56	0.40
West South Central	1.14	1.00	0.41	0.37	0.28	0.28
West	0.84	0.93	0.38	0.39	0.25	0.29
Mountain	1.20	1.15	0.73	0.54	0.36	0.33
Pacific	1.07	1.22	0.44	0.50	0.32	0.38
COUNTY TYPE						
Large Metro	0.56	0.52	0.24	0.20	0.19	0.15
Small Metro	0.66	0.57	0.29	0.27	0.17	0.19
250K - 1 Mil. Pop.	0.81	0.66	0.37	0.31	0.22	0.20
<250K Pop.	1.13	1.01	0.29	0.56	0.18	0.49
Nonmetro	0.65	0.63	0.29	0.24	0.20	0.15
Urbanized	1.22	1.25	0.46	0.35	0.31	0.29
Less Urbanized	0.83	0.73	0.42	0.34	0.29	0.19
Completely Rural	1.60	2.15	0.60	0.73	0.58	0.33

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.99P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 26 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0614		0.3253		0.6019	
GEOGRAPHIC DIVISION						
Northeast	0.0201		0.5433		0.1580	
New England	0.3782		0.5864		0.0237	
Middle Atlantic	0.0270		0.7232		0.6467	
Midwest	0.0033		0.0060		0.2792	
East North Central	0.1337		0.1460		0.9815	
West North Central	0.0023		0.0093		0.0245	
South	0.2290		0.2996		0.9900	
South Atlantic	0.3926		0.9204		0.6411	
East South Central	0.5265		0.1190		0.6034	
West South Central	0.5274		0.4078		0.8186	
West	0.0709		0.4160		0.3265	
Mountain	0.8584		0.3379		0.9749	
Pacific	0.0407		0.1483		0.3058	
COUNTY TYPE						
Large Metro	0.4256		0.6646		0.4586	
Small Metro	0.0363		0.7361		0.5133	
250K - 1 Mil. Pop.	0.0388		0.1915		0.4084	
<250K Pop.	0.6717		0.0244		0.0050	
Nonmetro	0.7231		0.1854		0.4054	
Urbanized	0.6163		0.2227		0.6153	
Less Urbanized	0.5217		0.2862		0.6506	
Completely Rural	0.2332		0.8720		0.5593	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.100C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Nonmedical Users of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	639	581	290	258	206	178
GEOGRAPHIC DIVISION						
Northeast	291	218	122	121	100	69
New England	134	121	58	68	47	24
Middle Atlantic	256	183	108	100	88	65
Midwest	223	224	120	97	81	79
East North Central	186	201	95	87	69	73
West North Central	123	98	72	44	43	28
South	417	330	181	144	127	98
South Atlantic	324	249	128	110	101	69
East South Central	151	112	103	53	58	43
West South Central	213	185	79	76	53	54
West	317	354	149	147	98	106
Mountain	129	123	79	60	40	35
Pacific	290	331	127	135	90	100
COUNTY TYPE						
Large Metro	487	447	222	179	169	130
Small Metro	344	304	159	151	95	105
250K - 1 Mil. Pop.	317	260	149	126	89	83
<250K Pop.	145	144	47	80	28	66
Nonmetro	225	227	97	90	69	55
Urbanized	124	132	48	43	29	34
Less Urbanized	164	157	84	73	58	39
Completely Rural	76	92	29	31	27	15

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.100D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.32	0.29	0.15	0.13	0.10	0.09
GEOGRAPHIC DIVISION						
Northeast	0.75	0.56	0.32	0.31	0.26	0.18
New England	1.33	1.20	0.57	0.68	0.47	0.23
Middle Atlantic	0.90	0.64	0.38	0.35	0.31	0.23
Midwest	0.48	0.48	0.26	0.21	0.18	0.17
East North Central	0.57	0.62	0.29	0.27	0.21	0.23
West North Central	0.89	0.70	0.53	0.31	0.31	0.20
South	0.59	0.46	0.26	0.20	0.18	0.14
South Atlantic	0.89	0.67	0.35	0.30	0.28	0.19
East South Central	1.23	0.90	0.84	0.42	0.47	0.35
West South Central	1.00	0.85	0.37	0.35	0.25	0.25
West	0.74	0.81	0.35	0.34	0.23	0.24
Mountain	1.04	0.97	0.64	0.47	0.32	0.28
Pacific	0.94	1.07	0.41	0.43	0.29	0.32
COUNTY TYPE						
Large Metro	0.49	0.46	0.22	0.18	0.17	0.13
Small Metro	0.58	0.50	0.27	0.25	0.16	0.17
250K - 1 Mil. Pop.	0.71	0.57	0.33	0.28	0.20	0.18
<250K Pop.	0.97	0.91	0.32	0.50	0.19	0.42
Nonmetro	0.57	0.56	0.25	0.22	0.17	0.13
Urbanized	1.06	1.10	0.41	0.36	0.25	0.29
Less Urbanized	0.73	0.65	0.37	0.30	0.26	0.16
Completely Rural	1.47	1.97	0.56	0.67	0.52	0.32

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.100P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Nonmedical Use of Any Prescription-Type Psychotherapeutic Among Persons Aged 18 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0222		0.1713		0.5515	
GEOGRAPHIC DIVISION						
Northeast	0.0291		0.4223		0.3524	
New England	0.5354		0.6236		0.0297	
Middle Atlantic	0.0276		0.5259		0.9607	
Midwest	0.0011		0.0005		0.1511	
East North Central	0.1659		0.0754		0.8465	
West North Central	0.0001		0.0002		0.0069	
South	0.1264		0.2152		0.9707	
South Atlantic	0.2517		0.8006		0.7222	
East South Central	0.6246		0.1887		0.7962	
West South Central	0.3700		0.3825		0.6665	
West	0.1118		0.5571		0.4481	
Mountain	0.8425		0.3976		0.9752	
Pacific	0.0690		0.2432		0.4172	
COUNTY TYPE						
Large Metro	0.2363		0.5126		0.4341	
Small Metro	0.0227		0.5633		0.5265	
250K - 1 Mil. Pop.	0.0299		0.1799		0.4310	
<250K Pop.	0.5218		0.0864		0.0073	
Nonmetro	0.6669		0.1003		0.3316	
Urbanized	0.6280		0.1374		0.6763	
Less Urbanized	0.4529		0.2146		0.4873	
Completely Rural	0.2413		0.8747		0.5715	

*Low precision; no estimate reported.

NOTE: Nonmedical use of any prescription-type psychotherapeutic indicates using pain relievers, tranquilizers, stimulants, or sedatives at least once. Indicated use does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.101C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	787	704	341	306	242	209
GEOGRAPHIC DIVISION						
Northeast	368	261	142	155	108	87
New England	170	126	67	100	50	38
Middle Atlantic	328	234	126	119	96	78
Midwest	275	285	133	124	90	92
East North Central	234	239	109	113	76	86
West North Central	145	153	76	52	48	34
South	481	405	217	167	153	110
South Atlantic	375	304	155	124	118	81
East South Central	150	160	109	73	70	50
West South Central	259	213	106	84	69	54
West	427	415	178	165	125	125
Mountain	144	154	92	67	54	36
Pacific	402	385	151	150	113	120
COUNTY TYPE						
Large Metro	629	546	267	217	198	157
Small Metro	408	368	184	184	122	118
250K - 1 Mil. Pop.	370	312	164	151	116	97
<250K Pop.	168	189	76	92	38	66
Nonmetro	267	259	112	104	82	62
Urbanized	141	143	59	54	42	38
Less Urbanized	206	185	95	79	65	47
Completely Rural	91	99	34	38	31	18

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.101D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.36	0.32	0.15	0.14	0.11	0.09
GEOGRAPHIC DIVISION						
Northeast	0.86	0.61	0.33	0.36	0.25	0.20
New England	1.53	1.13	0.60	0.90	0.45	0.34
Middle Atlantic	1.04	0.74	0.40	0.38	0.31	0.25
Midwest	0.53	0.55	0.26	0.24	0.17	0.18
East North Central	0.65	0.66	0.30	0.31	0.21	0.24
West North Central	0.94	0.99	0.49	0.33	0.31	0.22
South	0.61	0.51	0.28	0.21	0.20	0.14
South Atlantic	0.93	0.74	0.38	0.30	0.29	0.20
East South Central	1.09	1.16	0.80	0.53	0.51	0.36
West South Central	1.07	0.87	0.44	0.34	0.29	0.22
West	0.88	0.84	0.37	0.34	0.26	0.25
Mountain	1.03	1.08	0.66	0.47	0.39	0.25
Pacific	1.17	1.11	0.44	0.43	0.33	0.34
COUNTY TYPE						
Large Metro	0.57	0.50	0.24	0.20	0.18	0.14
Small Metro	0.61	0.54	0.28	0.27	0.18	0.17
250K - 1 Mil. Pop.	0.74	0.61	0.33	0.30	0.23	0.19
<250K Pop.	1.00	1.07	0.45	0.52	0.23	0.37
Nonmetro	0.61	0.57	0.25	0.23	0.19	0.14
Urbanized	1.07	1.08	0.45	0.41	0.32	0.29
Less Urbanized	0.82	0.68	0.38	0.29	0.26	0.17
Completely Rural	1.59	1.89	0.59	0.72	0.54	0.34

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.101P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2462		0.2589		0.3178	
GEOGRAPHIC DIVISION						
Northeast	0.1189		0.3358		0.3932	
New England	0.7951		0.2690		0.1173	
Middle Atlantic	0.0609		0.7173		0.8382	
Midwest	0.0513		0.0045		0.0838	
East North Central	0.8242		0.2121		0.6404	
West North Central	0.0016		0.0006		0.0059	
South	0.3725		0.2324		0.5541	
South Atlantic	0.2942		0.7504		0.7709	
East South Central	0.4266		0.4379		0.9494	
West South Central	0.5933		0.2142		0.4636	
West	0.1330		0.6454		0.4865	
Mountain	0.3699		0.4538		0.7890	
Pacific	0.2122		0.3371		0.4073	
COUNTY TYPE						
Large Metro	0.5972		0.6961		0.5049	
Small Metro	0.2056		0.7217		0.9757	
250K - 1 Mil. Pop.	0.1591		0.4022		0.1875	
<250K Pop.	0.9462		0.3844		0.0071	
Nonmetro	0.9285		0.0521		0.1687	
Urbanized	0.9731		0.2272		0.3043	
Less Urbanized	0.9048		0.1065		0.5082	
Completely Rural	0.9376		0.9183		0.3907	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.102C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	73	74	59	60	38	39
GEOGRAPHIC DIVISION						
Northeast	33	28	28	26	17	13
New England	17	16	15	18	10	7
Middle Atlantic	29	24	24	19	14	11
Midwest	33	33	25	25	18	16
East North Central	26	25	21	22	15	15
West North Central	20	21	15	13	9	7
South	45	51	35	40	23	27
South Atlantic	32	32	24	24	15	16
East South Central	16	24	13	19	10	13
West South Central	27	31	22	26	15	17
West	33	32	29	26	18	18
Mountain	16	16	16	15	10	10
Pacific	28	28	24	22	15	15
COUNTY TYPE						
Large Metro	53	53	44	41	27	26
Small Metro	41	40	33	36	23	22
250K - 1 Mil. Pop.	35	33	28	29	20	17
<250K Pop.	21	23	17	21	12	14
Nonmetro	29	32	24	25	15	19
Urbanized	16	18	13	13	9	9
Less Urbanized	22	24	18	19	11	15
Completely Rural	8	13	8	12	5	9

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.102D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.31	0.32	0.26	0.26	0.17	0.17
GEOGRAPHIC DIVISION						
Northeast	0.81	0.68	0.69	0.61	0.41	0.31
New England	1.57	1.45	1.36	1.62	0.90	0.65
Middle Atlantic	0.93	0.77	0.79	0.61	0.45	0.36
Midwest	0.60	0.60	0.46	0.46	0.33	0.30
East North Central	0.70	0.67	0.54	0.58	0.41	0.39
West North Central	1.15	1.24	0.86	0.77	0.51	0.42
South	0.54	0.61	0.43	0.48	0.28	0.32
South Atlantic	0.80	0.79	0.59	0.59	0.38	0.40
East South Central	1.13	1.70	0.92	1.37	0.69	0.91
West South Central	0.95	1.08	0.80	0.91	0.52	0.59
West	0.61	0.60	0.54	0.49	0.34	0.33
Mountain	1.01	0.99	1.02	0.92	0.61	0.62
Pacific	0.76	0.74	0.63	0.58	0.41	0.40
COUNTY TYPE						
Large Metro	0.46	0.46	0.38	0.36	0.24	0.23
Small Metro	0.59	0.56	0.47	0.50	0.33	0.31
250K - 1 Mil. Pop.	0.66	0.61	0.52	0.53	0.38	0.32
<250K Pop.	1.22	1.29	0.99	1.15	0.68	0.76
Nonmetro	0.62	0.66	0.52	0.53	0.32	0.40
Urbanized	1.04	1.37	0.89	0.98	0.63	0.67
Less Urbanized	0.83	0.82	0.68	0.66	0.42	0.51
Completely Rural	1.49	2.12	1.43	2.01	0.85	1.56

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.102P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 12 to 17, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.6289		0.3114		0.6965	
GEOGRAPHIC DIVISION						
Northeast	0.3627		0.1121		0.3109	
New England	0.9527		0.5870		0.7310	
Middle Atlantic	0.2788		0.1182		0.3221	
Midwest	0.9419		0.2725		0.4583	
East North Central	0.7331		0.5568		0.4464	
West North Central	0.6837		0.2483		0.8955	
South	0.3003		0.2242		0.0862	
South Atlantic	0.1653		0.1454		0.0184	
East South Central	0.2675		0.1997		0.4050	
West South Central	0.5490		0.6611		0.8176	
West	0.0679		0.2105		0.8654	
Mountain	0.1530		0.5021		0.6308	
Pacific	0.1812		0.2890		0.9182	
COUNTY TYPE						
Large Metro	0.2597		0.1669		0.9366	
Small Metro	0.6707		0.6717		0.7865	
250K - 1 Mil. Pop.	0.8029		0.9730		0.8671	
<250K Pop.	0.2843		0.5017		0.4226	
Nonmetro	0.9348		0.4233		0.7308	
Urbanized	0.9431		0.8511		0.5895	
Less Urbanized	0.8014		0.3954		0.8568	
Completely Rural	0.2877		0.5173		0.9427	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.103C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	139	133	104	98	64	62
GEOGRAPHIC DIVISION						
Northeast	65	53	47	45	26	28
New England	39	25	30	20	14	13
Middle Atlantic	50	47	35	40	21	26
Midwest	64	56	50	39	29	24
East North Central	53	45	41	33	25	20
West North Central	35	35	29	21	15	13
South	79	70	61	55	38	34
South Atlantic	53	51	40	37	29	23
East South Central	27	28	26	27	15	15
West South Central	52	40	37	31	20	20
West	69	83	50	55	33	36
Mountain	34	27	25	24	15	18
Pacific	60	78	43	49	29	31
COUNTY TYPE						
Large Metro	100	104	78	73	49	44
Small Metro	82	65	58	57	34	36
250K - 1 Mil. Pop.	61	54	45	47	28	29
<250K Pop.	54	38	35	33	19	20
Nonmetro	54	46	38	36	22	23
Urbanized	28	23	22	22	14	15
Less Urbanized	43	39	29	27	15	17
Completely Rural	16	13	12	10	10	5

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.103D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.49	0.46	0.37	0.34	0.22	0.21
GEOGRAPHIC DIVISION						
Northeast	1.28	1.03	0.92	0.87	0.51	0.56
New England	3.02	1.87	2.30	1.49	1.09	0.97
Middle Atlantic	1.34	1.24	0.94	1.06	0.57	0.67
Midwest	0.94	0.82	0.73	0.57	0.43	0.35
East North Central	1.13	0.93	0.86	0.69	0.52	0.43
West North Central	1.72	1.66	1.40	1.02	0.72	0.60
South	0.78	0.68	0.60	0.53	0.38	0.33
South Atlantic	1.09	1.03	0.84	0.75	0.59	0.46
East South Central	1.48	1.54	1.45	1.45	0.84	0.82
West South Central	1.50	1.14	1.06	0.87	0.59	0.58
West	1.06	1.25	0.76	0.82	0.50	0.53
Mountain	1.77	1.35	1.30	1.21	0.78	0.89
Pacific	1.31	1.68	0.94	1.05	0.63	0.66
COUNTY TYPE						
Large Metro	0.72	0.75	0.56	0.52	0.35	0.32
Small Metro	0.89	0.68	0.63	0.60	0.37	0.38
250K - 1 Mil. Pop.	0.95	0.82	0.70	0.71	0.44	0.44
<250K Pop.	1.98	1.35	1.26	1.14	0.70	0.68
Nonmetro	1.00	0.83	0.71	0.65	0.41	0.42
Urbanized	1.44	1.20	1.14	1.16	0.71	0.77
Less Urbanized	1.47	1.19	1.00	0.83	0.53	0.54
Completely Rural	3.00	2.59	2.29	2.05	1.95	1.11

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.103P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 to 25, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.0383		0.0825		0.5779	
GEOGRAPHIC DIVISION						
Northeast	0.8313		0.5505		0.1623	
New England	0.5134		0.7031		0.5488	
Middle Atlantic	0.4416		0.6533		0.2118	
Midwest	0.1129		0.0177		0.0508	
East North Central	0.8040		0.1715		0.2980	
West North Central	0.0016		0.0271		0.0436	
South	0.0613		0.3582		0.8941	
South Atlantic	0.0905		0.3026		0.6023	
East South Central	0.5296		0.5597		0.1902	
West South Central	0.1098		0.4262		0.6102	
West	0.6263		0.5073		0.7549	
Mountain	0.9536		0.8321		0.6314	
Pacific	0.5713		0.5197		0.5271	
COUNTY TYPE						
Large Metro	0.3431		0.3582		0.6383	
Small Metro	0.2953		0.4488		0.5530	
250K - 1 Mil. Pop.	0.4911		0.6103		0.6334	
<250K Pop.	0.3879		0.5395		0.7038	
Nonmetro	0.0309		0.0956		0.1262	
Urbanized	0.5959		0.9759		0.6660	
Less Urbanized	0.0245		0.0299		0.2783	
Completely Rural	0.7052		0.7540		0.2364	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.104C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	746	662	308	274	225	195
GEOGRAPHIC DIVISION						
Northeast	351	244	129	136	105	80
New England	168	116	60	92	53	32
Middle Atlantic	310	217	115	100	92	73
Midwest	257	258	120	107	83	83
East North Central	220	217	93	97	68	76
West North Central	132	138	76	46	46	32
South	455	384	196	153	140	103
South Atlantic	357	290	139	121	104	76
East South Central	143	157	107	60	72	45
West South Central	238	195	90	73	62	53
West	406	389	159	148	114	119
Mountain	138	144	78	60	49	36
Pacific	383	361	137	136	103	114
COUNTY TYPE						
Large Metro	600	508	239	197	182	152
Small Metro	380	345	168	160	116	106
250K - 1 Mil. Pop.	350	298	159	137	114	85
<250K Pop.	152	166	56	76	27	65
Nonmetro	257	247	98	91	75	55
Urbanized	136	135	50	39	36	30
Less Urbanized	196	176	83	75	62	43
Completely Rural	91	94	30	32	27	14

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.104D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.44	0.39	0.18	0.16	0.13	0.11
GEOGRAPHIC DIVISION						
Northeast	1.05	0.73	0.39	0.41	0.31	0.24
New England	1.91	1.32	0.69	1.04	0.61	0.37
Middle Atlantic	1.26	0.88	0.47	0.40	0.37	0.30
Midwest	0.65	0.65	0.31	0.27	0.21	0.21
East North Central	0.80	0.78	0.34	0.35	0.25	0.28
West North Central	1.13	1.17	0.65	0.39	0.40	0.27
South	0.76	0.63	0.33	0.25	0.23	0.17
South Atlantic	1.13	0.90	0.44	0.38	0.33	0.24
East South Central	1.37	1.48	1.02	0.56	0.68	0.42
West South Central	1.32	1.07	0.50	0.40	0.35	0.29
West	1.11	1.05	0.43	0.40	0.31	0.32
Mountain	1.33	1.35	0.75	0.56	0.47	0.33
Pacific	1.46	1.37	0.52	0.51	0.39	0.43
COUNTY TYPE						
Large Metro	0.71	0.61	0.28	0.23	0.21	0.18
Small Metro	0.75	0.66	0.33	0.31	0.23	0.20
250K - 1 Mil. Pop.	0.91	0.77	0.42	0.35	0.30	0.22
<250K Pop.	1.24	1.28	0.46	0.58	0.22	0.50
Nonmetro	0.76	0.70	0.29	0.26	0.22	0.16
Urbanized	1.40	1.34	0.51	0.39	0.37	0.30
Less Urbanized	1.00	0.84	0.43	0.36	0.31	0.21
Completely Rural	1.97	2.25	0.64	0.77	0.58	0.33

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.104P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 26 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.4316		0.6127		0.3108	
GEOGRAPHIC DIVISION						
Northeast	0.1424		0.2622		0.2744	
New England	0.9187		0.2594		0.1201	
Middle Atlantic	0.0906		0.5949		0.6876	
Midwest	0.0783		0.0446		0.2577	
East North Central	0.7190		0.4221		0.9957	
West North Central	0.0065		0.0247		0.0370	
South	0.4441		0.2050		0.3434	
South Atlantic	0.3129		0.7564		0.5731	
East South Central	0.6120		0.1794		0.5586	
West South Central	0.8544		0.3293		0.5730	
West	0.0639		0.3430		0.4367	
Mountain	0.3014		0.5233		0.5750	
Pacific	0.1165		0.1545		0.2998	
COUNTY TYPE						
Large Metro	0.7739		0.9258		0.5349	
Small Metro	0.2373		0.8025		0.7848	
250K - 1 Mil. Pop.	0.1808		0.4282		0.1311	
<250K Pop.	0.8625		0.2500		0.0094	
Nonmetro	0.7526		0.1903		0.2665	
Urbanized	0.9274		0.2323		0.3312	
Less Urbanized	0.7643		0.3195		0.6029	
Completely Rural	0.9933		0.8472		0.5444	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.105C Standard Errors of Estimated Numbers (in Thousands) of Lifetime, Past Year, and Past Month Users of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	776	693	330	300	239	208
GEOGRAPHIC DIVISION						
Northeast	365	254	136	149	106	88
New England	173	120	66	91	52	37
Middle Atlantic	322	228	120	118	93	80
Midwest	273	285	132	120	89	90
East North Central	231	236	106	108	74	83
West North Central	146	157	78	51	50	35
South	473	393	211	166	152	110
South Atlantic	372	298	153	127	117	82
East South Central	144	153	107	69	71	49
West South Central	251	205	99	82	68	54
West	421	411	169	163	123	125
Mountain	142	152	85	63	52	37
Pacific	397	381	145	150	111	119
COUNTY TYPE						
Large Metro	620	537	257	215	195	158
Small Metro	402	358	182	177	122	117
250K - 1 Mil. Pop.	366	304	164	147	117	95
<250K Pop.	165	185	72	89	34	66
Nonmetro	264	256	105	100	79	60
Urbanized	141	139	58	52	41	37
Less Urbanized	201	183	90	77	64	44
Completely Rural	91	98	30	34	28	15

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.105D Standard Errors of Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 1999 and 2000

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.39	0.35	0.17	0.15	0.12	0.10
GEOGRAPHIC DIVISION						
Northeast	0.95	0.66	0.35	0.39	0.27	0.23
New England	1.72	1.19	0.66	0.90	0.52	0.37
Middle Atlantic	1.13	0.80	0.42	0.41	0.33	0.28
Midwest	0.59	0.61	0.29	0.26	0.19	0.19
East North Central	0.71	0.73	0.33	0.33	0.23	0.25
West North Central	1.06	1.13	0.57	0.37	0.37	0.25
South	0.67	0.55	0.30	0.23	0.22	0.15
South Atlantic	1.02	0.81	0.42	0.34	0.32	0.22
East South Central	1.17	1.23	0.87	0.56	0.58	0.40
West South Central	1.17	0.94	0.47	0.38	0.32	0.25
West	0.98	0.94	0.39	0.37	0.28	0.29
Mountain	1.15	1.20	0.69	0.50	0.42	0.29
Pacific	1.29	1.23	0.47	0.48	0.36	0.38
COUNTY TYPE						
Large Metro	0.63	0.55	0.26	0.22	0.20	0.16
Small Metro	0.67	0.58	0.31	0.29	0.21	0.19
250K - 1 Mil. Pop.	0.82	0.67	0.37	0.32	0.26	0.21
<250K Pop.	1.11	1.17	0.48	0.57	0.22	0.42
Nonmetro	0.67	0.63	0.27	0.25	0.20	0.15
Urbanized	1.21	1.17	0.49	0.43	0.35	0.31
Less Urbanized	0.89	0.76	0.40	0.32	0.28	0.18
Completely Rural	1.76	2.10	0.59	0.73	0.55	0.32

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.105P P-Values from Tests of Differences in Percentages Reporting Lifetime, Past Year, and Past Month Use of Any Illicit Drug Other Than Marijuana Among Persons Aged 18 or Older, by Geographic Characteristics: 2000 Versus 1999

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	1999	2000	1999	2000	1999	2000
TOTAL	0.2597		0.3386		0.2835	
GEOGRAPHIC DIVISION						
Northeast	0.1439		0.2048		0.4837	
New England	0.7875		0.2054		0.1513	
Middle Atlantic	0.0750		0.5210		0.9521	
Midwest	0.0459		0.0065		0.1000	
East North Central	0.7876		0.2343		0.7446	
West North Central	0.0014		0.0014		0.0076	
South	0.3041		0.1560		0.3724	
South Atlantic	0.2347		0.5874		0.5318	
East South Central	0.5267		0.3167		0.8252	
West South Central	0.6297		0.2372		0.4982	
West	0.0947		0.4808		0.4986	
Mountain	0.3088		0.5253		0.7138	
Pacific	0.1685		0.2435		0.3969	
COUNTY TYPE						
Large Metro	0.6643		0.8813		0.4929	
Small Metro	0.1796		0.6555		0.9356	
250K - 1 Mil. Pop.	0.1598		0.3992		0.2065	
<250K Pop.	0.9258		0.4495		0.0094	
Nonmetro	0.9201		0.0715		0.1246	
Urbanized	0.9548		0.2681		0.2760	
Less Urbanized	0.9258		0.1225		0.4495	
Completely Rural	0.9392		0.9093		0.2960	

*Low precision; no estimate reported.

NOTE: Any Illicit Drug Other Than Marijuana indicates use at least once of cocaine (including crack), heroin, hallucinogens (including LSD and PCP), inhalants, or any prescription-type psychotherapeutic used nonmedically, regardless of marijuana/hashish use; marijuana/hashish users who also have used any of the other listed drugs are included.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.106C Standard Errors of Estimated Numbers (in Thousands) of Lifetime Users of Specific Hallucinogens, Inhalants, Needles, and Heroin, by Age Group: 1999 and 2000

Drug or Method of Administration	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Hallucinogens								
PCP	557	487	45	43	118	118	525	453
LSD	298	279	17	18	39	37	295	275
Peyote	479	433	35	36	105	99	456	408
Mescaline	268	275	11	12	34	35	266	270
Psilocybin (Mushrooms)	364	308	12	12	42	36	358	302
"Ecstasy" (MDMA)	440	395	28	31	98	91	412	381
	215	225	25	31	75	89	189	188
Inhalants								
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	443	381	54	49	98	94	416	355
Correction Fluid, Degreaser, or Cleaning Fluid	328	277	25	21	47	43	323	273
Gasoline or Lighter Fluid	135	96	26	24	33	29	129	90
Glue, Shoe Polish, or Toluene	155	136	32	32	44	39	142	131
Halothane, Ether, or Other Anesthetics	169	161	36	34	39	33	157	151
Lacquer Thinner or Other Paint Solvents	102	103	12	10	25	21	95	103
Lighter Gases (Butane, Propane)	120	103	24	22	32	27	109	97
Nitrous Oxide or "Whippets"	79	66	18	17	28	20	73	62
Spray Paints	305	246	25	24	88	83	278	223
Other Aerosol Sprays	88	98	29	25	30	27	76	93
	106	119	25	21	33	29	95	113
Any Needle Use^{1,2}								
Heroin Needle Use ^{1,2}	226	188	11	8	28	24	226	185
Cocaine Needle Use ^{1,2}	163	132	6	5	18	18	163	129
Stimulant Needle Use ^{1,2}	191	146	5	4	18	14	190	144
	122	134	5	4	17	12	122	134
Heroin								
Smoke Heroin ²	229	202	13	10	38	33	224	198
Sniff or Snort Heroin ²	124	99	8	6	21	20	120	97
	173	160	6	5	29	24	170	158

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.² These drug estimates are generated from data collected on a "non-core" module of the NHSDA questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NHSDA data are not edited for consistency across the "core" and "non-core" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.106D Standard Errors of Percentages Reporting Lifetime Use of Specific Hallucinogens, Inhalants, Needles, and Heroin, by Age Group: 2000 Versus 1999

Drug or Method of Administration	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Hallucinogens								
PCP	0.25	0.22	0.19	0.18	0.42	0.41	0.31	0.26
LSD	0.13	0.12	0.07	0.08	0.14	0.13	0.17	0.16
Peyote	0.22	0.19	0.15	0.15	0.37	0.34	0.27	0.24
Mescaline	0.12	0.12	0.05	0.05	0.12	0.12	0.16	0.16
Psilocybin (Mushrooms)	0.16	0.14	0.05	0.05	0.15	0.13	0.21	0.18
"Ecstasy" (MDMA)	0.20	0.18	0.12	0.13	0.35	0.31	0.24	0.22
"Ecstasy" (MDMA)	0.10	0.10	0.11	0.13	0.26	0.31	0.11	0.11
Inhalants								
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	0.20	0.17	0.23	0.21	0.34	0.32	0.25	0.21
Correction Fluid, Degreaser, or Cleaning Fluid	0.15	0.12	0.11	0.09	0.17	0.15	0.19	0.16
Gasoline or Lighter Fluid	0.06	0.04	0.11	0.10	0.12	0.10	0.08	0.05
Glue, Shoe Polish, or Toluene	0.07	0.06	0.14	0.14	0.16	0.13	0.08	0.08
Halothane, Ether, or Other Anesthetics	0.05	0.05	0.05	0.04	0.09	0.07	0.06	0.06
Lacquer Thinner or Other Paint Solvents	0.05	0.05	0.11	0.09	0.11	0.09	0.06	0.06
Lighter Gases (Butane, Propane)	0.05	0.05	0.08	0.07	0.10	0.07	0.04	0.04
Nitrous Oxide or "Whippets"	0.04	0.03	0.11	0.10	0.31	0.29	0.16	0.13
Spray Paints	0.04	0.04	0.12	0.11	0.11	0.09	0.04	0.05
Other Aerosol Sprays	0.05	0.05	0.11	0.09	0.12	0.10	0.06	0.07
Any Needle Use^{1,2}								
Heroin Needle Use ^{1,2}	0.10	0.08	0.05	0.03	0.10	0.08	0.13	0.11
Cocaine Needle Use ^{1,2}	0.07	0.06	0.03	0.02	0.06	0.06	0.10	0.08
Stimulant Needle Use ^{1,2}	0.09	0.07	0.02	0.02	0.06	0.05	0.11	0.08
Heroin								
Heroin	0.10	0.09	0.06	0.04	0.13	0.11	0.13	0.12
Smoke Heroin ²	0.06	0.04	0.03	0.02	0.07	0.07	0.07	0.06
Sniff or Snort Heroin ²	0.08	0.07	0.03	0.02	0.10	0.08	0.10	0.09

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.² These drug estimates are generated from data collected on a "non-core" module of the NHSDA questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NHSDA data are not edited for consistency across the "core" and "non-core" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.106P P-Values from Tests of Differences in Percentages Reporting Lifetime Use of Specific Hallucinogens, Inhalants, Needles, and Heroin, by Age Group: 1999 and 2000

Drug or Method of Administration	Total	AGE GROUP (Years)						
		12-17		18-25		26 or Older		
	1999	2000	1999	2000	1999	2000	1999	2000
Hallucinogens								
PCP	0.2561		0.7335		0.9899		0.2595	
LSD	0.8955		0.2710		0.5382		0.8832	
Peyote	0.7023		0.3984		0.1915		0.4486	
Mescaline	0.7570		0.5895		0.8934		0.7250	
Psilocybin (Mushrooms)	0.5342		0.5656		0.8799		0.5261	
"Ecstasy" (MDMA)	0.5319		0.3794		0.3540		0.4330	
	0.0000		0.0000		0.0000		0.0444	
Inhalants	0.0000		0.2871		0.4586		0.0047	
Amyl Nitrite, "Poppers," Locker Room Odorizers, or "Rush"	0.0284		0.0284		0.4823		0.0107	
Correction Fluid, Degreaser, or Cleaning Fluid	0.0200		0.0200		0.4902		0.0016	
Gasoline or Lighter Fluid	0.7950		0.7950		0.6124		0.3403	
Glue, Shoe Polish, or Toluene	0.6978		0.6978		0.7587		0.0231	
Halothane, Ether, or Other Anesthetics	0.6849		0.6849		0.2503		0.0244	
Lacquer Thinner or Other Paint Solvents	0.6821		0.6821		0.2034		0.0373	
Lighter Gases (Butane, Propane)	0.5859		0.5859		0.7080		0.0625	
Nitrous Oxide or "Whippets"	0.5859		0.3993		0.9142		0.4099	
Spray Paints	0.3993		0.9428		0.0503		0.3041	
Other Aerosol Sprays	0.9428		0.8822		0.3569		0.4609	
	0.8822		0.0000		0.0000		0.8044	
Any Needle Use^{1,2}	0.1321		0.1321		0.3159		0.2818	
Heroin Needle Use ^{1,2}	0.1766		0.1766		0.4934		0.9543	
Cocaine Needle Use ^{1,2}	0.1101		0.1101		0.1545		0.2672	
Stimulant Needle Use ^{1,2}	0.8585		0.8585		0.5960		0.0659	
Heroin	0.3171		0.3171		0.2640		0.0424	
Smoke Heroin ²	0.9962		0.9962		0.5736		0.3727	
Sniff or Snort Heroin ²	0.6650		0.6650		0.8415		0.0298	
	0.6650		0.0000		0.0000		0.4092	

*Low precision; no estimate reported.

¹ Refers to lifetime use of a needle to inject a drug that was not prescribed, or that was taken only for the experience or feeling it caused.

² These drug estimates are generated from data collected on a "non-core" module of the NHSDA questionnaire. Therefore, these estimates may be somewhat inconsistent with drug estimates based on data collected from the "core" modules because NHSDA data are not edited for consistency across the "core" and "non-core" modules of the interview.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.107C Standard Errors of Estimated Numbers (in Thousands) of Lifetime Users of Specific Pain Relievers, by Age Group: 1999 and 2000

Pain Relievers	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Darvocet®, Darvon®, or Tylenol® with Codeine	413	381	40	43	80	72	406	361
Percocet®, Percodan®, or Tylox®	283	258	20	20	58	54	275	247
Vicodin®, Lortab®, or Loracet®	261	229	23	29	65	68	249	207
Codeine	236	213	21	25	52	50	226	203
Demerol®	188	154	11	14	34	32	185	145
Dilaudid®	108	110	4	5	13	12	107	108
Fioricet®	67	48	3	4	11	8	66	47
Fiorinal®	77	55	5	3	8	8	77	54
Hydrocodone	129	114	12	17	35	41	125	103
Methadone	62	109	10	9	19	22	58	106
Morphine	110	134	14	13	28	31	107	126
Oxycontin®	46	47	5	7	12	21	44	42
Phenaphen® with Codeine	75	101	8	7	18	21	73	99
Propoxyphene	56	33	5	4	9	7	55	32
SK-65®	10	29	5	4	8	3	*	28
Stadol®	28	14	3	*	7	8	26	12
Talacen®	28	30	5	2	3	6	27	30
Talwin®	84	69	3	3	6	7	83	69
Talwin NX®	22	24	4	2	3	2	22	24
Tramadol	10	31	3	5	8	8	*	30
Ultram®	57	53	7	9	16	21	55	47

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.107D Standard Errors of Percentages Reporting Lifetime Use of Specific Pain Relievers, by Age Group: 1999 and 2000

Pain Relievers	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Darvocet®, Darvon®, or Tylenol® with Codeine	0.19	0.17	0.17	0.18	0.28	0.25	0.24	0.21
Percocet®, Percodan®, or Tylox®	0.13	0.12	0.09	0.09	0.20	0.19	0.16	0.14
Vicodin®, Lortab®, or Loracet®	0.12	0.10	0.10	0.12	0.23	0.23	0.15	0.12
Codeine	0.11	0.10	0.09	0.11	0.18	0.17	0.13	0.12
Demerol®	0.08	0.07	0.05	0.06	0.12	0.11	0.11	0.08
Dilaudid®	0.05	0.05	0.02	0.02	0.04	0.04	0.06	0.06
Fioricet®	0.03	0.02	0.01	0.02	0.04	0.03	0.04	0.03
Fiorinal®	0.03	0.02	0.02	0.01	0.03	0.03	0.05	0.03
Hydrocodone	0.06	0.05	0.05	0.07	0.12	0.14	0.07	0.06
Methadone	0.03	0.05	0.04	0.04	0.07	0.07	0.03	0.06
Morphine	0.05	0.06	0.06	0.06	0.10	0.11	0.06	0.07
Oxycontin®	0.02	0.02	0.02	0.03	0.04	0.07	0.03	0.02
Phenaphen® with Codeine	0.03	0.05	0.03	0.03	0.06	0.07	0.04	0.06
Propoxyphene	0.03	0.01	0.02	0.02	0.03	0.02	0.03	0.02
SK-65®	0.00	0.01	0.02	0.02	0.03	0.01	*	0.02
Stadol®	0.01	0.01	0.01	*	0.02	0.03	0.02	0.01
Talacen®	0.01	0.01	0.02	0.01	0.01	0.02	0.02	0.02
Talwin®	0.04	0.03	0.01	0.01	0.02	0.02	0.05	0.04
Talwin NX®	0.01	0.01	0.02	0.01	0.01	0.01	0.01	0.01
Tramadol	0.00	0.01	0.01	0.02	0.03	0.03	*	0.02
Ultram®	0.03	0.02	0.03	0.04	0.06	0.07	0.03	0.03

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.107P P-Values from Tests of Differences in Percentages Reporting Lifetime Use of Specific Pain Relievers, by Age Group: 2000 Versus 1999

Pain Relievers	Total	AGE GROUP (Years)					
		12-17		18-25		26 or Older	
		1999	2000	1999	2000	1999	2000
Darvocet®, Darvon®, or Tylenol® with Codeine	0.0783	0.5342		0.0029		0.1894	
Percocet®, Percodan®, or Tylox®	0.8299	0.8571		0.7113		0.8619	
Vicodin®, Lortab®, or Loracet®	0.0892	0.0003		0.0205		0.4946	
Codeine	0.9393	0.0577		0.7959		0.7050	
Demerol®	0.4464	0.1007		0.5945		0.4122	
Dilaudid®	0.6486	0.5500		0.8551		0.6452	
Fioricet®	0.7399	0.7236		0.3333		0.8329	
Fiorinal®	0.3747	0.3462		0.9283		0.4003	
Hydrocodone	0.4019	0.0438		0.0003		0.5853	
Methadone	0.0054	0.7809		0.1309		0.0123	
Morphine	0.0777	0.3377		0.0662		0.2171	
Oxycontin®	0.0100	0.0172		0.0000		0.4332	
Phenaphen® with Codeine	0.1130	0.8627		0.4607		0.1451	
Propoxyphene	0.2968	0.4911		0.6560		0.3608	
SK-65®	0.2730	0.9855		0.0982		*	
Stadol®	0.5063	*		0.8127		0.5949	
Talacen®	0.9728	0.0416		0.1608		0.9319	
Talwin®	0.2829	0.8657		0.5729		0.3070	
Talwin NX®	0.9833	0.1971		0.0798		0.7225	
Tramadol	0.2224	0.6129		0.8876		*	
Ultram®	0.4598	0.2944		0.0756		0.9546	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.108C Standard Errors of Estimated Numbers (in Thousands) of Lifetime Users of Specific Tranquilizers, by Age Group: 1999 and 2000

Tranquilizers	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Klonopin® or Clonazepam	129	110	10	11	29	32	123	102
Xanax®, Alprazolam, Ativan®, or Lorazepam	258	215	18	21	55	53	251	198
Valium® or Diazepam	403	367	21	24	65	63	395	350
Atarax®	73	72	5	5	9	10	72	71
BuSpar®	67	53	5	9	13	18	65	49
Equanil®	12	39	4	4	6	5	11	38
Flexeril®	87	111	6	6	17	22	86	109
Librium®	125	103	6	5	11	14	124	102
Limbitrol®	13	24	4	3	3	7	12	23
Meprobamate	34	57	4	3	6	8	34	56
Miltown®	30	46	3	3	7	4	29	45
Rohypnol®	36	39	7	7	17	18	30	31
Serax®	44	19	4	4	10	8	43	17
Soma®	108	80	9	10	27	31	104	73
Tranxene®	60	34	5	4	9	8	61	33
Vistaril®	49	42	5	4	6	9	49	41

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.108D Standard Errors of Percentages Reporting Lifetime Use of Specific Tranquilizers, by Age Group: 1999 and 2000

Tranquilizers	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Klonopin® or Clonazepam	0.06	0.05	0.04	0.05	0.10	0.11	0.07	0.06
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.12	0.10	0.08	0.09	0.19	0.18	0.15	0.12
Valium® or Diazepam	0.18	0.16	0.09	0.10	0.23	0.22	0.23	0.20
Atarax®	0.03	0.03	0.02	0.02	0.03	0.03	0.04	0.04
BuSpar®	0.03	0.02	0.02	0.04	0.04	0.06	0.04	0.03
Equanil®	0.01	0.02	0.02	0.02	0.02	0.02	0.01	0.02
Flexeril®	0.04	0.05	0.02	0.03	0.06	0.08	0.05	0.06
Librium®	0.06	0.05	0.03	0.02	0.04	0.05	0.07	0.06
Limbitrol®	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01
Meprobamate	0.02	0.03	0.02	0.01	0.02	0.03	0.02	0.03
Miltown®	0.01	0.02	0.01	0.01	0.02	0.01	0.02	0.03
Rohypnol®	0.02	0.02	0.03	0.03	0.06	0.06	0.02	0.02
Serax®	0.02	0.01	0.02	0.02	0.03	0.03	0.03	0.01
Soma®	0.05	0.04	0.04	0.04	0.09	0.11	0.06	0.04
Tranxene®	0.03	0.02	0.02	0.02	0.03	0.03	0.04	0.02
Vistaril®	0.02	0.02	0.02	0.02	0.02	0.03	0.03	0.02

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.108P P-Values from Tests of Differences in Percentages Reporting Lifetime Use of Specific Tranquilizers, by Age Group: 2000 Versus 1999

Tranquilizers	Total	AGE GROUP (Years)					
		12-17		18-25		26 or Older	
		1999	2000	1999	2000	1999	2000
Klonopin® or Clonazepam	0.7987		0.2092		0.0510		0.3482
Xanax®, Alprazolam, Ativan®, or Lorazepam	0.2125		0.6260		0.4060		0.2397
Valium® or Diazepam	0.1016		0.5051		0.2063		0.1343
Atarax®	0.9855		0.6912		0.2408		0.8828
BuSpar®	0.3486		0.0054		0.0356		0.9742
Equanil®	0.1357		0.6172		0.7619		0.1300
Flexeril®	0.1524		0.7305		0.2218		0.2300
Librium®	0.7156		0.6753		0.2737		0.6355
Limbitrol®	0.1805		0.5011		0.0943		0.2827
Meprobamate	0.6663		0.4664		0.6541		0.6710
Miltown®	0.2358		0.5766		0.1571		0.1731
Rohypnol®	0.7918		0.3107		0.9410		0.9109
Serax®	0.4506		0.8319		0.4196		0.2982
Soma®	0.9777		0.6155		0.2809		0.7021
Tranxene®	0.1228		0.7970		0.4489		0.1631
Vistaril®	0.3901		0.4832		0.0355		0.2548

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.109C Standard Errors of Estimated Numbers (in Thousands) of Lifetime Users of Specific Stimulants, by Age Group: 1999 and 2000

Stimulants	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Methamphetamine, Desoxyn®, or Methedrine	383	296	20	19	58	54	369	289
Prescription Diet Pills ¹	342	300	18	19	36	36	339	297
Ritalin® or Methylphenidate	160	153	27	26	52	48	144	144
Cylert®	82	51	6	5	10	10	81	49
Dexedrine®	198	193	7	9	19	20	197	191
Dextroamphetamine	96	93	4	6	12	15	95	92
Didrex®	58	48	3	5	6	6	57	47
Eskatrol®	42	42	3	3	6	11	42	40
Ionamin®	79	61	3	5	6	13	78	59
Mazanor®	29	32	3	5	6	5	28	31
Obedrin-LA®	18	38	2	2	4	5	18	38
Plegine®	11	36	2	4	2	2	10	35
Preludin®	81	110	4	3	11	10	80	109
Sanorex®	41	39	5	4	7	7	39	38
Tenuate®	23	28	3	3	6	7	22	27

*Low precision; no estimate reported.

¹ For example, Amphetamines, Benzedrine®, Biphetamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.109D Standard Errors of Percentages Reporting Lifetime Use of Specific Stimulants, by Age Group: 1999 and 2000

Stimulants	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Methamphetamine, Desoxyn®, or Methedrine	0.17	0.13	0.09	0.08	0.20	0.19	0.22	0.17
Prescription Diet Pills ¹	0.15	0.13	0.08	0.08	0.13	0.12	0.20	0.17
Ritalin® or Methylphenidate	0.07	0.07	0.11	0.11	0.18	0.17	0.09	0.08
Cylert®	0.04	0.02	0.02	0.02	0.04	0.04	0.05	0.03
Dexedrine®	0.09	0.09	0.03	0.04	0.07	0.07	0.12	0.11
Dextroamphetamine	0.04	0.04	0.02	0.03	0.04	0.05	0.06	0.05
Didrex®	0.03	0.02	0.01	0.02	0.02	0.02	0.03	0.03
Eskatrol®	0.02	0.02	0.01	0.01	0.02	0.04	0.02	0.02
Ionamin®	0.04	0.03	0.01	0.02	0.02	0.05	0.05	0.03
Mazanor®	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.02
Obedrin-LA®	0.01	0.02	0.01	0.01	0.01	0.02	0.01	0.02
Plegine®	0.00	0.02	0.01	0.02	0.01	0.01	0.01	0.02
Preludin®	0.04	0.05	0.02	0.01	0.04	0.04	0.05	0.06
Sanorex®	0.02	0.02	0.02	0.02	0.03	0.02	0.02	0.02
Tenuate®	0.01	0.01	0.01	0.01	0.02	0.03	0.01	0.02

*Low precision; no estimate reported.

¹ For example, Amphetamines, Benzedrine®, Biphetamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.109P P-Values from Tests of Differences in Percentages Reporting Lifetime Use of Specific Stimulants, by Age Group: 2000 Versus 1999

Stimulants	Total	AGE GROUP (Years)					
		12-17		18-25		26 or Older	
		1999	2000	1999	2000	1999	2000
Methamphetamine, Desoxyn®, or Methedrine	0.1567	0.5994		0.0000		0.4534	
Prescription Diet Pills ¹	0.0870	0.1353		0.0735		0.1124	
Ritalin® or Methylphenidate	0.6600	0.6737		0.8337		0.6190	
Cylert®	0.6061	0.8192		0.8821		0.5707	
Dexedrine®	0.6866	0.0936		0.2656		0.8221	
Dextroamphetamine	0.1997	0.1294		0.7936		0.2421	
Didrex®	0.8035	0.0621		0.8611		0.6993	
Eskatrol®	0.6886	0.4818		0.4263		0.8468	
Ionamin®	0.4670	0.4380		0.2610		0.3427	
Mazanor®	0.7669	0.0185		0.8784		0.9256	
Obedrin-LA®	0.2830	0.6090		0.2555		0.3871	
Plegine®	0.1458	0.2127		0.6038		0.2048	
Preludin®	0.2856	0.5583		0.6421		0.2510	
Sanorex®	0.4563	0.6629		0.8285		0.4982	
Tenuate®	0.6406	0.8173		0.7959		0.5574	

*Low precision; no estimate reported.

¹ For example, Amphetamines, Benzedrine®, Biphetamine®, Fastin®, or Phentermine.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.110C Standard Errors of Estimated Numbers (in Thousands) of Lifetime Users of Specific Sedatives, by Age Group: 1999 and 2000

Sedatives	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Methaqualone, Sopor®, or Quaalude®	309	281	10	8	26	24	308	277
Barbiturates ¹	249	213	7	8	18	18	246	211
Restoril® or Temazepam	85	94	7	7	14	12	83	93
Amytal®	56	49	6	6	4	5	56	48
Butisol®	27	40	3	3	5	9	27	39
Chloral Hydrate	53	45	3	4	6	6	53	45
Dalmane®	56	90	5	5	6	5	56	90
Halcion®	87	78	5	4	12	8	87	78
Phenobarbital	119	115	6	5	14	12	118	113
Placidyl®	96	95	3	4	9	6	96	95
Tuinal®	122	138	3	3	5	6	122	137

*Low precision; no estimate reported.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.110D Standard Errors of Percentages Reporting Lifetime Use of Specific Sedatives, by Age Group: 1999 and 2000

Sedatives	Total		AGE GROUP (Years)					
			12-17		18-25		26 or Older	
	1999	2000	1999	2000	1999	2000	1999	2000
Methaqualone, Sopor®, or Quaalude®	0.14	0.13	0.04	0.03	0.09	0.08	0.18	0.16
Barbiturates ¹	0.11	0.10	0.03	0.03	0.06	0.06	0.15	0.12
Restoril® or Temazepam	0.04	0.04	0.03	0.03	0.05	0.04	0.05	0.05
Amytal®	0.03	0.02	0.02	0.02	0.02	0.02	0.03	0.03
Butisol®	0.01	0.02	0.01	0.01	0.02	0.03	0.02	0.02
Chloral Hydrate	0.02	0.02	0.01	0.02	0.02	0.02	0.03	0.03
Dalmane®	0.03	0.04	0.02	0.02	0.02	0.02	0.03	0.05
Halcion®	0.04	0.04	0.02	0.02	0.04	0.03	0.05	0.05
Phenobarbital	0.05	0.05	0.03	0.02	0.05	0.04	0.07	0.07
Placidyl®	0.04	0.04	0.01	0.02	0.03	0.02	0.06	0.06
Tuinal®	0.06	0.06	0.01	0.01	0.02	0.02	0.07	0.08

*Low precision; no estimate reported.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.

Table 1.110P P-Values from Tests of Differences in Percentages Reporting Lifetime Use of Specific Sedatives, by Age Group: 2000 Versus 1999

Sedatives	Total	AGE GROUP (Years)					
		12-17		18-25		26 or Older	
		1999	2000	1999	2000	1999	2000
Methaqualone, Sopor®, or Quaalude®	0.1313	0.0509		0.2268		0.1814	
Barbiturates ¹	0.6005	0.4189		0.2060		0.6561	
Restoril® or Temazepam	0.8554	0.7067		0.1816		0.9630	
Amytal®	0.8429	0.9513		0.3090		0.9074	
Butisol®	0.3896	0.9385		0.9267		0.3644	
Chloral Hydrate	0.8994	0.2732		0.9267		0.8233	
Dalmane®	0.0623	0.9454		0.8299		0.0575	
Halcion®	0.5668	0.7208		0.0866		0.7326	
Phenobarbital	0.5638	0.9507		0.4104		0.4920	
Placidyl®	0.6409	0.7702		0.1451		0.7207	
Tuinal®	0.3202	0.9105		0.3276		0.3350	

*Low precision; no estimate reported.

¹ For example, Nembutal®, Pentobarbital, Seconal®, Secobarbital, or Butalbital.

Source: SAMHSA, Office of Applied Studies, National Household Survey on Drug Abuse, 1999 and 2000.